Additional Statistical Methodology

Where not available, the HR was estimated using the ratio between probabilities and the 95% CI was estimated using the p-value .

$$HR_{OS}=\frac{ln\ (\pi_{experimental})}{ln\ (\pi_{control})}$$

Calculating the confidence interval for a ratio (log transformation needed)

For example, in the study reported by Witzens-Harig M, the HR was 0.843, p-value was 0.65, we calculate the CI as follows:

z = –0.862+ √[0.743 – 2.404×log(0.65)] = 0.472;

Est = log (0.843) = −0.171;

SE = −0.171/0.472 = −0.362 but we ignore the minus sign, so SE = 0.362, and 1.96×SE = 0.710;

95% CI on log scale = −0.171 − 0.710 to −0.171 + 0.710 = −0.881 to 0.539;

95% CI on natural scale = exp (−0.881) =0.414 to exp (0.539) =1.714.

Hence 95% CI of HR is 0.414 to 1.714.

TableS1. Bayesian network meta-analysis of OS.

enzastaurin	0.96 (0.34, 2.75)	0.9 (0.26, 3.17)	1.12 (0.26, 5.03)	0.38 (0.02, 7.43)	0.72 (0.16, 3.26)	1.02 (0.11, 9.62)
1.04 (0.36, 2.97)	placebo/observation	0.94 (0.49, 1.84)	1.17 (0.42, 3.36)	0.39 (0.02, 6.48)	0.75 (0.25, 2.19)	1.05 (0.14, 7.86)
1.11 (0.32, 3.79)	1.07 (0.54, 2.04)	rituximab	1.26 (0.36, 4.28)	0.42 (0.02, 7.39)	0.8 (0.22, 2.78)	1.13 (0.14, 9.13)
0.89 (0.2, 3.88)	0.85 (0.3, 2.4)	0.8 (0.23, 2.78)	lenalidomide	0.34 (0.02, 4.65)	0.64 (0.14, 2.88)	0.9 (0.1, 8.32)
2.65 (0.13, 50.58)	2.56 (0.15, 41.17)	2.4 (0.14, 41.78)	2.97 (0.21, 41.58)	lenalidomide & rituximab	1.93 (0.1, 36.66)	2.69 (0.09, 82.32)
1.39 (0.31, 6.22)	1.34 (0.46, 3.92)	1.25 (0.36, 4.45)	1.56 (0.35, 6.97)	0.52 (0.03, 10.51)	everolimus	1.41 (0.26, 7.85)
0.98 (0.1, 9.17)	0.95 (0.13, 7.01)	0.89 (0.11, 7.36)	1.11 (0.12, 10.47)	0.37 (0.01, 11.49)	0.71 (0.13, 3.92)	thalidomide

FigureS1. Surface under the cumulative ranking(SUCRA) of the treatments.

