Supplementary Table 1. Demographic information

References	Comparison	Age	Male/female
Sumarriva(2019)	DR cases(n=83)	57.9 ±11.1	52/31
	diabetic controls(n=90)	59.7 ±10.0	47/43
Haines(2018)	rhegmatogenous RD (n=25)	68.0±6.0	-
	PDR(n=9)	62.0±10.0	-
	controls(n=8)	41.0±10.0	-
Zhu(2019)	NDR(n=21)	55(50-58)	9/12
	PDR(n=21)	49(46-56.5)	9/12
Chen(2016)	NPDR(n=40) (discovery set)	59(53-66)	20/20
	diabetes patients without DR(n=40)	62(52-69)	23/17
Rhee(2018)	DR cases(n=32)		12/20
	no DR cases(n=32)		17/15
Wang(2019)	Vitreous samples: PDR(n=28)	49.61(26-65)	12/6
	non-diabetic patients with MH(n=22)	53.95(32-71)	8/14
	AH samples: PDR(n=23)	54.39(35-79)	12/11
	non-diabetic patients with cataract(n=25)	58.39(48-79)	13/12
Paris(2015)	PDR(n=20)		-
	nondiabetic controls(n=31)		-
Jin(2019)	patients with type 2 DM and cataract(n=14)	69.86 ± 9.30	14/-
	patients with DR and cataract(n=13)	61.77 ± 8.39	13/-
	senile cataract patients(n=7)	66.71 ± 12.16	7/-
Barba(2010)	PDR(n=22)	46.1±9.2	12/10

	non-diabetic patients with MH(n=22)	45.3±11.5	14/8
		No difference	No difference

AH=aqueous humor

DR=diabetic retinopathy; PDR= proliferative diabetic retinopathy; NPDR=non-proliferative diabetic retinopathy; MH=macular hole.; DM= diabetes mellitus; RD= retinal detachment