Comparison of faecal protein biomarkers diagnostic accuracy for colorectal advanced neoplasms: a systematic review and meta-analysis

Atefeh Nasir Kansestani[1•], Mohammad Erfan Zare[1•], Qingchao Tong[1], Jun Zhang[1*]

[1] Department of Clinical Laboratory, Sir Run Run Shaw Hospital, Zhejiang University, School of Medicine, Hangzhou, China

• These two authors work equally.

*** Correspondence:** Professor Jun Zhang, Department of Clinical Laboratory, Sir Run Run Shaw Hospital, Zhejiang University, School of Medicine, Hangzhou, China. Tel: (+86 571) 8600 2260, Email:jameszhang2000@zju.edu.cn

Supplemental Table 1. Strategies for database search.

Database	Search Strategy
PubMed http://www.ncbi.nlm.nih.gov/pubmed	**MeSH terms:** ("Colorectal neoplasms") AND ("Diagnosis" OR "Early detection of cancer") **Text words:** ("Colorectal cancer" OR "CRC" OR "Colorectal malignancy" OR "Colorectal tumour" OR "Adenoma" OR "Colorectal neoplasms") AND ("Faecal biomarker" OR "Laboratory tests" "Diagnostic biomarker" OR "Screening Biomarker")
Web of Science http://apps.webofnowledge.com/	**Text words:** ("Colorectal cancer" OR "CRC" OR "Colorectal malignancy" OR "Colorectal tumour" OR "Adenoma" OR "Colorectal neoplasms") AND ("Faecal biomarker" OR "Laboratory tests" "Diagnostic biomarker" OR "Screening Biomarker")
Scopus http://www.scopus.com/	**Text words:** ("Colorectal cancer" OR "CRC" OR "Colorectal malignancy" OR "Colorectal tumour" OR "Adenoma" OR "Colorectal neoplasms") AND ("Faecal biomarker" OR "Laboratory tests" "Diagnostic biomarker" OR "Screening Biomarker")

Supplemental Table 2. Relative-diagnostic Odds Ratio of combined biomarkers compared to individual biomarkers

Biomarker	iFOBT+PK-M2	iFOBT+FC	PK-M2+FC	iFOBT+PK-M2+FC
Colorectal Cancer				
iFOBT	0.54 (0.07-4.02); P= 0.51	0.57 (0.10-2.86); P= 0.46	0.12 (0.01-2.30); P= 0.14	0.11 (0.01-2.07); P= 0.12
PK-M2	2.49 (0.27-22.87); P= 0.40	1.71 (0.33-8.91); P= 0.51	0.95 (0.04-21.56); P= 0.97	0.95 (0.04-22.11); P= 0.97
FC	2.25 (0.53-9.56); P= 0.26	1.75 (0.58-5.27); P= 0.30	0.90 (0.12-7.05); P= 0.91	1.00 (0.13-7.90); P= 0.99
Advance Adenoma				
iFOBT	1.14 (0.06-21.17); P= 0.92	1.47 (0.11-20.34); P= 0.75	0.13 (0.01-3.40); P= 0.18	0.15 (0.01-3.16); P= 0.19
PK-M2	7.91 (0.33-188.14); P= 0.17	4.92 (0.33-72.93); P=0.21	0.72 (0.02-22.43); P= 0.83	0.52 (0.02-14.30); P= 0.66
FC	3.25 (0.42-24.93); P= 0.22	1.61 (0.28-9.27); P= 0.55	0.56 (0.10-3.01); P= 0.44	1.93 (0.33-11.42); P= 0.42
Advance Adenoma				
iFOBT	0.43 (0.08-2.24); P= 0.28	0.59 (0.09-3.76); P= 0.54	0.07 (0.01-0.43); P< 0.01	0.05 (0.01-0.33); P< 0.01
PK-M2	1.31 (0.58-2.97); P= 0.46	3.45 (0.81-14.67); P= 0.86	0.86 (0.28-2.66); P= 0.77	0.85 (0.29-2.47); P= 0.73
FC	4.00 (0.51-31.68); P= 0.16	3.68 (0.57-23.74); P= 0.15	0.45 (0.04-4.68); P= 0.45	0.36 (0.04-3.58); P= 0.33



Supplemental Figure 1. Deeks' funnel plot asymmetry test for CRC biomarkers. **A:** Immunochemical Fecal occult blood tests; **B:** pyruvate kinase-M2; **C:** fecal calprotectin.



Supplemental Figure 2. Deeks' funnel plot asymmetry test for AA biomarkers. **A:** Immunochemical Fecal occult blood tests; **B:** pyruvate kinase-M2; **C:** fecal calprotectin.



Supplemental Figure 3. Deeks' funnel plot asymmetry test for AN biomarkers. **A:** Immunochemical Fecal occult blood tests; **B:** pyruvate kinase-M2; **C:** fecal calprotectin.