Table S4 - Characteristics of studies reporting on treatment statisfaction in acne

#	Study	Design	Treatment	Duration weeks	Setting	Country	Number pts	Age (mean)	Men	Women	Gender unknown	Tool	Comments
1	Abdel Hay 2019[68]	RCT	Azelaic acid (AA) 20% in combination with salicylic acid (SA) 20% vs trichloroacetic acid 25%, 4 sessions	8	Dermatology outpatient clinic, Faculty of Medicine, Cairo University, single-centre	Egypt	34	21,8	3	31	0	Satisfaction questionnaire 4 point Likert scale 0 = dissatisfied - patient feels worse than before/same as before 1 = slightly satisfied - patient feels slightly better but still not worth it 2 = moderately satisfied - patient feels good with need for slight improvement 3 = highly satisfied - patient feels optimal cosmetic result	"Patient's satisfaction evaluation revealed that the patients were more significantly satisfied by the side treated by combined SA and AA peels (P=0.001); the results showed that in the group treated by SA and AA peel, 59% (n=20) were highly satisfied (grade 3), 20% (n=7) were moderately satisfied (grade 2), 15% (n=5) were slightly satisfied (grade 1) and 6% (n=2) were dissatisfied. While in the group treated by TCA peel, 12% (n=4) were highly satisfied, 50% (n=17) were moderately satisfied, 26% (n=9) were slightly satisfied and 12% (n=4) were dissatisfied. Patients were more satisfied by the less downtime and the fewer incidences of brown crusts associated with the combined solution." Note: it says questionnaire, but unclear if it was one or more questions. Seems one question.
2	Adalatkhah 2011[69]	RCT	Systemic flutamide vs cyproteron acetate-ethinyl estradiol	26	Specialty dermatology clinic, single-centre	Iran	64	22	0	64	0	Not reported	"Satisfaction regarding different aspects of drug efficacy and tolerability was not found to be significantly different between the trial groups except for improvement in greasy skin ($P < 0.05$)."

3	Alanazi 2020[70]	Cross-sectional	Any treatment as prescribed by the dermatologist (topical and/or systemic)	N/A	Public areas located in Riyadh	Saudi Arabia	346	22,4	173	173	0	Questionnaire consisting of 15 questions, including question regarding satisfaction with treatment	"The most satisfying medication for the patient was isotretinoin, and 71% were happy with the results of using it compared to only 38% who used topical antibiotics (P = 0.012) [Figure 2]."
4	Alsubeeh 2019[71]	Cross-sectional	Any treatment as prescribed by the dermatologist (topical and/or systemic)	N/A	Outpatient dermatology clinic at King Khalid University Hospital, Riyadh, single-centre	Saudi Arabia	2330	25,3	478	1852	0	Questionnaire (5 sections with questions), the 12-item Medication Adherence Scale was used to quantify the mean adherence score. 5 point Likert scale 1 (never) to 5 (always)(maximum of 60 points) One question on treatement satisfaction.	Five dermatological diseases (psoriasis, chronic dermatitis, acne vulgaris, hair growth disorders, and vitiligo) were included. No separate data for acne. "Psoriasis and chronic dermatitis patients had the lowest mean adherence scores. Patients who were on oral medication had the highest adherence score, while those on topical medication had the lowest score. The thoughtful consideration of factors associated with high adherence is important for optimal therapeutic outcomes." 1563 reported to be satisfied with treatment versus 767 who reported they were not (Table 3).
5	Baker 2001[72]	Open-label	Topical adapalene gel 0.1% QD	12	Dermatology practices (600 dermatologists)	USA	2545	22	572	1335	638	1) 4 point Likert scale Very statisfied, satisfied, moderately satisfied, not satisfied 2) Dichotomous Would continue to use: yes/no.	"Furthermore, 69% of patients were "satisfied" or "very satisfied" with adapalene gel 0.1%, and most patients (80%) said they would continue to use adapalene gel 0.1% (Figure 5)."

6	Barakat 2017[73]	Open-label	IPL, 6 sessions with one week interval	26	Dermatology Out-patient Clinic of Minia University Hospital, Minia, single-centre	Egypt	24	22,4	4	20	0	Cardiff Acne Disability Index (CADI): Questionnaire consisting of 5 questions (score 0-15) Higher scores means more impairment of quality of life	"Regarding CADI score, it was significantly decreased at 2 weeks" and "3 months after IPL with no significant change between them indicating significant patient satisfaction even at 3 months after IPL." Although they reported treatment satisfaction, they actually used a tool to measure quality of life.
7	Baugh 2005[74]	RCT split-face	KTP laser vs no treatment (2 treatments pw for 2 weeks)	6	Single-centre	USA	26	27,8	4	21	0	Nonstandardized overall treatment percent satisfaction scale 0-100%, with intervals of 10 percentage points	The mean overall percentage treatment satisfaction varied from 50-59% to 90-100% with the majority between 60-79% satisfaction with treatment.
8	Brodell 2012[75]	RCT	Topical combined with systemic; adalapene 0.1%-BPO 2.5% QD vs vehicle QD both combined with doxycycline 100 mg for 12 weeks followed by 24 weeks topicals only.	36	Multi-centre	USA/Canada	459	NR	NR	NR	459	Treatment satisfaction questionnaire: - Satisfied with time taken for treatment to work - Satisfied with treatment effectiveness - Feel better about self since starting treatment - Not bothered by side effects - Satisfied overall with treatment - Would use treatment again	No answer categories mentioned. Presented as percentages satisfied. "At weeks 12 and 36, a significantly higher proportion of patients receiving A–BPO vs. vehicle reported high satisfaction for five out of six treatment satisfaction items."
9	Burger 2009[76]	Cross-sectional	Isotretinoin	N/A	N/A: via 30 pharmacies	South Africa	57	20,2	33	24	0	Structured questionnaire consisting of open and closed questions including perceived patient satisfaction or dissatisfaction with isotretinoin	Details not reported. Might be Likert. The response rate was 29.2%, yielding 57 participants. The questionnaire as on "perceived patient satisfaction or dissatisfaction" - "More than 80% of patients felt either very satisfied or satisfied with the perceived efficacy of isotretinoin."

10	Callender 2012[256]	Part of fixed combination of clindamycin phosphate 1.2% and low concentration BPO 2.5% gel study, see for details Gold 2009. This is a post-hoc analysis comparing Skintypes I-III and IV-VI.											
11	Campbell 2006[77]	Open-label	Adapalene gel 0.1 % QD when used with other acne treatments, either as part of an initial combination regimen or as add-on therapy	12	Multi-centre	USA	1979	NR	628	1092	259	5 point Likert scale Very satisfied, somewhat satisfied, indifferent, somewhat dissatisfied, very díssatisfied - verbally given	Data on 1602 patients, subjects without Investigator Global Assessment were not included. Of the 1187 on initial combination therapy 522 (44%) was very satisfied, 491 (41.4%) somewhat satisfied and the remainder (174 = 14.6%) was indifferent to very dissatisfied. Of the 415 who were in the add-on therapy group 164 (39.5%) was very satisfied, 175 (42.2%) was somewhat satisfied and the remainder (76= 8.2%) was indifferent to very dissatisfied
12	Cardinali 2015[78]	CCT	Topical clindamycin 1%/BPO3% QD vs tretinoin 0.05% QD	12	Unità operativa Dermatologia, Prato, single-centre	Italy	40	33,3	0	40	0	Satisfaction questionnaire as Global Tolerability on symptoms: dryness, erythema, exfoliation, itching and cutaneous tension 1 = minimum 2 = mild 3 = moderate 4 = severe	Not actual measuring patient satisfaction
13	Choi 2010[79]	RCT split-face	Cleanser A (papain, proteomax and soap powder) BID vs cleanser B (0.04% triclosan, 1% salicylic acid, and 1% azelaic acid BID in addition to the other constituents of cleanser A)	8	Department of Dermatology, Seoul National University Hospital, Chongno-gu, Seoul	Korea	13	26	6	7	0	NRS 0-10 0 = not satisfied, 10 = very satisfied	"Though patients were generally satisfied with both treatments, they were more satisfied with cleanser B." Estimated scores from figure mean scores were for cleanser A 2.9 vs 4 for cleanser B authors report $p < 0.05$. The degree of satisfaction seems to be low on a scale from 0-10.
14	Cook-Bolden 2006[257]	Part of MORE trial, see for details Campbell 2006. This paper describes subject preferences											

15	Cook-Bolden 2012[258]	Part of fixed combination of clindamycin phosphate 1.2% and low concentration BPO 2.5% gel study, see for details Gold 2009. This paper describes post-hoc data in the Hispanic population											
16	Ctri 2019[80]	RCT	Isotretinoin 20mg QD vs isotretinoin 20 mg and desloratidine 5 mg daily	12	NR	India	86	N/A	N/A	N/A	86	Unclear	Trial protocol and not yet recruiting. Very little information. "Improvement in acne severity assessed by the patient using participants self-assessment of acne severity (very satisfied to unsatisfied)." Probably Likert scale
17	del Rosso 2009[81]	RCT	6% BPO foaming cloth vs 6% BPO cleanser and 6% BPO foaming cloth vs 4% BPO wash	1 session	Two centres	USA	193	NR	69	124	0	17-item survey regarding various attributes of BPO cloth; and an 11-point comparative survey Rating items 5 point Likert scale: 1 (strongly agree) to 5 (strongly disagree)	The 17 items (of which 16 reported) are grouped in convenience with the product (5 items), use attributes (6) and cometic attributes (5). The comparative 11-items survey contained same 5 questions on convenience, use attributes (4) and preference based on cosmetic attributes (2). "These attributes supporting preference for the benzoyl peroxide foaming cloth cleanser may improve patient satisfaction with topical acne treatment resulting in improved patient compliance" This single session study is more about satisfaction with a BPO-based face cleaner, not so much about the treatment as a whole.
18	del Rosso 2018[82]	Open-label	0.3% adapalene and 2.5% benzoyl peroxide gel and oral doxycycline 100 mg BID	12	Multi-centre	USA	186	19,6	78	97	11	4 point Likert scale Very satisfied, satisfied, somewhat satisfied, not satisfied	One question "Are You Satisfied with the Study Treatment?" "A majority of subjects were 'Satisfied' or 'Very Satisfied' with the study treatment overall (83.2%)"

19	Dong 2016[83]	Open-label	Aminolevulinic acid 10% plus photo dynamic therapy - treated once every 30 days for two or three sessions	25	Clinic of the Department of Lab of Laser Surgery, located in the Cosmetic Laser Center of the Ninth People's Hospital, Shanghai JiaoTong University School of Medicine, single-centre	China	46	23	16	30	0	4 point Likert scale Very satisfied, satisfied, somewhat satisfied, not satisfied	"The subjects conducted a self-evaluation of their treatment outcome at the 12-week follow-up visit. Forty-four of the 46 patients (95%) were either satisfied or very satisfied with the improvement in their acne."
20	Donnarumma 2019[84]	RCT	0.3% adapalene and 2.5% benzoyl peroxide gel in 3 groups: G1= training/leaflet; G2= training/leaflet/SMS; G3= standard instructions only	12	Section of Dermatology, Department of Clinical Medicine and Surgery University of Naples Federico II, Naples, single-centre	Italy	126	19,5	39	86	1	Satisfaction questionnaire 4 point Likert scale Very much satisfied, somewhat satisfied, not much satisfied, not at all satisfied	Yet satisfaction results are presented as satisfaction is low, medium-low, medium-high or high. "The patient's satisfaction score after 12 weeks of treatment was higher in G2 and G1 than in G3." Table 4: Satisfaction high G1:32(76.2%); G2:34 (81.0%); G3 25 (61.0%)
21	Dréno 2010[85]	Cross-sectional	Any treatment as prescribed by the dermatologist (topical and/or systemic)	NR	> 1680 dermatologists (16 countries/>=105 per country)	Americas/ Asia/Europe	3751	21,2	NR	NR	3751	Questionnaire 5 point Likert scale Not satisfied, a little satisfied, moderately satisfied, satisfied, very satisfied	Reported graphically. No numbers could be extracted from the graph, and are also not mentioned in the text.
22	Dréno 2018[86]	RCT split-face	0.3% adapalene and 2.5% benzoyl peroxide gel vs vehicle	24	Multi-centre	Canada/France	67	21,5	23	44	0	Satisfaction questionnaire	"According to the subject satisfaction questionnaire at week 24, a total of 90.1% indicated they were satisfied to very satisfied with A0.3/BPO2.5 versus 59.0% for vehicle, and 59.0 versus 32.8% were very satisfied, respectively (p<0.0001). Similarly, 86.8% of subjects were satisfied to very satisfied with the time that A0.3/BPO2.5 took to work versus 50.9% for vehicle (p<0.0001)." Questions seem to be at least 'overall satisfaction with treatment', and 'satisfaction with onset of

													effect'. Answers suggest a Likert-scale, but not reported.
23	Dubois 2019[87]	Open-label	0.3% adapalene and 2.5% benzoyl peroxide gel + prescribed skin care	16	Multi-centre	Mauritius/ Singapore /USA	50	20,9	26	24	0	Satisfaction questionnaire 4 point Likert scale Not satisfied, somewhat satisfied, satisfied, very satisfied	"At week 16, over three-fourths (77%) of subjects were overall satisfied or very satisfied with treatment (from 72% of Asians to 82% of black/African-Americans; Figure 1 B). Three-fourths (approximately 75%) of all subjects were satisfied or very satisfied with treatment effectiveness (from 67% of Latin-Americans to 82% of black/African-Americans)." Besides satisfaction with treatment (graphically reported), also satisfaction with *treatment effect* was asked and reported in the text (not graphically).
24	Eichenfield 2019[88]	RCT post hoc analysis	Tretinoin 0.05% lotion QD vs vehicle QD	12	Multi-centre	USA	154	12,4	63	91	0	Patient satisfaction score (PSS) NRS 1-10 (10 was the most satisfied and a score of 5 or greater was considered as 'satisfied')	"Subject satisfaction with treatment was significantly greater with tretinoin 0.05% lotion than vehicle by Week 12 (P = 0.008)."
25	Fabbrocini 2011[89]	RCT split-face	Resveratrol hydrogel vs vehicle	60 days	Multi-centre	Italy	20	NR	12	8	0	10-point satisfaction scale (1 = lowest to 10 = highest) on: - compliance - smell - texture - results - quality of life	"According to the patients' subjective self-evaluation, using the 10-point scale, the mean satisfaction rate was 8 for compliance, 9 for smell, 8 for texture, 8 for results, and 9 for quality of life."
26	Fernandez-Obregon 2003[259]	Part of BEST study, see for details Weiss 2003. This paper describes efficacy by selected demographic subsets											
27	Franzke 2009[90]	Cross-sectional	N/A	N/A	Questionnaires through 48 pharmacies	Germany	504	21,9	200	302	2	Survey 4 point Likert scale Very unsatisfied, more unsatisfied, more satisfied, very satisfied	"Deficits in care did exist: Thus almost 50% of patients were “more” or “very unsatisfied” with previous acne treatment."

28	Fuchs 2019[91]	Open-label	Absorbing Gold Microparticles and Optical Pulses. 3 treatments (baseline, week 1, week 2) + follow-up at 12 weeks; subresponderders; treatment week 13, 14 and 15 + follow-up week 26 and week 36	36	Multi-centre	Denmark/ Switserland	28	18,9	18	10	0	5 point Likert scale 1 = very satisfied 2 = somewhat satisfied 3 = neither satisfied nor dissatisfied 4 = somewhat dissatisfied 5 = very dissatisfied	"At 12-week follow-up visit 81% of all patients reported that they were either “very satisfied” or “somewhat satisfied” with the treatment. For sub-responders 78% (seven of nine patients) responded “very satisfied” or “somewhat satisfied” with the treatment at 36-week follow-up visit."
29	Gold 2008[92]	Open-label	Photopneumatic technology - 4 treatments at 3-week intervals with follow-up visits at 1 and 3 months	13	NR	USA	11	NR	4	7	0	5 point Likert scale 1 = not satisfied 2 = slightly satisfied 3 = moderately satisfied 4 = very satisfied 5 = extremely satisfied	"Satisfaction (Figure 5) scores increased steadily as the study progressed. Subject satisfaction was slightly but consistendly higher than investigator satisfaction." Numbers are not mentioned in the text. From figure 5 it is estimated the mean subject satisfaction went from about 2.6 to 3.5.
30	Gold 2009[93]	Literature review	Clindamycin phosphate 1.2%/BPO 2.5% gel vs clindamycin phosphate 1.2% gel vs BPO 2.5% gel vs vehicle	12	Multi-centre	USA	2813	19,3	1338	1475	0	NRS (1-10) (1 = least satisfied, 10 = most satisfied)	"At Baseline, subjects randomized to receive clindamycin-BPO 2.5% gel had a mean satisfaction score of 4.2 (dissatisfied) with prior acne therapies. At Week 12, these subjects had a mean score of 7.5 (satisfied), which was statistically significant compared to Baseline ($p<0.001$, Week 12 versus Baseline). Therefore, subjects were significantly more satisfied with clindamycin-BPO 2.5% gel than prior acne therapies (Figure 4). In addition, 81.2 percent of clindamycin-BPO 2.5% gel subjects were satisfied with their treatment at Week 12 (Figure 5)." In the post-

													hoc analysis, subjects with a Baseline score of 5 or less were considered dissatisfied with their prior acne therapy and subjects with a score of 6 to 10 were considered satisfied with their prior acne therapy.
31	Gold 2013[260]	Part of fixed combination of clindamycin phosphate 1.2% and low concentration BPO 2.5% gel study, see for details Gold 2009. This paper describes a post-hoc analysis of efficacy and tolerability											
32	Gold 2014[94]	Open-label	Silk'n Blue Device; eight treatments over a 4-week period. Follow-up visits were at 1 month and 3 months.	17	Tennessee Clinical Research Center in Nashville, TN, single-centre	USA	17	38,8	1	11	5	5 point Likert scale 1 = not satisfied 2 = somewhat satisfied 3 = slightly more satisfied 4 = satisfied 5 = very satisfied	"91.6% of the patients were satisfied (with a rating of 2 or above) and that 75% of the participants reported they were more than satisfied (rating of 3 or above) with the Silk'n Blue device"
33	Gollnick 2015[95]	Open-label	Adapalene 0.1%-BPO 2.5% alone or in combination with other drugs	39	Multi-centre	Germany	5131	18	2899	2196	36	4 point Likert scale Very satisfied, satisfied, hardly satisfied, not satisfied	Overall satisfaction (n= 3920): 43.7% was very satisfied, 41.1% satisfied, 8.5% hardly satisfied, and 6.6% not satisfied. "patient satisfaction was significantly higher in patients with less severe acne."
34	Gonzalez 2012[96]	RCT split-face (2 weeks) + open-label (6 weeks)	Clindamycin 1%/BPO5% vs adapalene 0.1%/BPO2.5%	8	Multi-centre	Argentina	48	27,6	10	38	0	The Product Acceptability and Preference Questionnaire. Satisfaction with treatment: 5 point Likert scale 1 = very satisfied to 5 = very unsatisfied	"Overall treatment satisfaction was high, with 72% of patients rating themselves as "satisfied" or "very satisfied" with C/BPO and 68% with A/BPO at week 1, and corresponding rates at week 2 were 89% with C/BPO and 83% with A/BPO. At week 8, after an additional 6 weeks of full-face C/BPO treatment, 85% of patients rated themselves as "satisfied" or "very satisfied" with treatment."

35	Haedersdal 2008[97]	RCT split-face	Long-pulsed dye laser versus long-pulsed dye laser-assisted photodynamic therapy; 3 treatments in 2 weeks + 4 weeks follow-up + 12 weeks follow-up	14	Department of Dermatology, Bispebjerg Hospital, University of Copenhagen, single-centre	Denmark	15	NR	5	10	0	NRS (0-10) 0 = no satisfaction and 10 = best imaginable satisfaction	"Patient satisfaction was slightly higher with the MAL-LPDL treatment (median [25-75 percentiles] at week 4: 7 [range, 4.75-8]; week 12: 8 [6.25-9]) versus the LPDL treatment (week 4: 6 [3.75-8]; week 12: 7.5 [5-8.75]) at weeks 4 (P = .031) and 12 (P = .034)."
36	Hermes 1998[98]	Open-label	Isotretinoin (start 10mg/day, then increasing; 11 patients starting 40-50 mg/day)	12-120	Department of Dermatology, Charité-Virchow Clinic, Humboldt University, Berlin, single-centre	Germany	94	23,5	76	18	0	4 point Likert scale Very good, good, satisfactory, not satisfactory	"Few patients rated their overall improvement as only satisfactory, and only one male patient failed to respond, also on subsequent treatment with diverse other agents."
37	Ianosi 2013[99]	RCT	Vacuum + IPL (group A) weekly for 5 weeks vs IPL (group V) weekly for 5 weeks vs topical placebo (Sebium H2O Micellaire Solution) (Group 0) daily for 5 weeks	21	Medical Center Dr. Ianosi, Surgical Department, University of Medicine and Pharmacy of Craiova, single-centre	Romania	180	24	37	86	57	Not reported	"Concerning quality of life, patients from Group A were more satisfied with the treatment compared with those belonging to Group V (p=0.004) and significantly more satisfied compared with Group 0 (p< 0.001). Patients from Group V were also statistically significantly more satisfied after the treatment compared with the control group (p<0.001) (Figure 6)." Figure 6 contains no explanatory legend nor data.
38	IRCT2017013132319N1[100]	RCT	Metformin semisolid formulation + azithromycin 250mg every other day vs placebo + azithromycin 250mg every other day	8	Shahid Faghihi Skin and Dermatology Clinic, Shiraz, single-centre	Iran	20	N/A	N/A	N/A	20	NRS (1-10)	Status: recruiting. No data.

39	ISRCTN12892056[101]	RCT	Usual care + spironolactone 50mg/day vs usual care + placebo. Dose escalating to 100 mg.	52	Multi-centre	United Kingdom	434	N/A	N/A	N/A	434	6 point Likert scale	Status: recruiting. No data.
40	Jung 2018[102]	Cross-sectional	Acne patches	N/A	Department of Dermatology, National Medical Center, Seoul	Korea	80	NR	28	52	0	Probably NRS 1-10	English abstract: "Overall, the satisfaction score for the use of acne patches was 5.05" Google translate text: "The average overall satisfaction with the patch is 10 out of 10. It was 5.05 when asked to respond."
41	Kellett 2006[103]	RCT cross-over 4 arms	Over 4 weeks, the patients used each of four topical antibiotics for 1 week: erythromycin/zinc solution, cclindamycin phosphate lotion, benzoyl peroxide (BP)/ erythromycin gel (each applied BID) and clindamycin phosphate gel (applied QD)	4	Inveresk Consumer Research Unit, Edinburgh, single-centre	United Kingdom	67	23	25	42	0	NRS (0-100) Higher is better	"For overall satisfaction with the product, clindamycin phosphate gel received the top score of 67.0, compared with 60.6 for clindamycin phosphate lotion, 57.7 for BP/erythromycin gel (P=0.028) and 56.0 for erythromycin/zinc solution (P=0.029)."
42	Khodaeiani 2012[104]	RCT split-face	2% metronidazol gel (right) vs placebo (left), both BID	8	Department of Dermatology, Sina Hospital, Tabriz University of Medical Sciences, Tabriz, single-centre	Iran	78	22,6	26	44	8	Not reported	"At the last visit, 88% of the patients stated that they were satisfied with the results of treatment on the right side [metronidazol], whereas none reported satisfaction with the treatment results on the left side (P <0.001)."

43	Kim 2008[106]	Open-label split-face	Twice weekly for 4 weeks on the face with the Aurora AC. Immediately after, only one facial half of all the patients treated with the Matrix IR.	13	Department of Dermatology, Yonsei-Zium Skin Laser Clinic, Seoul, single-centre	Korea	10	22,6	8	2	0	Satisfaction survey	Results not reported. Only mention of satisfaction is: "They demonstrated a 75% reduction in active lesion counts in most patients and high patient satisfaction." But that is from their ref 8: Ruiz-Esparza J, Gomez JB. Nonablative radiofrequency for active acne vulgaris: The use of deep dermal heat in the treatment of moderate to severe active acne vulgaris (thermotherapy): A report of 22 patients. Dermatol Surg.2003;29:333–9. Although satisfaction was a predefined outcome, no results were provided.
44	Kim 2018[105]	Open-label	BPO 5% gel QD, liquid cleanser BID and QD moisturizer SPF 30	12	Proinnovera GmbH, Center of Dermatology Excellence, Münster, single-centre	Germany	50	21,8	17	33	0	4 point Likert scale Very satisfied, satisfied, dissatisfied, very dissatisfied	"The majority of subjects (approximately 87%) were satisfied to very satisfied, including 33% very satisfied (Figure 1)." From figure 1: Very satisfied 33%; Satisfied 54%; Dissatisfied 10%; Very dissatisfied 2%.
45	Kwon 2018[107]	RCT split-face	Dual mode vs conventional single pass 1450nm diode laser, 3 sessions	20	Single-centre	Korea	25	23,4	9	16	0	5 point Likert scale 0 = none 1 = slight improvement 2 = moderate improvement 3 = good improvement 4 = excellent improvement	"Patients' mean scores evaluated at the final visit for acne improvement were 2.9 ± 0.8 at the dual-mode side and 2.2 ± 0.6 at the stamp mode-only side, respectively ($P < 0.05$). Improvement for hyperseborrhea and skin texture was also observed at both sides, while degrees of both subjects were better in dual-mode side (2.3 ± 1.1 vs 1.7 ± 1.3 [$P < 0.05$] for seborrhea; 2.2 ± 0.8 vs 1.6 ± 1.1 [$P < 0.05$] for skin texture; Figure 4)." Note: not really

													satisfaction.
46	Kwon 2018[108]	RCT split-face	1450-nm diode laser (DL) vs fractional microneedling radiofrequency, 3 sessions	20	Single-centre	Korea	26	23,4	10	16	0	5 point Likert scale 0 = no improvement 1 = slight improvement 2 = moderate improvemen 3 = good improvement 4 = excellent improvement	"Patients' self-assessments for the degree of acne improvements generally paralleled physicians' evaluation. Patients' mean scores evaluated at the follow-up visit for hyperseborrhoea were 2.1± 0.8 at the FMR side and 2.2±0.9 at the DL side, respectively (P > 0.05). On the other hand, degrees of improvements for acne, skin texture/enlarged pores, and atrophic acne scars were better in FMR side compared with DL side (2.8±1.1 vs. 2.1±1.3 (P < 0.05) for acne; 2.5± 0.8 vs. 1.8±1.1 (P < 0.05) for skin texture/enlarged pores; 2.9 ±1.1 vs. 1.9±1.8 (P < 0.05) for atrophic acne scars) (Fig. 4)." Note: not really satisfaction.
47	Kwon 2019[109]	RCT split-face	1450-nm diode laser (DL) + 450-nm blue light vs 450-nm blue light, 3 sessions	20	Single-centre	Korea	25	21,6	11	14	0	Unclear	"Patient's subjective assessments of satisfaction were also recorded at the final visit". Probably 5 point Likert, but unclear. Only data on improvement are reported not regarding satisfaction.

48	Laquieze 2006[110]	RCT split-face for the moisturizer	Oral isotretinion 10-20 mg + Cetaphil Moisturizing Cream on 1 side of the face vs 20mg or topical tretinoin + Cetaphil Moisturizing Cream on 1 side of the face	2	PERITESCO, Paris, single-centre	France	30	25,4	6	24	0	Satisfaction questionnaire Scores 1-10 (1 = bad, 10 = excellent) 1. Skin comfort is very good or good* 2. The product softens skin* 3. The product firms up skin* 4. The product hydrates skin* 5. The product is not sticky* 6. The product doesn't sting eyes* 7. Product's smell is very pleasant or pleasant* 8. The product provides a soothing sensation during application* 9. Intent to buy † 10. Global score * completely agree or agree † yes or perhaps"	"In both treatment groups, subjects gave Cetaphil Moisturizing Cream a very high rating on all items, and particularly on skin comfort and on softening and hydration properties (Table 2). About 93% of subjects expressed a wish to buy Cetaphil Moisturizing Cream (without considering the price). The mean global score given by the subjects was 7.6 out of 10 and 7.5 out of 10 in the isotretinoin and the tretinoïne group, respectively." Note: scale not given.
49	Lee 2012[111]	Open-label	Selective sebaceous gland electrothermolysis, 3 sessions	52	Department of Dermatology, Chung-Ang University College of Medicine, Seoul, single-centre	Korea	12	24,6	6	6	0	4 point Likert scale 1 = unsatisfied 2 = slightly satisfied 3 = satisfied 4 = very satisfied	"Of the 12 patients, one (8%) reported being ‘‘slightly satisfied’’ with the treatment, four (33%) reported being ‘‘satisfied’’, and seven (59%) reported being ‘‘very satisfied.’’ The mean score for patient satisfaction was 3.50 ± 0.67 (out of 4)."

50	Lee 2013[112]	Open-label	Minimally invasive fractional radiofrequency microneedling	8	Single-centre	Korea	20	26,5	10	10	0	Questionnaire 5 point Likert scale 0= worsened 1 = no improvement 2 = somewhat improved 3 = fairly improved 4 = noticeably improved	"Patients' satisfaction scores for clinical improvement were also assessed at every follow-up visit using a 5-point scale. The mean satisfaction scores for the acne severity were 2.1, 2.0, and 2.0 at the 2-, 4-, and 8-week follow-ups, respectively (Table 2). Those for the facial oiliness were 2.1, 2.2, and 2.1, respectively, at the same follow-up assessments (Table 3). Only one patient graded 0 (worsened) on both criteria. Patients showed relatively high overall satisfaction even though the global improvement scores by physicians diminished at later follow-up"
51	Lekwuttikarn 2017[113]	RCT split-face	595-nm pulse dye laser (PDL) treatment, 2 sessions	8	Department of Pediatrics, Chulalongkorn University, Bangkok, single-centre	Thailand	30	18	17	13	0	NRS (0–10) 0 = dissatisfied, 10 = very satisfied	"The participants were provided a high level of satisfaction with the PDL treatment. The satisfaction scores were 7.53, 8.13, and 8.10 (of 10) at the 2nd, 4th, and 8th weeks, respectively."
52	Lubtikulthum 2019[114]	RCT	Herbal extract gel vs benzoyl peroxide 2.5% gel BID	12	Two centres of Srinakharinwirot University, Bangkok	Thailand	77	21,8	30	47	0	Satisfaction score 7 point Likert scale: −3 = extremely dissatisfied −2 = very dissatisfied −1 = dissatisfied 0 = neither satisfied nor dissatisfied 1 = satisfied 2 = very satisfied 3 = extremely satisfied	"No statistically significant difference between the groups in terms of satisfaction with efficacy was found (P-value = 0.391). However, the median satisfaction with drug administration in the HBE group (2 (25th percentile = 1, 75th percentile = 2)) was higher than in the BP group (1.5 (25th percentile = 0, 75th percentile = 2)), which resulted in a statistically significant difference (P-value = 0.011)" HBE =

													topical herbal extract formula
53	Marron 2013[115]	Observational prospective	Oral isotretinoin up to cumulative dose of 120mg/kg	30	Multi-centre	Spain	346	20,7	143	203	0	Questionaire with three dimension. VAS (0-100)	"The treatment satisfaction assessment, completed at the end of the programme, revealed a significant improvement in clinical symptoms (88.4 ± 12.2), satisfaction regarding the treatment (86.2 ± 11.9) and social life (87.6 ± 15.1). The results are shown in Table III." Note: three dimensions 1) Improvement of symptoms: clinical lesions and greasy skin 2) Treatment: information: efficacy, secondary effects and administration of medication 3) Improvement in social life: carrying out daily activities and emotional state Note: HADS, DLQI and SF-36 were aso measured.
54	Martínez-Carpio 2012[116]	Open-label, 4 arms (per indication, not intervention). Only acne reported here.	Photodynamic Therapy (PDT) in the treatment of actinic keratosis (AK), basal cell carcinoma (BCC), acne vulgaris (AV) and photoaging (PA), in the short and medium term.	52	Clínica Alcolea. Barcelona, single-centre	Spain	20	25,3	9	11	0	5 point Likert scale 1 = very dissatisfied 2 = dissatisfied 3 = somewhat satisfied 4 = satisfied 5 = very satisfied	"2/19 patients declared they were somewhat satisfied 2/19 (11%), 5/19 satisfied (26%) and 12/19 were very satisfied (63%)"
55	Maruguchi 2006[117]	Open-label split-face	1450-nm diode laser vs conventional topical treatment	8	Maruguchi Skin Clinic, Kobe, single-centre	Japan	17	23,3	0	17	0	Questionnaire	"From the assessment of patient satisfaction, seven patients preferred laser treatment, two patients preferred physical treatment and three patients found laser treatment equal to physical treatment." Note: Details of the questionnaire are not mentioned.

56	Mazzetti 2019[118]	RCT	1) 0.1% clascoterone cream BID vs 2) 0.5% clascoterone cream BID vs 3) 1% clascoterone cream QD vs 4) 1% clascoterone cream BID vs 5) vehicle cream QD or BID	12	Multi-centre	Italy	363	19,7	167	196	0	5 point Likert scale 1 = excellent (very satisfied) 2 = good (moderately satisfied) 3 = fair (slightly satisfied) 4 = poor (not satisfied at all)	"Subject satisfaction at week 12/EOS was similar across all five treatment groups. The highest satisfaction score (ie, excellent and good) was reported by subjects treated with clascoterone cream 1% BID (72.6%), followed by clascoterone 0.1% BID (68.3%), clascoterone 1% QD (66.7%), vehicle (64.2%), and clascoterone 0.5% BID (61.4%)."
57	Min 2015[119]	RCT split-face	Fractional microneedling radiofrequency (FMr) vs bipolar microneedling radiofrequency (BR), 2 sessions	12	Department of Dermatology, Seoul National University Hospital, single-centre	Korea	20	22,8	11	9	0	Satisfaction score unclear, probably NRS10 or NRS11, higher is better	"Satisfaction score was higher in BR than in FMR right after treatment and at day 1 (FMR: 7.6 ± 0.72, BR: 8.46 ± 0.42). However, higher score was marked in FMR than in BR at day 7 and 84 (FMR: 7.41 ± 0.59, BR: 6.35 ± 0.73, P = 0.004)."
58	Min 2016[120]	Retrospective analysis	Fractional microneedling radiofrequency (FM) vs topical therapies and/or oral antibiotics	N/A	Department of Dermatology, Seoul National University Hospital, single-centre	Korea	52	20,8	31	21	0	VAS (0-10) 0 = dissatisfied, 10 = fully satisfied	"The satisfaction score and the therapeutic effectiveness for the FMR treatment were 7.8 ± 0.3 and 3.6 ± 0.3, respectively. All subjects scored greater than 5 for satisfaction."

59	Moftah 2016[121]	RCT split-back	IPL vs PDT, 3 sessions	15	Outpatient Clinics of the Dermatology and Venereology Department, Al-Zahraa University Hospital and Cairo Hospital, Caïro	Egypt	35	23,7	21	14	0	Cardiff Acne Disability Index (CADI): Questionnaire consisting of 5 questions (score 0-15) Higher scores means more impairment of quality of life	"There is significant decrease in CADI score 1 and 3 months after treatment (5.88 ± 2.84, 6.42 ± 3.66) compared with that before treatment (11.28 ± 3.07) indicating a significant patient satisfaction with the outcome of treatment on both sides (P < 0.005). The patients' satisfaction (CADI score) after treatment (1 month after the last session) showed a significant positive correlation with the percentage of improvement in total lesions count (r = 0.661, P < 0.005)." Although they reported treatment satisfaction, they actually used a tool to measure quality of life.
60	Mohanan 2012[122]	Case-series	IPL up to 5 sessions	72	Department of Dermatology, Madras Medical College, Chennai, Tamil Nadu, single-centre	India	10	NR	NR	NR	10	Unclear	"Patients were (…) asked to grade their satisfaction levels after all the treatment sessions were completed." No tool reported. "The overall patient satisfaction with IPL was 87.5% in our study."

61	Mokhtari 2016[123]	RCT	Azithromycin gel 2% BID vs clindamycin gel 1% BID	16	Skin Diseases and Leishmaniasis Research Center, Department of Dermatology, Isfahan University of Medical Sciences, Isfahan, single-centre	Iran	40	21,8	7	33	0	VAS (0-10)	"Average satisfaction score in patients 4 weeks after treatment in clindamycin gel 1% group and azithromycin gel 2% was 6.98 ± 1.4 and 6.88 ± 0.99, and no significant difference was observed between the two groups according to t-test results (P = 0.71). Eight weeks after treatment, average satisfaction score was 7.5 ± 1.4 and 7.3 ± 1.24 in two groups, respectively, and the difference between two groups was not significant (P = 0.51). Two months after treatment, average of this score was 7.7 ± 1.1 and 7.8 ± 1.1 in two groups, respectively and no significant difference was observed between them (P = 0.8). Variance analysis with replication of observations showed also no significant difference in two groups (P = 0.6)."
62	Mokhtari 2017[124]	RCT	Benzoyl peroxide 5% QD vs benzoyl peroxide 5% QD + IPL, 3 sessions	16	Skin Diseases and Leishmaniasis Research Center, Department of Dermatology, Isfahan University of Medical Sciences, Isfahan, single-centre	Iran	72	25,6	29	29	14	VAS (0-10) 0 = dissatisfied, 10 = very satisfied	"Patients in IPL groups were more satisfied than control group (P < 0.001)." Details reported by graph only, not data.

63	Monfrecola 2016[125]	Open-label	Face compact cream (FCC) QD for cosmetic camouflage	4	Department of Clinical Medicine and Surgery, Section of Dermatology, University of Naples Federico II, Naples, single-centre	Italy	20	33,4	0	20	0	Questionnaire 6 questions: 1) Is the FCC effective? 2) Did the FCC improve your skin? 3) Did the FCC help you to be happier by giving you a more even complexion? 4) Is the FCC pleasant? 5) Are you satisfied with the FCC? 6) Would you continue to use the FCC?" 5 point Likert scale Disagree, neither agree no disagree, agree, strongly agree	"The patients in this study observed that the FCC led to an improvement in their acne. Before application, the mean VAS score for the patients' rating of their facial skin was 4.7. After 28 days of use, there was a 64% improvement in the mean VAS score for the patients' rating of their facial skin, which was reduced to 1.7 (Figure 3A). The patients also provided a positive opinion on the FCC (Figure 3B): 90% agreed that the cream was effective; 80% agreed that the cream improved their skin; 95% agreed that the cream helped them to be happier by providing a more even complexion; and 100% agreed that the cream is pleasant. Furthermore, 100% of patients reported that they were satisfied with the FCC and 95% stated that they would continue to use the cream."

64	Myhill 2017[126]	RCT	Adapalene 0.1%/benzoyl peroxide 2.5% gel A/BPO) + supplementary patient eduction (SEM) vs A/BPO and standard of care (SCOPE) vs A/BPO and extra vists.	12	Multi-centre	United Kingdom	97	22,5	30	67	0	Questionnaire	"According to the subject appreciation questionnaire, the SEM was more helpful to adhere to the treatment (56.7%) versus more visits (32.3%) and A/BPO alone (15.2%), better use the product (70%, 61.3%, and 54.5%, respectively), and better manage skin irritation (53.3%, 48.4%, and 36.4%, respectively) (Fig. 4)." "Fig. 4 Subject appreciation questionnaire: the information you have received in the study about the treatment helps you to a) better manage skin irritation? b) adhere (or be compliant) to the treatment? c) understand when you should expect to see an mprovement of your acne? and d) understand the correct way to use the study treatment?" Note: data given in percentages, no scales.
65	NCT01082523[127]	RCT	Clindamycin/benzoyl peroxide gel in the mornings and adapalene gel in the evenings vs same + text message on BID use	12	Multi-centre	USA	40	22,7	15	25	0	Dermatology Life Quality Index (DLQI), and patient satisfaction surveys	Note: Study published as Boker 2012. No details given on tool nor any results reported on satisfaction. DLQI is not measuring satisfaction although the report states this.

66	Ohlsen 2019[128]	Open-label	Topical clindamycin phosphate 1.2% and tretinoin 0.025% (Clin-RA) QD	12	Multi-centre	Sweden	84	22,6	17	67	0	4 point Likert scale Very satisfied, satisfied, dissatisfied, or very dissatisfied	"Patient satisfaction data were available for 50 patients at week 12. The majority of patients (n = 35; 70%) were satisfied or very satisfied with Clin-RA treatment, while 15 (30%) patients were dissatisfied or very dissatisfied."
67	Paradisi 2011[129]	Retrospective analysis	Flutamide (Flu) vs flutamide + oral contraceptive Ginoden (Flu+OC)	156	Department of Obstetrics and Gynecology and Reproductive Biology, S. Orsola Hospital, University Alma Mater Studiorum of Bologna, Bologna, single-centre	Italy	230	21,9	0	230	0	3 point Likert scale Highly satisfied, satisfied, or dissatisfied	"Patients' self-evaluations of the clinical outcome of the treatment were also consistent with changes in the acne and seborrhea scores and indicated practically a plebiscitary satisfaction with more than 97% of the subjects satisfied or highly satisfied after the first year of treatment reaching 100% at the third year (Table I). All the patients continued to express complete satisfaction also in the following years of treatment (data not shown due to the striking dropout of subjects)."
68	Pariser 2014[130]	RCT	Clindamycin phosphate 1.2 % and benzoyl peroxide 3.75% QD vs vehicle QD	12	Multi-centre	USA	498	18,7	256	242	0	Patient satisfaction score (PSS)(1-10) 10 = the most satisfied	At baseline, patients in both treatment groups also had similar mean PSS scores with their prior facial acne treatment (4.8 and 4.4, respectively). By week 12, the mean PSS score for patients treated with clindamycin-BP 3.75% had increased to 7.5 (a 56% increase) compared to 6.2 with vehicle."

69	Picosse 2016[131]	Open-label	Isotretinoin 0.5mg/kg up to 120mg/kg	52	Two centres	Brazil	50	20	35	15	0	Not reported	"There were 3 premature withdrawals due to personal reasons (not for adverse events), meaning that 47 patients completed the treatment, with 41 (87.2%) becoming very satisfied and 6 (12.8%) satisfied with the outcome." Note: scale not reported.
70	Politi 2015[132]	Open-label	Cooling-vacuum-assisted erbium:glass (Er:glass) 1540-nm laser, 4-6 sessions	18	Single-centre	Israel	12	19	7	5	0	5 point Likert scale 1 = not satisfied, 5 = very satisfied	"Patients' self-rated satisfaction was ranged between "satisfied" (four patients) to "very satisfied" (eight patients)—at the final study visit."
71	Prilepskaya 2003[133]	Open-label	Desogestrel (50/100/150µg) and ethinylestradiol (35/30/30µg) for three phases of 7 days followed by a 7-day pill-free interval, for 6 cycles	12	Scientific Center of Obstetrics, Gynecology and Perinatology, Russian Academy of Medical Science, Moscow and Moscow State Medico-Stomatological University, Moscow	Russia	60	23,1	0	60	0	5 point Likert scale Very satisfied, satified, somewhat satisfied, not satisfied, very dissatisfied	"Subjective assessments indicated that after one, three and six cycles, 69%, 93% and 98%, respectively, of women were satisfied or very satisfied with the DSG-OC." Tool actually not reported. Note: figure 4 shows categories very satisfied, satified, somewhat satisfied, not satisfied, very dissatisfied.
72	Rapp 2004[134]	Cross-sectional	All treatments	N/A	Respondents were recruited from a high school, two universities, an academic medical centre dermatology Clinic and through a database of persons with acne maintained by Johnson & Johnson	USA	479	23,3	172	307	0	Questionnaire on four dimensions: how well it works, cost, ease of use and time spent using it 6 point Likert scale 1 = extremely dissatisfied, 6 = extremely satisfied A sum of the ratings was used with possible scores ranging from 4 to 24	Mean satisfaction was 15.86 (out of a score of 4-24)(Cronbach's alfa was 0.78). Satisfaction scores were further reported as coefficients per predictor (age, sex, ethnicity, income adequacy, education, global health, acne severity and trait anger). The questions are not completely reported.

73	Rassai 2013[135]	RCT	Azithromycin 500 mg per day, 3 days a week plus oral levamisole 150 mg per day, 2 days a week vs azithromycin 500 mg per day, 3 days a week	8	Outpatient dermatology clinic, Department of Jundishapur University of medical sciences, Ahvazm, single-centre	Iran	169	NR	60	88	21	4 point Likert scale Weak, intermediate, good, excellent	In Group 1 satisfaction rates were medium" in 4 patients, good in 54 and excellent in 16 patients. In Group 2 these numbers were 27, 45 and 2, respectively. Totally regarding treatment results all patients in Group 1 were more satisfied than Group 2 and this differences were more significant in excellent group ($p < 0.05$)."
74	Rendon 2015[136]	Cross-sectional	N/A	N/A	Web-based observational survey was conducted with subjects in the US	USA	312	35,3	0	312	0	Survey 5 point Likert scale Strongly disagree, disagree, neutral, agree, strongly agree	No details on the tool are provided, but becomes clear from Table 3 "among subjects currently using an acne treatment, fewer than half “agreed” or “strongly agreed” that they were satisfied with their treatment (OTC BP, 47.0% [47/100]; OTC SA, 43.0% [46/107]; Table 3). The most frequently reported reasons for not being satisfied with BP or SA were skin dryness (BP, 26.3%; SA, 44.3%) and flakiness (BP, 12.3%; SA, 31.1%); bleaching of clothes or bedding with BP (8.8%), and “cakey” make-up appearance with SA (19.7%)." BP = benzoyl peroxide, SA = salicylic acid
75	Rist 2003[261]	Part of BEST study, see for details Weiss 2003. This paper describes the study design and the selection criteria											
76	Rodriguez 2003[262]	Part of BEST study, see for details Weiss 2003. This paper describes results according to prior treatment											
77	See 2018[137]	Review	Topical retinoids	N/A	N/A	Australia/ Singapore /Japan/ Korea/Philippines	N/A	N/A	N/A	N/A	N/A	N/A	Review summarizes the results of impact of supplemental patient education on satisfaction with retinoid therapy of the study of Myhill 2017 (row 59)

78	Shamban 2008[138]	Retrospective analysis	Photopneumatic device, that delivers broadband light (400 to 1200 nm), 2-4 sessions	N/A	Multi-centre	Japan/USA	56	25	5	51	0	4 point Likert scale 1 = not satisfied 2 = somewhat satisfied 3 = satisfied 4 = very satisfied	No details on the tool are provided. Legend of Figure 3 states "Overall satisfaction as judged by treating physicians and patients. Both data sets show an increase in satisfaction with number of treatments" From figure 3 it looks that after 1 treatment the overall satisfaction score was 3 (satisfied) and after 4 treatments 4 (very satisfied)
79	Shavin 2003[263]	Part of BEST study, see for details Weiss 2003. This paper describes implications of the BEST study											
80	Shin 2012[139]	RCT split-face	CO2 laser vs fractional microneedle radiofrequency, 1-2 sessions	12	Department of Dermatology, Severance Hospital, Cutaneous Biology Research Institute, Yonsei University College of Medicine, Seoul, single-centre	Korea	20	23,5	11	9	0	Survey, 4 point Likert scale 0 = unsatisfied 1 = slightly satisfied 2 = satisfied 3 = very satisfied	"After MRF device treatment, surveys evaluating overall patient satisfaction revealed that three of 20 patients (15%) were very satisfied, eleven (55%) were satisfied, five (10%) were slightly satisfied, and one (5%) was unsatisfied (Table I and Figure 5). After CO2 FS treatment, two of 20 patients (10%) were very satisfied, eight (40%) were satisfied, seven (35%) were slightly satisfied and three (15%) were unsatisfied. The overall satisfaction levels of MRF device and CO2 FS were not significantly different (p = 0.435)."

81	Smith 2006[140]	RCT	Benzoyl peroxide 5,5% microsphere cream QD vs benzoyl peroxide 6% gel	12	Division of Dermatology, University of California, San Diego and Department of Dermatology, University of Minnesota Medical School Minneapolis	USA	48	17,1	26	22	0	Survey questionnaire	No details were provided regarding the tool, most likely a Likert scale. "A similar number of subjects from both treatment groups rated their overall satisfaction as good or excellent (67% in the 5.5% BP-MS cream group and 68% in the 6% BP gel group). However, more subjects in the 5.5% BP-MS cream group rated their overall satisfaction as excellent compared with the 6% BP gel group (38% vs 11%, respectively)(Figure 4)."
82	Son 2010[141]	RCT	*Ah shi point* acupuncture vs general acupuncture, 12 sessions	6	Department of Oriental Dermatology, Kyung Hee University, East-West Neo Medical Center, Seoul	Korea	36	26,1	10	16	10	5 point Likert scale 1 = dissatisfied to 5 = very satisfied	"The mean final satisfaction scores were 4.0±0.6 for the acupuncture group and 3.5±0.9 for the acupuncture + *ah shi* point group, which are not significantly different (p=0.098)."

83	de Souza Sittart 2015[142]	Open-label	Adapalene 0.1% and benzoyl peroxide 2.5% gel QD	12	Multi-centre	Brazil	79	18,3	47	26	6	Satisfaction questionnaire: - “were you uncomfortable with the adverse events of the treatment?” - “was it easy to incorporate the treatment regimen in your daily life” - "compared with other treatments for acne used this treatment was a litte worse to much better?" - "would like to use again"	The satisfaction questionnaire showed to the question, “were you uncomfortable with the adverse events of the treatment?”: 38 patients (52.1%) answered “not at all uncomfortable”, 34 (46.6%) answered “a little uncomfortable” and 1 (1.4 %) answered “uncomfortable”. - To the question, “was it easy to incorporate the treatment regimen in your daily life”: 43 patients (58.9%) strongly agreed, 28 (38.4%) agreed and 2 (2.7%) disagreed. - Compared with other treatments for acne used by patients, 52 (75.4%) found the study treatment “much better”, 13 (18.8%) “a little better”, 3 (4.3%) “equal”, and 1 (1.4%) “a little worse”. - 69 patients (94.5%) stated they would like to use this treatment again and 4 (5.5%) declared they would not like." and "Of the 73 patients, 69 (94.5%) were satisfied or very satisfied with the treatment. Only 4 participants (5.5%) declared to feel a little satisfied or dissatisfied". The last sentence suggest there was also a Likert scale, however this is not further mentioned in the reference. "Considering the patient satisfaction questionnaire (...) most patients reported being satisfied with the results"

84	Stein-Gold 2016[143]	RCT	Adalapene 0.3%/benzoyl peroxide 2.5% gel QD vs adalapene 0.1%/benzoylperoxide 2.5% QD vs vehicle QD	12	Multi-centre	Canada/USA	503	19,3	250	253	0	Questionnaire	No details regarding the questionnaire were provided. "Regarding the subject's satisfaction questionnaire, more than twice as many subjects in the 0.3 % A/BPO group were satisfied with the product than in the vehicle group (71.6 vs. 32.3 %, respectively). Over half of subjects (about 53 %) considered that 0.3 % A/BPO did not lead to skin irritation compared with 76.9 % for vehicle."
85	Suh 2008[144]	Cross-sectional	N/A	N/A	Multi-centre	Korea	1400	23,8	480	756	164	Questionnaire	No details regarding the questionnaire were provided. "Seventy-seven percent of patients were unsatisfied with the results of acne treatment. The major reasons for dissatisfaction were ineffectiveness (84%), cost (7%), and side effects (4%). The average total cost of acne treatment was higher in dissatisfied patients than in satisfied patients (1,155 vs. 699 US dollar, $p<0.01$) (Fig. 5). Moreover, the forehead was the first presenting site in 71% of satisfied patients and in 62% of dissatisfied patients ($p<0.05$). On the other hand, cheeks were more often the first presenting area in dissatisfied patients (61% vs. 51%) ($p<0.05$). When completing the questionnaire, 14% of dissatisfied and 9% of satisfied patients had chest lesions ($p<0.05$). Seventy-seven percent of satisfied patients and

													87% of dissatisfied patients thought that emotional stress aggravates acne ($p<0.01$)."
86	Tan 2012[145]	Two RCTs, second is follow-up study of the first	1. Doxycycline 100mg per day + adapalene 0.1%/benzoylperoxide 2.5% (A/BPO) vs doxycycline 100mg per day + vehicle for first 12 weeks 2. In case of at least 50% improvement re-randomized in to A/BPO or vehicle for an additional 24 weeks	36	Multi-centre	Canada/USA	243	18,5	132	111	0	Satisfaction questionnaire	No details regarding the questionnaire were provided. Probably Likert scale (Figure 4) "At week 36, the highest level of overall treatment satisfaction and satisfaction with treatment effectiveness was for (A/BPO + DCN) + A/BPO, which was significantly greater than the regimens of (A/BPO DCN) + vehicle and (vehicle + DCN) + vehicle (both $P < 0.05$).....Finally, 85.0% of subjects in the group of (A/BPO + DCN) + A/BPO were "very satisfied" or "satisfied" with the overall treatment scheme, the highest among four treatment regimens (85.0% vs 52.4-75.5%)

87	Tan 2018[146]	RCT	Adapalene 0.1%/benzoylperoxide 2.5% (A/BPO) overnight (A/BP-EN) vs A/BPO for 3 h (A/BPO-3h) vs A/BPO overnight + daily moisturizer (A/BPO-moisturizer) vs A/BPO every other night (A/BPO-EoN)	4	Multi-centre	Canada	123	20,6	52	71	0	Questionnaire	No details regarding the questionnaire were provided. Probably Likert scale (Figure 5) "A high percentage of subjects reported to be “very satisfied” or “satisfied” with treatment effectiveness at Week 4. A very high percentage of subjects were “very satisfied” or “satisfied” with the treatment and its instructions for use at both Weeks 4 and 12. A relatively lower percentage of subjects were satisfied with the A/BPO-moisturizer regime at Week 12, compared with the other three groups, while the responses at Week 4 were similarly positive for all four groups."
88	Tanghetti 2019[147]	RCT	Tazarotene 0.045% lotion QD vs tazarotene 0.1% cream QD vs vehicle GD	12	Multi-centre	USA	210	22,2	94	106	0	Patient satisfaction survey NRS 1-10 (10 = most satisfied)	"There were no statistically significant differences in PSS mean scores at week 12 between tazarotene 0.045% lotion and tazarotene 0.1% cream (P = 0.372) or combined vehicle (P = 0.242). Overall, patients treated with tazrotene 0.045% lotion assessed their treatment satisfaction higher than tazarotene 0.1% cream (mean score of 7.7 versus 7.4)." As Minimal Clinical Important Difference (MCID) is not known a difference of 0.3 might not be clinically meaningful.

89	Ting 2012[148]	RCT split-face	Clindamycin phosphate 1.2%/benzoyl peroxide (BPO) 2.5% gel QD vs adalapene 0.1%/BPO 2.5% gel QD	2	California Dermatology Care, San Ramon, single-centre	USA	21	21,3	11	10	0	Satisfaction questionnaire -Which side of your face did the medication absorb or dry more quickly? -Which side of your face feels more moisturized or hydrated? -Which side of your face stings or burns more? -Immediately following product application, which side of your face do you prefer from an overall appearance standpoint? -Which side of your face feels smoother or softer? -Which product (side of your face) do you prefer for facial application overall? -Which product are you more satisfied with in relation to the improvement of your acne? -Which product (side of your face) are you most likely to continue using in the future?	Dichotomous questions on preference of the split-face two treatments. "After 2 weeks of treatment, results of the participant satisfaction questionnaire showed that more participants preferred clindamycin phosphate 1.2%–BPO 2.5% gel than adapalene 0.1%–BPO 2.5% gel for overall facial application and from an overall appearance standpoint. Participants rated that clindamycin phosphate 1.2%–BPO 2.5% gel was absorbed or dried more quickly, moisturized or hydrated better, stung or burned less, and left the face smoother or softer (Table 2). More participants were satisfied with the improvement of their acne after 2 weeks of treatment with clindamycin phosphate 1.2%–BPO 2.5% gel and nearly twice as many participants were likely to continue using clindamycin phosphate 1.2%– BPO 2.5% gel (65% [13/20]) compared with adapalene 0.1%–BPO 2.5% gel (35% [7/20])(Table 2)."
90	Tremblay 2006[149]	Open-label	Narrowband 415nm light-emitting diode blue light, 2 sessions per week	4-8	Three different centres in southern California	USA	45	26	14	31	0	Questionnaire	No details were provided regarding the tool. "Of the patients surveyed, 10% were not satisfied, 40% were mildly to moderately satisfied, and 50% were highly satisfied with the treatment and results. A significant number of patients reported a decrease in skin oiliness or seborrhea, as well as a perceived decrease in

													pore size."
91	Troielli 2016[150]	Open-label	Adalapene 0.1%/benzoyl peroxide (BPO) 2.5% gel QD	12	Multi-centre	Argentina	105	19,8	28	63	14	Satisfaction survey 5 questions about the therapeutic results in terms of efficacy, tolerability, speed of the therapeutic response, and adherence Each variable was rated on 5 point Likert scale: Excellent, very good, good, regular, or bad	No precise data were provided. "Overall patient satisfaction was similar to the effectiveness rating, with 94% of patients indicating that they were very satisfied or satisfied with treatment. The majority (81%) felt that adapalene/BPO treatment met their expectations, and 95% felt the regimen was easy to follow."
92	Tucker 2017[151]	Cross-sectional	N/A	N/A	N/A 9 pharmacies	United Kingdom	40	NR	NR	NR	40	Questionnaire	Only 4 patients had acne, the remainder had dermatitis. Satisfaction with pharmacy-based management. "When asked, 52.2% (12/23) of participants stated that they were 'very satisfied' with the advice and treatment they received at the pharmacy, 39.1% (9/23) were 'satisfied', but 8.7% (2/23) reported that they were 'very dissatisfied'."
93	Tyring 2018[152]	Two RCTs	Tretinoin 0.05% lotion QD vs vehicle QD	12	Multi-centre	USA/Dominica Republic/ Honduras/ El Salvador	1640	20,5	731	909	0	Patient satisfaction score (PSS). NRS 1-10; 10 was the most satisfied and a score of 5 or greater was considered as 'satisfied', a score of 5 or greater was considered as 'satisfied'	"Patient satisfaction (as assessed by PSS) was significantly greater with tretinoin 0.05% lotion by week 12. At baseline, both patient groups had a similar satisfaction with prior facial acne treatment (4.7 and 4.6 respectively). By week 12, the mean PSS for patients trfeated with tretinoin 0.05% lotion had increased by 53% (compared with 43% with vehicle; $P < 0.01$) and 9 out of 10 patients reported to be satisfied with their treatment."

94	Veraldi 2012[153]	Open-label	Retinoid treatment + Myrtacine® 0.2% and vitamin PP (pellagra preventing) 4% BID vs retinoid treatment + emollient cream BID	4	Multi-centre	Italy	167	22,8	69	98	0	Not reported	No data provided. "0.2% Myrtacine® + 4% vitamin PP provided greater patient satisfaction and acceptability (data not shown)".
95	Veraldi 2016[154]	Cross-sectional	Cleansers	N/A	N/A survey	Italy	786	23,9	248	538	0	Survey consisting of 10 question of which one was "Are you satisfied about this cleanser?"	"The fourth question was: ‘‘Are you satisfied about this cleanser?’’: 338 patients answered ‘‘yes’’ (62.2%), 200 patients answered ‘‘no’’ (36.8%) and 5 patients answered ‘‘I do not know’’ (0.9%)."
96	Veraldi 2016b[155]	RCT	Hydrogen peroxide-based kit (cleanser, day and night treatment) vs benzoyl peroxide based kit (cleanser, toner and repair treatment)	8	Two centres in Catania and Rome	Italy	120	NR	NR	NR	120	Patient self-satisfaction questionnaire	Althought the authors state in the abstract "at Week 8, a patient self-satisfaction questionnaire was administered", there is no mentioning of this at all-in the methods, nor the results.
97	Weiss 2002[265]	Part of BEST study, see for details Weiss 2003. This paper describes preliminary results regarding patient satisfaction											

98	Weiss 2003[156]	Open label BEST study; BenzaClin ® (benzoyl peroxide/clindamycin topical gel) Efficacy and Satisfaction Trial (BEST)	Benzoyl peroxide/clindamycine gel BID	8	Multi-centre	USA	1669	19,6	681	988	0	11 point VAS 0 = not satisfied, 10 = very satisfied	"At baseline, the mean patient satisfaction score in 1389 patients was 2.6 ± 1.3, in dicating a dissatisfaction with their current acne treatment regimens." and "Following 8 weeks of treatment with benzoyl peroxide/clindamycin topical gel, patient satisfaction scores improved more than 250% from baseline (Figure 1). This improvemen was significant compared with baseline levels (P≤0.0001)." Figure 1 shows mean score of 7.4
99	Weiss 2003b[264]	Part of BEST study, see for details Weiss 2003											
100	Wheeland 2011[157]	Open-label	Blue light BID	8	NR	USA	33	NR	NR	NR	33	5 point Likert scale Not satisfied, slightly satisfied, satisfied, very satisfied, extremely satisfied	"Overall, 82 percent were satisfied, very satisfied , or extremely satisfied with the blue light treatment regimen."
101	Yu 2011[158]	Open-label	TriPollar radio-frequency, 6 sessions	6	Skin and Cancer Foundation Inc., Manila	Phillipines	20	NR	NR	NR	20	5 point Likert scale 0 = none 1 = mild improvement 2 = good 3 = very good 4 = excellent	Degree of improvement is not the same as satisfaction. "Average improvement as rated by patients at the 4-week follow-up visit was 2.5 on the rating scale, indicating good to very good satisfaction with the clinical results."

A = adapalene, BID = twice daily, BP or BPO = benzoyl peroxide, CCT = controlled clinical trial, DL = diode laser, IPL = intense pulse light therapy, KTP = potassium titanyl phosphate, LPDL = long pulsed dye laser, MAL = methylaminolevulinic acid, N/A = not applicable, NR = not reported, NRS = numeric rating scale, OC = oral contraceptive, OTC = over-the-counter, PDL = pulsed dye laser, PDT = photodynamic therapy, PSS = patient satisfaction score, pw = per week, QD = once daily, RCT = randomised controlled trial, SA = salicylic acid, SPF = sun protection factor, VAS = visual analogue scale, vs = versus