Table S5 - Characteristics of studies reporting on treatment statisfaction in acne scars

#	Study	Design	Treatment	Duration weeks	Setting	Country	Number pts	Age (mean)	Men	Women	Gender unknown	Tool	Comments
1	Abdel Kareem 2020[159]	CCT split-face	Combined subcision using carboxytherapy plus fractional carbon dioxide (CO2) laser resurfacing vs only fractional CO2 laser, 3 sessions	22	Department of Dermatology and Venereology, Faculty of Medicine, Al-Azhar University, Cairo, single-centre	Egypt	20	28,4	0	0	20	Patient satisfaction was also evaluated for the degree of improvement on both sides of the face	No details on the tool/scale were provided "For patient satisfaction, we found that 20% of the patients were highly satisfied, 35% of the patients were moderatelyvsatisfied, 25% showed mild satisfaction and 20% were not satisfied after treatment for the right side, while on the left side there was no high satisfaction and only 20% of the patients shows moderate satisfaction, 35% showed mild satisfaction and 45% showed no satisfaction as regarding the left side, so the satisfaction is higher in the right than the left side but does not reach statistical difference." The rights side was the side treated with the combination of treatments
2	Afra 2018[160]	RCT	Topical tazarotene gel 0.1% QD vs microneedling therapy (4 sessions)	26	Dermatology outpatient department clinic of a tertiary care institute, single-centre	India	36	23,4	13	23	0	Patient Global Assessment NRS 0 (no improvement) to 10 (maximum improvement) poor (0-3), good (4-7), and excellent (8-10)	"A significant difference was observed in the comparison of the PGA scores (maximum of 10) of both treatment groups at the 3-month follow-up(mean[SD] microneedling, 5.96 [1.96] vs tazarotene, 2.14 [0.59]; mean difference, −3.82; 95% CI, −4.32 to −3.32; P < .001) and 6-month follow-up (mean [SD] microneedling, 5.86 [2.77] vs tazarotene, 5.76 [2.23]; mean difference,−0.97; 95%CI, −1.67 to −0.27; P < .001). The independent

													dermatologist scores did not show a significant difference between the treatment groups at the 3- and 6-month follow-up visits (median [interquartile range], 5.5 [2.0-7.0] vs 4.5 [2.2-7.0]; estimated median difference assessed with the Hodges-Lehmann-Sen statistic, 0.0; 95% CI, −1.0 to 0.5; P = .67)."
3	Alajlan 2011[161]	Retrospective analysis	1,550nm erbium-doped fiber fractional laser (average 5 sessions) vs fractional 10,600nm CO2 laser (average 2.5 sessions)	52	Dermatology department, College of Medicine, King Saud University, Riyadh, single-centre	Saudi Arabia	82	NR	NR	NR	82	4 point Likert scale Mild = < 25% improvement Moderate = 26–50% improvement Good = 51–75% improvement Very good = > 75% improvement	"An overall patients' satisfaction was 71% for NAF 1,550 nm laser group and 65% for AF CO2 laser group"
4	Al-Dhalimi 2015[162]	Open-label	Fractional 1,540nm erbium glass laser, 3 sessions	12	Laser Research Unit, Faculty of Medicine, University of Kufa, single-centre	Iraq	21	27,6	10	11	0	4 point Likert scale Grade 0 = not satisfied Grade 1 = moderately or partially satisfied Grade 2 = greatly but not fully satisfied Grade 3 = fully or completely satisfied	"4 patients (21.4%) reported being greatly satisfied regarding the results of their treatment, 6 patients (31.6%) were partially or moderately satisfied, while 9 patients (47.3%) reported no satisfaction."
5	Al-Dhalimi 2012[163]	Open-label	Subcision, 1-3 sessions	> 26	Department of Dermatology and Venereology, Al-Sadir teaching hospital in Najaf, single-centre	Iraq	40	23,5	21	19	0	4 point Likert scale Grade 0 = not satisfied Grade 1 = moderately or partially satisfied Grade 2 = greatly but not fully satisfied Grade 3 = fully or completely satisfied	"Patients satisfaction was recorded as follows: 7 (20.59%), patients were grade 1 (partially or moderately satisfied), 22 (64.7%) patients were grade 2 (greatly satisfied) and 5 (14.7%) patients were grade 3 (fully satisfied)."
6	Alster 2007[164]	Prospective case-series	1,550nm erbium-doped fiber fractional laser vs fractional 10,600nm CO2 laser, 2-5 sessions	> 26	Washington Institute of Dermatologic Laser Surgery, single-centre	USA	53	39,6	14	39	0	Satisfaction survey	No details provided and the results are NOT reported

7	Asilian 2019[165]	RCT split-face	Nokor needle method subcision vs blunt blade subcision (BB), 1 session	26	Department of Dermatology, Skin Disease and Leishmaniasis Research Center, Isfahan University of Medical Sciences, Isfahan,	Iran	28	30,5	14	14	0	VAS scale 0 to 10 (5 = very much: 8-10; 4 = high: 6-8; 3 = moderate: 4-6; 2 = low: 2-4; and 1 = no: 0-2)	"According to patients' satisfaction, there was no significant difference between two groups during the first 2 weeks (P > 0.05), but after the first and the second months of treatment, it was significantly higher in BB group than Nokor group (P < 0.05) (Table 4)." Patient satisfaction after 2 months BB 5.01 ± 1.99 vs Nokor needle 3.89 ± 2.07, P = 0.043.
8	Azzam 2013[166]	CCT	Fractional CO2 laser vs fat grafting, 3 sessions	12	Department of Dermatology, Kasr El-Ainy Hospital, Faculty of Medicine, Cairo University, Cairo, single-centre	Egypt	20	NR	9	11	0	Satisfaction via pretreatment and posttreatment digital photos; 3 questions scar improvement, texture improvement, and overall improvement in appearance 4 point Likert scale + (0% to 25%, mild) ++ (25% to 50%, moderate) +++ (50% to 75%, marked) ++++ (75% to 100%, excellent)	"Patient satisfaction in both groups turned out to be statistically insignificant as regards scar improvement (P = 0.244), texture improvement (P = 0.500), and overall improvement in appearance (P = 0.123)."
9	Batra 2003[167]	Open-label	Combination CO2/erbium YAG laser, 1 session	130	Referral-based academic practice Beth Israel Deaconess Medical Center Cosmetic Surgery and Laser Center, Chestnut Hill, Mass, single-centre	USA	27	50,4	2	25	0	Unclear	Conclusions are drawn while method of measuring is not described, nor results are reported. "While patient responses remained very positive, satisfaction decreased over time" and "Patients undergoing laser resurfacing to treat acne were as satisfied as patients treated for photodamage"

10	Bernstein 2001[168]	RCT split-face	Non-ablative skin remodeling using a 532nm, 2 ms pulse-duration, frequency-doubled Nd:YAG laser vs no treatment, 2-4 sessions	12	Laser Surgery and Cosmetic Dermatology Centers, Marlton, NJ, single-centre	USA	22	NR	0	22	0	Unclear	Female with rhytides or acne scars were included. Authors report in Abstract subject satisfaction will be evaluated, but in Methods section information is not provided. In the Result section the authors report "Although subjective assessment of improvement and thus subject satisfaction were favorable". In the discussion it is concluded "Although the typical results with treatment appear to be difficult to measure with non-standardized photography, subject satisfaction is high."
11	Bernstein 2017[169]	Open-label	Picosecond-domain, 532 and 1,064nm Nd:YAG laser used in combination with a fractionating holographic optic, 4 sessions	30	Main Line Center for Laser Surgery, Ardmore, Pennsylvania, single-centre	USA	31	45	4	23	4	5 point Likert scale -2 = very dissatisfied -1 = dissatisfied 0 = no opinion 1 = satisfied 2 = very satisfied	"Subjects reported high satisfaction rates with the treatment results. For subjects treated with the 1,064 nm laser, 15 of 19 subjects were satisfied or very satisfied (nine very satisfied, six satisfied, two no opinion, two dissatisfied) with their treatment results. Of the eight subjects treated with the 532 nm, all were satisfied or very satisfied (four very satisfied, four satisfied)."
12	Bhatia 2006[170]	Retrospective analysis	1,320nm Nd:YAG laser, 6 sessions	26	Referral-based academic practice	USA	40	NR	0	0	40	3 point Likert scale Satisfied, somewhat satisfied, not satisfied	Included both people with acne scars or photodamage. "Overall 58.8% of patients reported satisfaction with the results from their treatments. The percentage of patients reporting satisfaction with the treatments was higher in the acne-scarring group, 62.5%, versus the photoaging group,

													55.6%."
13	Bjørn 2014[171]	RCT split-face	Fractional CO2 laser at 1 month vs 3 month intervals, 2 sessions	52	Department of Dermatology, Aarhus University Hospital, single-centre	Denmark	13	NR	NR	NR	13	11-point NRS 0 (unsatisfied) to 10 (best imaginable satisfaction)	"The patients were moderately satisfied with the treatments and the satisfaction scores were not influenced by the treatment interval 1 month postoperatively (1-month interval 5.68±1.74; 3- months interval 5.73±1.92, P=0.93) and 6 months postoperatively (1-month interval 5.83±1.70; 3-months interval 6.00±1.95, P=0.79)."
14	Boss 2000[172]	Open-label	Injection autologous cultured fibroblasts (3) with 2 week intervals, 3 sessions	208	Department of Plastic Surgery, University of Medicine and Dentristry of New Jersey, single-centre	USA	94	NR	NR	NR	94	Subjective patient satisfaction survey: 1) Satisfaction with correction: 10-point NRS 2) Satisfaction with process: 10-point NRS	Patients with rhytides, acne and traumatic scars were included. "A subjective patient satisfaction survey showed 92% of the patients were satisfied with the grade of correction". Little details were provided.
15	Brunner 2000[173]	Cross-sectional	Facial laser resurfacing with the Sharplan CO2 laser and SilkTouch Scanner, 1 session	NA	Private Practice, Princeton, New Jersey and Department of Otolaryngology, University of Toronto, Toronto, single-centre	USA/Canada	47	49,2	8	39	0	Questionnaires evaluating patient satisfaction in the following categories: (1) extent and duration of the postoperative recovery period; (2) lifestyle disruption of skin care regimes, social routines, and occupational duties; (3) surgical outcome; and (4) overall goal achievement. Responses were rated on a 5-point analogue scale	Patients received a facial resurfacing for rhytides, more youthful appearance, facial scars or acne scarring. 47/100 completed the questionnaires Satisfaction with the treatment is not explicitly mentioned. It would fit best a surgical outcome. However the scale used for surgical outcome was % of improvement, not satisfaction.

16	Cachafeiro 2016[174]	RCT	Nonablative fractional erbium laser 1,340nm vs microneedling, 3 sessions with 1 month interval, 3 sessions	38	Dermatology Outpatient Clinic of Hospital de Clínicas de Porto Alegre, single-centre	Brazil	46	26,3	21	21	4	Questionnaire on 1) perception of the degree of improvement of their scars, 2) pain during treatment and 3) recovery time 11-point NRS scale: 0 (maximum of dissatisfaction) to 10 (maximum satisfaction)	Although this was a prespecified outcome, the results are not reported.
17	Casabona 2018[175]	Retrospective pilot study	Combined use of microfocused ultrasound and a calcium hydroxylapatite dermal filler, 1 session	12	Clínica Vida, São Paulo, single-centre	Brazil	10	37,6	NR	NR	10	Questionnaire Question 1: Do you think this procedure was effective? Question 2: How long lasting was the effect of this procedure? Question 3: Do you want to undergo this procedure again? Question 4: Would you recommend this procedure to friends? 5 point Likert scale (very unsatisfied, unsatisfied, OK, satisfied, or very satisfied)	"The mean post-treatment Patient Satisfaction score was 4.9"(±0.3). Subjects were very satisfied (n = 9) or satisfied (n = 1)."
18	Chan 2004[176]	Open-label	1,320nm Nd:YAG laser monthly for 6 months, 6 sessions	32	NR	Hong Kong/Japan	27	41	0	27	0	VAS scale 0-10	Women with acne scars and wrinkles were included. the "the overall degree of patients' satisfaction was rated as 4.9"(range 0–9.8) for wrinkle reduction and 4 (range 0–10) for the treatment of acne scarring."
19	Chan 2010[177]	Open-label	CO2 laser, 1 session	26	Division of Dermatology, Department of Medicine, University of Hong Kong, Hong Kong SAR, single-centre	China	9	44,8	5	4	0	Questionnaire 4 point Likert scale Not satisfied, slight satisfied, satisfied, very satisfied	Only 5 of the 9 patients had acne scars, the other 4 had photodamage. "At 3 and 6 months posttreatment, 100% and 85.8% of patients, respectively, were overall satisfied to very satisfied with the procedure"

20	Chapas 2008[178]	Open-label	CO2 laser, 3 sessions	38	Laser & Skin Surgery Center of New York, New York, New York, single-centre	USA	15	NR	NR	NR	15	5 point Likert scale 0 = no improvement 1 = less than 25% improvement 2= 26–50% improvement 3 = 51-75% improvement 4 = > 75% improvement	Both subjects and investigators assessed overall satisfaction with treatment results. "Subjects and investigators recorded average overall quartile grading scores of 2.42 and 2.46, respectively, at 3 months following the last treatment, correlating to 26–50% consistent overall satisfaction with the improvements obtained."
21	Cho 2012[179]	Open-label	Fractional radiofrequency microneedle treatment, 2 sessions	12	NR	Korea	30	NR	12	18	0	Questionnaire 5 point Likert scale Very satisfied, satisfied, ordinary, dissatisfied, very dissatisfied	"Patient satisfaction at 12 weeks of follow-up was very satisfied in 12 patients (40%), satisfied in 14 (46.7%), ordinary in three (10%), dissatisfied in one (3.3%), and very dissatisfied in none (0%)."
22	Chrastil 2008[180]	Open-label	Erbium-doped 1,550nm fractional photothermolysis laser, 2-6 sessions	26	NR	USA	29	NR	9	20	0	4 point Likert scale 1 = 1% to 25% 2 = 26 to 50% 3 = 51 to 75% 4 = > 75%	Little data regarding satisfaction with treatment provided. "At least 50% to 75% improvement in facial and back acne scarring was seen in 23 of 29 patients. Five patients had an improvement of greater than 75% in acne scarring, 18 patients had an improvement of 50% to 75% of acne scarring, 5 patients had a 25% to 50% improvement in acne scarring, and 1 patient had less than a 25% response to treatment. Clinical improvement tended to increase with each treatment. The patients' degree of satisfaction paralleled the physicians' assessment."

23	Cole 2008[181]	Open-label	Er:YAG laser, 1-2 sessions	40	Salisbury District Hospital, single-centre	UK	29	NR	NR	NR	29	Questionnaire SAS Questionnaire including 4 questions. 0-10 VAS	Patients included had epidermal naevi, tuberous sclerosis, angiofibromata, neurofibromatosis, and scarring caused by acne and other means. They state that they want to measure how satisfied patients are with the results of treatment, but the questions are regarding "their problem area" or their appearance. Not exactly treatment satisfaction. "There was a significant improvement in patient satisfaction with appearance (SAS)"
24	Dai 2019[182]	Open-label	1,064nm Nd:YAG picosecond-domain laser, 3 sessions	26	NR	Taiwan	20	28	13	7	0	5 point Likert scale -2 = very dissatisfied -1 = dissatisfied 0 = no opinion 1 = satisfied 2 = very satisfied	"Overall, 55% and 25% of subjects were satisfied and very satisfied with their results at the 1-month follow-up, respectively. Subjects maintained satisfaction at the 3-month follow-up, with 55% and 40% being satisfied and very satisfied with the treatment outcome, respectively. There was a statistically significant correlation between subject-rated improvement score and satisfaction score (Spearman's rank ρ = 0.653, P = 0.002)."

25	Datz 2018[183]	RCT split-face	Fractional CO2 laser vs no treatment, 2 sessions	56	Department of Dermatology of the University Medical Centre Regensburg, single-centre	Germany	28	60,1	10	16	2	4 point Likert scale 1 = dissatisfied 2 = somewhat satisfied 3 = satisfied 4 = very satisfied	"Overall, 55% and 25% of subjects were satisfied and very satisfied with their results at the 1-month follow-up, respectively. At visit 1, patient satisfaction was 2.14±0.99 (n=29) for both sides, which means somewhat satisfied. 1 year after the second treatment, patients were very satisfied with the appearance of the treated half of the skin graft (3.86 0.35; n=29) and satisfied with the control side (2.83±1.00; n=29) (P<0.001)."
26	de Andrade Lima 2015[184]	Retrospective analysis	88% phenol peeling followed by microneedling, 1 session	12	Dermatology department, Santa Casa de Misericórdia do Recife, Recife, single-centre	Brazil	28	NR	NR	NR	28	Satisfaction questionnaire	Patients with wrinkles and sagging were also included, there were only 5 patients that had just acne scars, and 10 had a combination with wrinkles and sagging. No details regarding the satisfaction questionnaire were provided. "In the patient satisfaction questionnaire, 100% of patients reported satisfaction with the results. All informed they would undergo the procedure again, if necessary."
27	de Andrade Lima 2017[185]	Retrospective analysis	Percutaneous collagen induction with needles (IPCA®) associated with pulsed radiofrequency with multineedles (RFPM®), 1 session	8	Dermatology department, Santa Casa de Misericórdia do Recife, Recife, single-centre	Brazil	13	NR	6	7	0	Satisfaction questionnaire	No details provided, probably Likert scale. "All patients were satisfied with the results, shown in the questionnaires as good and very good, according to the norms proposed".

28	Dierickx 2018[187]	RCT split-face	Picosecond domain 755nm Alexandrite laser standard pulse coverage vs high pulse coverage, 5 sessions	46	Skinperium Laser Clinic, Boom, singe-centre	Belgium	7	33	NR	NR	7	Satisfaction questionnaire	Patients recorded overall improvement, if they noticed a difference from one side over the other and whether the treatment was comfortable and easy to tolerate. This is not actually covering satisfaction. "The comparison of baseline to final follow-up images of each subject found both sides to be Much or Very Much improved with no statistically significant difference between the standard and high pulse sides (P>0.05, n = 7 paired t-test). Six of the seven subjects did not note any difference between the effect on different sides of the face and four of the seven rated their overall improvement after treatment as Good, three subjects as Reasonable and one subject with Slight Improvement. All subjects found the treatment comfortable and easy t tolerate and there was no increased incidence of side effects other than the mild occurrences typically observed for this type of treatment."
29	Dierickx 2018[186]	Open-label	Injections with hyaluronic acid, 3 sessions	8	Skinperium Laser Clinic, Boom, single-centre	Belgium	12	33	4	8	0	Satisfaction questionnaire	"Subjects' satisfaction with the treatment in terms of overall facial appearance and sensation and perception of the facial skin was evaluated" This is not exactly treatment satisfaction. "The percentage of subjects who were satisfied with the overall appearance of

													their face increased from 33% at baseline to 83% at Week 36." Table 4 shows percentage of "very and somewhat satisfied". Probably Likert scale.
30	Elawar 2018[188]	Open-label	Non-insulated fractional microneedle radiofrequency treatment with smooth motor insertion, 2-6 sessions	36	Two centres in Marseille and Toulouse	France	19	38	6	19	0	Global Aesthetic Improvement Scale (GAIS) 5 point Likert scale 0 = No improvement 1 = ≤ 25% improvement 2 = 26-50% improvement 3 = 51-75% improvement 4 = 76-100% improvement	The authors report "The patients undergoing the treatment reported satisfaction with the results of the treatment, as measured subjectively using the same scale (GAIS)." "Patients reported overall satisfaction from the treatment. All of the patients reported a noticeable improvement—57.89 percent of the"patients saw 51 to 75 percent improvement, 36.84 percent saw 26 to 50 percent improvement, and 5.26 percent saw 76 to 100 percent improvement" Subjective assessment of improvement is not the same as treatment satisfaction
31	Epstein 2010[189]	Open-label	Injection with Artefill (non-resorbable polymethacrylate), 1-2 sessions	36	Nova Southestern University/Largo Medical Center Dermatology, Largo, Florida, single-centre	USA	14	NR	NR	NR	14	5 point Likert scale Significant, moderate, mild, minimal, none	Subject satisfaction as described in Figure 6; 2 patients significant, 8 moderate, 2 mild, 1 minimal and 1 none
32	Erol 2008[190]	Open-label	Intense Pulsed Light (IPL), 6-24 sessions	96	NR	Turkey	144	NR	24	120	0	A patient satisfaction phone survey 5 point Likert scale 0 = no satisfaction 1 = minimal 2 = moderate 3 = good 4 = excellent	Only 6/109 patients that completed the treatment had acne scarring. Data on treatment satisfaction were available for 86 patients. "Seventy-sixty percent were satisfied with their treatment at good to excellent degrees. Only one patient was dissatisfied with it."

33	Faghihi 2017[191]	RCT split-face	Fractionated microneedle radiofrequency (FMR) vs FMR combined with subcision, 3 sessions	12	Department of Dermatology, Skin Diseases and Leishmaniasis Research Center, Isfahan, single-centre	Iran	25	30,1	9	16	0	VAS scale 0 = no satisfaction, 10 = best probable satisfaction	VAS in FMR + Subcision group 6 (2.2) and in FMR group 5.1 (1.6) p< 0.001
34	Firooz 2016[192]	Open-label	Fractional Er:YAG laser (2,940nm), 2-5 sessions	30	Center for Research and Training in Skin Diseases and Leprosy, Tehran University of Medical Sciences, Tehran, single-centre	Iran	9	32	1	8	0	5 point Likert scale Worse = deterioration No change = improvement ≤24% Fair = improvement of 25-49% Good = improvement of 50-74% Excellent = improvement of ≥75%	"Patient evaluation of treatment outcome showed that in 6 patients satisfaction was good and in 3 of them was fair"
35	Galal 2017[193]	RCT split-face	Fractional CO2 laser therapy vs fractional CO2 laser followed by intradermal PRP injection, 3 sessions	52	Two centres in Giza	Egypt	30	26,7	9	21	0	3 point Likert scale Satisfied, partially satisfied, or dissatisfied	"Patient satisfaction was higher for the sides that were treated with combined laser and PRP (Figure 6). Fifteen patients (50%) were very satisfied with the combined treatment versus only one patient (3.3%) for the laser treatment only (Figure 2)."
36	Gheisari 2018[194]	CCT	Nokor needle subcision (NNS) vs blunt cannula subcision (BCS), 1 session	12	Dermatology clinic of two hospitals in Tehran	Iran	34	NR	9	25	0	Unclear	"patients were significantly satisfied with BCS (P = .021) but not with NNS (P = .353), during 3-month monitoring."
37	Goel 2017[195]	Open-label	Nanofractional radiofrequency, 1-3 sessions	26	Cutis skin studio, Mumbai, single-centre	India	20	NR	6	14	0	Questionnaire	Little data provided "At the final followup visit, data showed that the vast majority of patients were very satisfied with the treatment and the cosmetic outcomes achieved"
38	Gold 2012[196]	Open-label	Fractional bipolar radiofrequency energy, 3 sessions	12	NR	USA	15	35,7	2	13	0	Questionnaire	Little data provided. "For satisfaction, 67 – 92% of subjects were satisfied to very satisfied at all-time points" "Grades were based on a 5-point scale". Probably 5 point Likert

													scale.
39	Goodman 1998[197]	Retrospective analysis	CO2 laser, 1 session	12	Dermatology practice, single-centre	Australia	100	NR	NR	NR	100	Questionnaire	30/100 had acne and traumatic scarring. Little data provided "22 of 30 (73%) with acne and traumatic scarring had their expectations met." and "most conditions were treated roughly to the same satisfaction level."
40	Gotkin 2019[198]	Open-label	Fractional CO2 laser, 1 session	26	NR	USA/Italy	32	NR	7	25	0	Questionnaire 5 point Likert scale 0 = no improvement 1 = < 25% improvement 2 = 26-50% improvement 3 = 51-75% improvement 4 = 76-100% improvement	Only 4 had acne scars. Satisfaction is not discussed in Results section, but in table 2 at 6 months an improvement of 2.5 was noted.
41	Hedelund 2012[199]	RCT split-face	Fractional CO2 laser vs no treatment, 3 sessions	42	Department of Dermatology, Bispebjerg Hospital, Aarhus, single-centre	Denmark	13	33	6	7	0	NRS 0 = unsatisfied, 10 = maximal satisfaction	"Patients were satisfied with treatments and satisfaction scores were similar at 1 month (median (interquartile range, IQR)) (4 (2–7)), 3 months (4 (1–5)), and 6 months (4.5 (2–7)) postoperatively (P = 0.117)."
42	Hedelund 2010[200]	RCT	Fractional nonablative 1,540nm laser vs no treatment, 3 sessions	12	Department of Dermatology, Bispebjerg Hospital, Aarhus, single-centre	Denmark	10	45,5	4	6	0	NRS 0 = unsatisfied, 10 = best imaginable satisfaction	"Patients were satisfied with the treatment, and satisfaction scores were similar at 4 and 12 weeks after treatment (median 25% to 75% percentiles 6, IQR 3–7.5, after 4 weeks; median 5.5, IQR 1–7, after 12 weeks; P=0.1875)."
43	Hu 2009[201]	Open-label	Nonablative 1,550nm erbium-doped fiber laser, 1 session	12	NR	Taiwan	45	29	NR	NR	45	4 point Likert scale Improvement none, fair, good of excellent	Degree of improvement is not the same as satisfaction with treatment. In table 1 it shows that 60% showed a good to excellent improvement

44	Hu 2011[202]	Open-label	Ablative fractional 2,940nm Er:YAG laser, 1 session	12	NR	Taiwan	34	34,2	NR	NR	34	4 point Likert scale Improvement: Poor = ≤ 25% Fair = 25-50% Good = 50-75% Excellent = ≥ 75%	Degree of improvement is not the same as satisfaction with treatment. In table 1 it shows that 72,2% showed a good to excellent improvement
45	Ibrahim 2018[203]	CCT split-face	Microneedling plus platelet-rich plasma vs skin microneedling alone, 4 sessions	26	Department of Dermatology and Venereology outpatient clinic,Faculty of Medicine, Al-Azhar University, Cairo, single-centre	Egypt	35	24,7	NR	NR	35	4 point Likert scale Improvement: Poor = 0-24% Fair = 25-49% Good = 50-74% Excellent = 75-100%	Degree of improvement is not the same as satisfaction with treatment. "Regarding patient's satisfaction grades there was a significant improvement after both treatment modalities with insignificant differences between both treatment modalities"
46	Issaría 2011[204]	Open-label	Non-ablative fractional 1,540nm erbium laser, 4 sessions	24	Instituto Médico Láser, Servicio de Dermatología, Madrid, single-centre	Spain	20	34,7	2	18	0	Global Aesthetic Improvement Scale (GAIS) 5 point Likert scale Worse = 0% improvement No change = 0-24% improvement Improved = 25-49% improvement Much iproved = 51-74% improvement Very much improved = 75-100% improvement	Degree of improvement is not the same as satisfaction with treatment. It was more about improvement of scar size, pore size, acne lesions, sebum secretion. The authors conclude "patient satisfaction is high"
47	Kang 2009[205]	Open-label	Triple combination of dot peeling, subcision and fractional laser, 3-4 sessions	52	NR	Korea	35	31	2	8	25	Interviews 4 point Likert scale 0 = no or minimal improvement (0-25%) 1 = light improvement (26-50%) 2 = significant improvement (51-75%) 3 = marked improvement (>75%)	Degree of improvement is not the same as satisfaction with treatment. "The mean patients' subjective scores were recorded as 2, which represents significant improvement (51–75%)."

48	Karnik 2014[206]	RCT	Polymethylmethacrylate microspheres injections vs saline injections, 1-2 sessions	32	Multi-centre	USA	175	44	68	107	0	6 point Likert scale 1 = very satisfied 2 = dissatisfied 3 = somewhat satisfied 4 = somewhat satisfied 5 = satisfied 6 = very satisfied	"A total of 84% of PMMA-collagen treated subjects were satisfied or better compared with 52% for the saline control group"
49	Keller 2007[207]	Open-label	Nonablative 1,064nm Nd:YAG laser, 5 sessions	26	Department of Dermatology and Department of Surgery, Universidade de São Paulo, single-centre	Brazil	12	27,6	0	0	12	NRS 1 = lowest satisfaction 10 = highest satisfaction	"Patient satisfaction surveys at the end of the study revealed a mean satisfaction score of 8.6; 91.6% of the patients reported Grades 8 to 10 (on a scale from 0 to 10), and one patient reported Grade 6."
50	Kim 2014[208]	Open-label	Fractional radiofrequency treatment, 4 sessions	26	NR	Korea	31	28,9	11	12	8	5 point Likert scale 0 = not satisfied 1 = somewhat satisfied 2 = satisfied 3 = very satisfied 4 = extremely satisfied	Data provided in Figure 3. One patient was somewhat satisfied, 16 were satisfied, 8 were very satisfied and 6 extremely satisfied
51	Kim 2017[209]	Open-label	Transcutaneous pneumatic injection of 20% hypertonic glucose solution, 1-5 sessions	19	NR	Korea	13	33,1	4	9	0	4 point Likert scale 0 = unsatisfied 1 = slightly satisfied 2 = satisfied 3 = very satisfied	Only 6 had acne scars. Results not reported by cause of atrophic scar. "Meanwhile, the overall mean subjective satisfaction with the TPI treatment was 2.3 ± 0.9. Six of the 13 (46.2%) patients achieved subjective satisfaction of grade 3, six (46.2%) patients grade 2 and one (7.7%) patient grade 0."
52	Knight 2019[210]	Open-label	Combined 400-600nm and 800-1200nm Intense Pulsed Phototherapy (IPL), 5 sessions at 1-2 weeks intervals	14	Knight Dermatology Institute, Orlando, FL, single-centre	USA	13	23,5	3	10	0	NR	It looks like a Likert scale was used. "Eighty percent of the patients reported some level of satisfaction with treatment outcomes (Figure 4)." Figure 4 shows that 20% considered satisfaction was worse, 10% no change, 10% improved, 40% much improved, 20% very much improved. Degree of improvement is not the

													same as satisfaction with treatment.
53	Koren 2019[211]	Retrospective analysis	Combination of diluted calcium hydroxylapatite-based filler and an energy-based device, 1-5 sessions	26	The Tel Aviv Medical Center, single-centre	Israel	352	28,7	121	231	0	5 point Likert scale 0 = not at all satisfied 1 = somewhat satisfied 2 = moderately satisfied 3 = satisfied 4 = highly satisfied	Four treatments: ablative fractional CO2 laser (FACL), a radiofrequency (RF) bipolar device, a 1540 nm nonablative fractional laser (NAFL) and injection of diluted calcium hydroxylapatite (CaHA). The EBDs energy based device) were used either as monotherapy or in combination with diluted CaHA. Satisfaction scores for the various options (mono or combination therapies) varied between 1.5 and 3.4, with FACL +CaHA in separate session scoring highest, and NAFL monotherapy scoring lowest.
54	Lan 2018[212]	Open-label	Fractional micro-plasma radio-frequency, 3 sessions	52	The Laser Research Unit, Department of Dermatology, Southwest Hospital, Chongqing, single-centre	China	95	23,2	53	33	9	4 point Likert scale 0 = dissatisfied 1 = somewhat satisfied 2 = satisfied 3 = very satisfied	"No patients were dissatisfied at the clinical response of the treatment for atrophic acne scars (Fig. 5). The average score for satisfaction for improvement of acne scars was 2.44 at last follow-up."
55	Lee 2009[213]	RCT split-face	585nm PDL vs 1064nm Nd:YAG laser, 4 sessions	14	NR	Korea	18	23	10	8	0	NRS 0 (neutral) to 10 (highly satisfied)	"For PDL treatment, patient satisfaction scores increased from 0 (baseline) to 4.5 (8 weeks after final treatment); for Nd:YAG laser treatment, patient satisfaction scores increased from 0 (baseline) to 5.1 (8 weeks after final treatment). However, again no significant difference was found between the two modalities."

56	Lee 2010[214]	Open-label	Subdermal minimal surgery technology, 3 sessions	26	NR	Korea	10	29,2	6	4	0	5 point Likert scale -1 = worse 0 = no change 1 = mild improvement 2 = moderate improvement 3 = strong improvement	Degree of improvement is not the same as satisfaction with treatment. Mean score (Table 2) 2.1. "all patients were satisfied with the procedure according to their self-assessments."
57	Loss 2018[215]	Open-label	Adapalene 0.3% gel, first weeks once daily for 4 weeks, then 20 weeks twice daily	72	Department of Dermatology, Johns Hopkins School of Medicine, Baltimore, single-centre	USA	20	35,7	11	9	0	7-item subject satisfaction questionnaire	No details of the 7-item questionnaire were provided. Not clear how the remaining patients scored. "In the subject satisfaction questionnaire, the vast majority of subjects (88.2%) were satisfied with the effectiveness of the treatment and reported that they would consider using this treatment again (83.3%). In addition, almost 90% of subjects (16/18) were satisfied with the rapid onset of efficacy."
58	Maas 2019[216]	Open-label	Microfocused ultrasound with visualization (MFU-V), 3 sessions	26	NR	USA	20	44,5	8	12	0	Satisfaction questionnaire	No details on the questionnaire. "The results of the 90-day subject satisfaction questionnaire show that 17 subjects (94.4%) noted improvement in their acne scores and expressed some degree of satisfaction"

59	Mahmoud 2010[217]	RCT	1,550nm erbium fractionated laser 10 mJ vs 40 mJ, 5 sessions	26	Department of Dermatology, Henry Ford Hospital, Detroit, Michigan, single-centre	USA	15	NR	3	12	0	Questionnaire 5 point Likert scale 0 = no improvement 1 = minor improvement (1-25%) 2 = moderate improvement (26-50%) 3 = marked improvement (51-75%) 4 = very significant improvement (76-100%)	Degree of improvement is not the same as satisfaction with treatment. "A post-treatment questionnaire distributed to the patients after the end of the study and completed by 11 of the 13 treated patients showed that 18% (n = 2) considered that they had 1% to 25% improvement, 55% (n = 6) had 51% to 75% improvement, and 27% (n = 3) had 76% to 100% improvement " and "patient satisfaction was significantly greater in patients treated with higher fluences (p<.05) but not in patients treated with higher densities, which implies that the higher the energy, the better the response was to the laser treatment."
60	Majid 2014[218]	Open-label	Fractional CO2 laser, 3-4 sessions	52	Cutis Skin and Laser Institute, Srinagar, Jammu and Kashmir, single-centre	India	60	NR	25	35	0	Survey (no details)	"Patient satisfaction survey revealed that about 73% (44 cases out of the total of 60) were satisfied with the treatment results. For the remaining 16 cases, the improvement achieved was not up to their expectations."
61	Montesi 2007[219]	Open-label	Bipolar radiofrequency, 6-8 sessions	28	NR	USA/Italy	30	44	4	26	0	NR	Only 4 patients had acne scars. In abstract "they expressed their satisfaction" and in Discussion "many patients told us they had received positive comments from family members and friends" "Patients see consistently improvements with the Aluma device with each treatment session"

62	NCT02025088[220]	RCT	Non-ablative fractional erbium 1,340nm laser vs microneedling, unclear how often	NR	NR	Brazil	23	NR	NR	NR	NR	NR	This is an ongoing trial, no details are provided "The secondary objectives are: assess the degree of satisfaction of participants with both techniques..."
63	NCT02955381[221]	RCT	Restylane® Silk vs placebo, 2 sessions	26	NR	USA	26	NR	NR	NR	26	VAS scale 0 = not satisfied 10 = extremely satisfied	This is an ongoing trial, no data are provided
64	NCT03170596[222]	RCT split-face	Microneedling vs tazaratone 0.1% gel, 1 time vs daily application of the gel	26	NR	India	36	NR	NR	NR	36	Patient satisfaction using Patient's global assessment score 0 (no response) to 10 (maximum)	It looks like more to be about improvement than satisfaction. This is an ongoing trial, no data are provided
65	Nirmal 2013[223]	Open-label	Er:YAG 2,940nm fractional laser, 4 sessions	20	Tertiary care teaching hospital, single-centre	India	25	NR	15	10	0	4 point Likert scale None = 0-25% Fair = 26-50% Good = 51-75% Excellent improvement = 76-100%	Degree of improvement is not the same as satisfaction with treatment. "Patient's satisfaction scores were higher in comparison to both observers' rating with a relatively higher proportion of improvement grading in the good and excellent categories"
66	Ortiz 2010[224]	Retrospective analysis	Fractional CO2 laser, 1 session	104	Department of Dermatology, University of California at Irvine, Irvine, California, single-centre	USA	10	NR	NR	NR	10	NR	"Only 6 patients had acne scars, remainder photodamage. As with any procedure, results tend to fade over time due to relaxation of the tightening process and the natural progression of aging. Long-term patient satisfaction was upheld regardless." and "All subjects reported that they were pleased with their immediate and long-term results. However, these subjective results were not graded."

67	Panchaprateep 2015[225]	Open-label	Low fluence 585nm Q-switched Nd:YAG laser, 3 sessions	12	NR	Thailand	25	20,1	17	8	0	Questionnaire 5 point Likert scale (excellent, good, fair, some, little/no/improvement)	Degree of improvement is not the same as satisfaction with treatment. "Subjective degrees of satisfaction were also evaluated." and "Eighty-nine percent of subjects assessed their overall improvement as good or excellent (Grade 4, 5)."
68	Peterson 2011[226]	Open-label	Fractional laser with radiofrequency and fractionated radiofrequency, 5 sessions	34	Dermatology Cosmetic Laser Associates of La Jolla, San Diego, California, single-centre	USA	15	42	3	12	0	Questionnaire 5 point Likert scale 0 = 0% 1 = 1-24% 2 = 25-49% 3 = 50-74% 4 = 75-100%	Degree of improvement is not the same as satisfaction with treatment. "Patients rated their satisfaction and overall improvement at day 60, 120, 150, and 210, which corresponded to treatment sessions 3 and 5 and postoperative visits 30 and 90 days from the final treatment session. Patient satisfaction scores showed no significant changes from day 60 to 210 (from 1.47 to 2, p = .21)" and "Patient satisfaction scores did not improve, although patient overall improvement scores did. We speculate that our patients had higher expectations of the device upon study entry"
69	Petrov 2016[227]	RCT	Fractional CO2 laser vs local treatments vs CO2 laser combined with platelet-rich plasma, 1-4 sessions	NR	Acibadem Sistina Clinical Hospital, at the Department of Dermatovenerology, Skopje, single-centre	Republic of Macedonia	232	NR	NR	NR	232	5 point Likert scale 1 = not satisfied 2 = partially not satisfied 3 = neutral 4 = partially satisfied 5 = totally satisfied	72 patients with acne scars were included. Details are provided for only acne scars. In the control group (local treatments) 56,52% (26/46) was partially not satisfied or not satisfied, compared to 0% in the fractional CO2 laser group. 17,39% (8/46) of the control group was neutral compared with 10,71% (3/28) in the fractional CO2 laser

													group. 26.09% (12/46) was partially or totally satisfied in the control group compared to 89,28% (25/28) in the fractional CO2 laser group.
70	Petrov 2016b[228]	Open-label	Fractional CO2 laser, unclear how often	NR	Acibadem Sistina Clinical Hospital, at the Department of Dermatovenerology, Skopje, single-centre	Republic of Macedonia	40	28,2	NR	NR	40	Scale of satisfaction by treatment (no details)	Although this was a prespecified outcome, the results are not reported.
71	Politi 2016[229]	Open-label	Mid-infrared non-fractional 1,540nm Er:Glass laser with integrated cooling-vacuum assisted technology, 3-6 sessions	40	NR	Israel	25	NR	10	15	0	5 point Likert scale Score 1-5 (1 being not satisfied and 5 being very satisfied)	"Patients' self-rated satisfaction was 3-5 (mean 4.2 ± 1.1) at 3 months follow-up."
72	Politi 2018[230]	Open-label	Mid-infrared non-fractional 1,540nm Er:Glass laser with integrated cooling-vacuum assisted technology, 3-7 sessions	34	NR	Israel	22	24,5	8	14	0	5 point Likert scale Score 1-5 (1 being not satisfied and 5 being very satisfied)	"Patients' self-rated satisfation ranged from 3 to 5 (mean 4.2± 0.8) at the 3-month follow-up visit."
73	Qin 2015[231]	Open-label	Fractional bipolar radiofrequency, 4 sessions	28	Department of Dermatology, Peking University Shenzhen Hospital, single-centre	China	26	22,6	10	16	0	4 point Likert scale 0 = dissatisfied 1 = somewhat satisfied 2 = moderately satisfied 3 = very satisfied	"Patient satisfaction was " very satisfied " in 3 patients (13.0%) at 4-week follow-up and in 13 patients (56.5%) at 12-week follow-up. 9 patients (39.1%) felt moderately satisfied. Only 1 patient (4.4%) felt somewhat satisfied, and none felt dissatisfied at 12-week visit."

74	Rojas 2015[232]	Retrospective analysis	Punch surgery and/or CO2 laser 10,600nm, unclear how often	NR	El servicio de dermatología del Hospital Clínico de la Universidad de Chile, single-centre	Chili	47	29	24	23	0	Validated questionnaire NRS 0-10 (higher means greater satisfaction)	Although the authors refer to a validated scale, there is no reference supporting this. 63,8% received treatment with CO2 laser only, 10,6% only with punch surgery and 25.6% with the combination of thes two. "Respecto al grado de satisfacción de los pacientes, el 9% refirió un puntaje de 10 luego del procedimiento, el 32% entre 7-9, el 43% entre 4-6 y un 17% entre 0-3". Translated: with respect to the grade of satisfaction, 9% scored a 10 after treatment, 32% 7-9, 43% between 4-6 and 17% between 0-3.
75	Sadaawi 2018[233]	RCT	Glycolic acid peel 35% vs microneedling vs combination of both procedures, 6 sessions	12	Outpatient clinics of Dermatology, Venereology and Andrology Department, Faculty of Medicine, Zagazig University Hospitals, single-centre	Egypt	30	29,2	10	20	0	3 point Likert scale Mild, good, very good	"Patients' satisfaction was higher in group III (P = 0.04)". In group I (dermapen) 2 scored mild, 6 good and 2 very good, in group II (microneedling) 6 scored mild, 3 good and 1 very good, in group III (combination of the two procedures). 3 scored mild, 3 good and 4 very good.
76	Samuthrsindh 2011[234]	Open-label	Ablative fractional CO2 laser, 1-3 sessions	NR	NR	Thailand/ Germany	85	40	18	67	0	4 point Likert scale 0 = no improvement 1 = minimal improvement ie improvement of 1-25% 2 = moderate improvement ie improvement 26-50% 3 = marked improvenment ie improvement >50%	44/85 patients had acne scars. Patient satisfaction was scored by a blinded observer. Degree of improvement is not the same as satisfaction with treatment. For acne scars "mean satisfaction score: 1.7±1.0"

77	Sapra 2015[235]	Open-label	Injectable poly-L-lactic acid, 3-4 sessions	26	Institute of Cosmetic and Laser Surgery, Oakville, Ontario, single-centre	Canada	22	37,6	9	13	0	Subject treatment satisfaction score (STSS) 5 point Likert scale Score of 0 indicating no satisfaction with treatment and a score of 4 indicating excellent satisfaction	"An increase in mean STSS score was observed for images taken by the VISIA-CR system from Session 2 (1.4) to the follow-up visit (2.5), indicating an improvement in treatment satisfaction over time. The proportion of patients achieving an STSS score of ≥3, indicating very good to excellent satisfaction with treatment, was 54.5%"
78	Tanzi 2004[236]	RCT split-face	Long-pulsed 1,320nm Nd:YAG laser vs 1450nm diode laser, 3 sessions	64	Washington Institute of Dermatologic Laser Surgery, Washington, DC, single-centre	USA	20	36,7	0	0	20	NRS Degree of satisfaction on a scale of 1 (lowest) to 10 (highest)	"Patient satisfaction surveys at the end of the study revealed a mean satisfaction score of 4.6 on the 1320- nm Nd:YAG-treated facial half and 5.7 on the 1450-nm diode laser-treated facial half."
79	Tatliparmak 2019[237]	Retrospective analysis	Combined fractional carbon dioxide laser and fractional microneedle radiofrequency, 3-6 sessions	116	Multi-centre	Turkey	72	27,6	29	43	0	5 point Likert scale –2: not at all satisfied –1: not satisfied 0: not sure 1: satisfied 2: very satisfied	"Overall, 45.8% of patients were very satisfied and 44.4% were satisfied with the treatment (Table 1, Figures 1 and 2).". 33 were very satisfied, 32 satisfied, 0 not satisfied, 7 not sure.
80	Tay 2008[238]	Open-label	Low-fluence 2,940nm Er:YAG laser, 1-2 sessions	8	Division of Dermatology, Changi General Hospital, Singapore, single-centre	Singapore	9	33	3	6	0	Patient satisfaction score 3 point Likert scale A = not satisfied, B = somewhat satisfied, and C= highly satisfied	"All patients were somewhat satisfied with the treatment, and all requested further treatments."
81	Taylor 2017[239]	Retrospective analysis	Combination of a superficial chemical peel (trichloracetic acid), subcision, and fractional CO2 laser, 1 session	4	NR	USA	114	NR	39	75	0	NR	"Ninety percent of the patients were satisfied with their final result and desired no additional treatment."

82	Tenna 2017[240]	RCT	Autologous fat grafts plus platelet-rich plasma in combination with fractional CO2 laser vs only autologous fat grafts plus platelet-rich plasma, 1 session	26	NR	Italy	30	NR	0	0	30	Italian version of the FACE-Q module	FACE-Q scales for health-related quality of life, early life impact, satisfaction with outcomes, and decision to have treatment. Klassen AF *et al*. *Plast Reconstr Surg* 2015; 135:378-86. "We applied a t test on unpaired data, comparing the results obtained with the FACE-Q postoperative module in both group A and group B and found no significant difference (p value = 1.000)." and "Finally, the FACE-Q module, which measures patient satisfaction after treatment, confirmed great satisfaction in face image perception after treatment." The FACE-Q for facial appearance is a non-specidic general satifsfaction tool to exactly measure that. Not to assess trestment satisfaction. https://eprovide.mapi-trust.org/instruments/face-q2
83	Thomas 2017[241]	Open-label	Laser therapy, not specified, unclear how often	NR	Skin & Cancer Foundation Australia, Westmead, single-centre	Australia	147	33	49	98	0	Comprehensive Satisfaction Questionnaire (PSQ-18)	Patients responded to the satisfaction statements on a five-point categorical scale. To achieve the final score, responses were organized so that high scores were allocated to reflect satisfaction with medical care and items within the same domain were averaged to create the seven subscale scores. 25/147 had acne scarring, the others suffered from rosacea (n=54) or photoaging

													(n=68). "Overall, satisfaction with laser therapy and subjective benefit derived from laser therapy was high across all three skin conditions. 44.4% of patients rated the overall satisfaction with the highest maximum score (5/5) with 37.0% of patients rating overall satisfaction as 4/5. Treatment was viewed as beneficial, 40.7% of patients scored it 5/5 and 44.4% scored treatment as 4/5. Patients were more likely to view treatment as being beneficial for their skin condition at younger ages (P = 0.032) and had greater overall satisfaction with increasing age (P = 0.031)." The PSQ-18 does not handle satisfaction with treatments such. Domains are: General Satisfaction; Technical Quality; Interpersonal Manner; Communication; Financial Aspects; Time Spent with Doctor; Accessibility and Convenience. https://www.rand.org/health-care/surveys_tools/psq.html

84	Trelles 2014[242]	Open-label	Fractional ablative microplasma radiofrequency and acoustic pressure ultrasound, 4 sessions	38	Instituto Medico Vilafortuny, Cambrils, Tarragona, single-centre	Spain	19	35,2	5	14	0	5 point Likert scale Very dissatisfied, dissatisfied, somewhat satisfied, satisfied, and very satisfied	"Patients reported to be Somewhat Satisfied (16%), Satisfied (53%), and Very Satisfied (31%)."
85	Tzikas 2004[243]	Open-label	Radiance fine needle soft tissue filler, 1 session	26	Private practice in Delray Beach, Florida, single-centre	USA	90	NR	5	85	0	4 point Likert scale Excellent, good, fair, poor	Only two patients suffered from acne scarring. No separate data for acne patients. "Patient satisfaction with treatment was uniformly high: 47% of patients rated their satisfaction as excellent; 41% rated their satisfaction as good. Only 2% reported poor overall satisfaction"
86	Verner 2016[244]	Open-label	Fractional bipolar radiofrequency, 3-5 sessions	34	Clinic Tel Aviv, Kiriat Ono, single-centre	Israel	12	36	4	8	0	Global Aesthetic Improvement Scale (GAIS).	"The patient satisfaction survey revealed that half of the patients (6 out of 12) were very satisfied, while the other half (6 out of 12) reported to be satisfied with the procedure and results." These results do not match the outcomes of the GAIS.
87	Wang 2010[245]	Open-label	Fractional ablative CO2 laser, 2 sessions	26	Department of Dermatology, Changi General Hospital, Singapore, single-centre	Singapore	5	27	4	1	0	Patient satisfaction score 3 point Likert scale A = not satisfied, B = somewhat satisfied, and C= highly satisfied	"All patients were somewhat satisfied with the treatment."
88	Weinstein 2019[246]	RCT split-face	Tranilast 8% liposomal gel + isotretinoin	12	NR	Israel	40	28	8	32	0	4 point Likert scale 1 = high and 4 = none	"The average satisfaction rate was higher among subjects treated with the combined isotretinoin and tranilast protocol compared with the subjects who were treated with 8% tranilast liposomal gel alon (2.56 vs. 3.50, respectively, P = 0.001).

89	Yang 2016[247]	RCT split-face	Fractional 1,550nm Er:Glass laser vs no treatment, 4 sessions	24	Department of Dermatology, Huashan Hospital, Fudan University, Shanghai, single-centre	China	30	NR	0	0	30	4 point Likert scale Very good, good, medium, inadequate	"As for the patient self-evaluation, three patients (11.54%) rated their satisfaction as very good, 11 (42.31%) rated as good, and 8 (30.77%) rated as medium in the treated side. In contrast, one patient (3.85%) rated his or her satisfaction as very good, 5 (19.23%) rated as good, and 2 (7.69%) rated as medium in the control side."
90	Yeung 2012[248]	Open-label	Combination of bipolar radiofrequency and fractional diode laser, 1-5 sessions	32	NR	Hongkong /USA	24	NR	0	0	24	Questionnaire 4 point Likert scale 0 = not satisfied 1 = somehwat satisfied 2 = moderately satisfied 3 = very satisfied	"after the final treatment; 43.8% of patients reported being satisfied or very satisfied with the treatment 1 month after the final treatment. The degree of patient satisfaction was increased at the 3-month visit, with 63.2% of patients reporting being satisfied or very satisfied with the procedure."
91	Yoo 2009[249]	Open-label	Fractional 1,540nm photothermolysis, 3 sessions	24	NR	Korea	16	29,8	6	10	0	NRS 1 (lowest) to 10 (highest)	"All the patients indicated satisfaction with the results of their treatment and stated that they would undergo treatment again. The patient satisfaction survey at the end of the study revealed a mean satisfaction score of 8.4. Eighty-seven percent of the patients (14 patients) reported a satisfaction rating of Grades 8–10 (on a scale of 0–10), and two patients reported a satisfaction rating of Grades 6 and 7, respectively."

92	Yu 2016[250]	Open-label	Hynbrid energy (HE) and trifractional technology (TF), 4 sessions	9	Ospital ng Makati and the Quirino Memorial Medical Center, two centres	Philippines	13	NR	9	4	0	4 point Likert scale Very satisfied, satisfied, slightly satisfied, and unsatisfied	"After HE treatments, 62.5% of subjects said that they were satisfied with their improvement, while 37.5% reported that they were slightly satisfied. Level of satisfaction increased after additional TF treatments with 75% of subjects reporting that they were satisfied and 25% of subjects were slightly satisfied. No subject reported dissatisfaction with treatment.
93	Yu 2016b[251]	Open-label	Dermabrasion combined with ReCell®, 1 session	78	Plastic Surgery Hospital, Chinese Academy of Medical Sciences and Peking Union Medical College, single-centre	China	19	NR	0	0	19	Patient Satisfaction Score	7/19 suffered from acne scars. No details of the Patient Satisfaction Score are provided. In table 2 values between 0 and 9 are listed. In the post acne group the improvement was 4.4 ±1.7.
94	Yug 2016[252]	Open-label	Serial high-concentration (95%) trichloroacetic acid, 6 sessions	52	NR	USA	3	NR	1	2	0	NR	"All three patients reported high satisfaction with the procedure"
95	Zhang 2019[253]	Open-label	Picosecond Alexandrite Laser With DLA, 3 sessions	26	Department of Cosmetic Laser Surgery, Hospital of Dermatology, Chinese Academy of Medical Sciences, Peking Union Medical College, Nanjing, single-centre	China	20	21,2	13	7	0	5 point Likert scale –2: very dissatisfied –1: dissatisfied 0: no opinion 1: satisfied 2: very satisfied	"In addition, patient satisfaction data showed that 50% and 30% of the patients were “satisfied” and “very satisfied” with the treatment results, respectively, with no patients “very dissatisfied” or “dissatisfied”."

96	Zhou 2016[254]	RCT split-face	Conditioned medium of adipose-derived stem cells (ADSC-CM) after fractional CO2 laser vs fractional CO2 laser combined with Dulbecco's modified Eagle's media (DMEM), 3 sessions	16	Single-centre	China	22	36,4	10	12	0	Subjective satisfaction scale 0 (aggravated) to 4 (much improved)	13 patients suffered from acne scars. Degree of improvement is not the same as satisfaction with treatment. "In acne scar group, the subjective satisfaction scores in both sides also increased successively with treatment times (Figure 3). One month after the third treatment, ADSC-CM side subjective satisfaction score value was 2.35 ± 0.69, which was significantly higher than that in DMEM side (2.08 ± 0.76)."
97	Zhu 2013[255]	Open-label	Autologous platelet-rich plasma combined with erbium fractional laser therapy, 2 sessions	12	NR	China	22	28	5	17	0	4 point Likert scale 0 = unsatisfied 1 = slightly satisfied 2 = satisfied 3 = very satisfied	"According to the quartile grading scale, the self-evaluation was 3.3±0.36, and 91% of patients described themselves as 'satisfi ed' or 'very satisfied' with their overall improvement, and 45% wanted to receive further treatment."

ADSC-CM = conditioned medium of adipose-derived stem cells, BB = blunt blade, CCT = controlled clinical trial, CO2 = carbon dioxide, DMEM = Dulbecco's modified Eagle's media, GAIS = Global Aesthetic Improvement Scale, IPL = intense pulse light therapy, N/A = not applicable, nd: YAG = neodymium-doped yttrium aluminum garnet, NR = not reported, NRS = numeric rating scale, PMMA = Polymethylmethacrylate, PDL = pulsed dye laser, QD = once daily, RCT = randomised controlled trial, STSS = Subject treatment satisfaction score, VAS = visual analogue scale, vs = versus