Supplemental file

Supplementary methods: PubMed Search Strategy

PubMed Search strategy

("hydroxychloroquine"[MeSH Terms] OR "hydroxychloroquine"[All Fields] OR ("chloroquin"[All Fields] OR "chloroquine"[MeSH Terms] OR "chloroquine"[All Fields] OR "chloroquine s"[All Fields] OR "chloroquines"[All Fields])) AND ("severe acute respiratory syndrome coronavirus 2"[Supplementary Concept] OR "severe acute respiratory syndrome coronavirus 2"[All Fields] OR "ncov"[All Fields] OR "2019 ncov"[All Fields] OR "covid 19"[All Fields] OR "sars cov 2"[All Fields] OR (("coronavirus"[All Fields] OR "cov"[All Fields]) AND 2019/11/01:3000/12/31[Date - Publication]) OR ("severe acute respiratory syndrome coronavirus 2"[Supplementary Concept] OR "severe acute respiratory syndrome coronavirus 2"[All Fields] OR "sars cov 2"[All Fields]))



Supplementary Figure S1. Flowchart of study selection



Supplementary Figure S2. Risk of bias assessment of randomized controlled trials



Supplementary Figure S3. Effect of early treatment with HCQ on COVID-19 symptom resolution by 14 to 30 days



Supplementary Figure S4. Effect of early treatment with HCQ on time to COVID-19 symptom resolution in days



Supplementary Figure S5. Effect of early treatment with HCQ on viral clearance at 14 days