S12 Table

S12 Table. Reported outcome statistics from experimental and quasi-experimental studies that implemented interventions with consumables provision components.

Significant improvement ($p \leq 0.05$) | Non-significant impact ($p > 0.05$) | Significant deterioration ($p \leq 0.05$) | No statistical significance reported | Not measured

Study	Intervention Arm	Indicator Type	Availability of drinking water	Availability of handwashing water	Availability of soap	Supplies for latrine cleaning
Alexander et al. (2013)	Budget	Observed	p=0.015 of Poisson regression coefficient of difference between intervention and control.	p=0.003 of Poisson regression coefficient of difference between intervention and control.	p<0.001 of Poisson regression coefficient of difference between intervention and control.	
	Accountability	Observed	p<0.001 of Poisson regression coefficient of difference between intervention and control.	p<0.001 of Poisson regression coefficient of difference between intervention and control.	p<0.001 of Poisson regression coefficient of difference between intervention and control.	
	Maintenance	Observed	p=0.015 of Poisson regression coefficient of difference between intervention and control.	p=0.01 of Poisson regression coefficient of difference between intervention and control.	p<0.001 of Poisson regression coefficient of difference between intervention and control.	
	Latrine construction	NA				

Alexander et al. (2014)	Water and handwashing	Observed		p<0.001 for Yates uncorrected chi-squared test between intervention and non-intervention schools.	No comparative statistic reported.	No comparative statistic reported.
		Reported		p=0.07 for Yates uncorrected chi-squared test between intervention and non-intervention schools.	No comparative statistic reported.	No comparative statistic reported.
				p=0.03 for Yates uncorrected chi-squared test between intervention and non-intervention schools.		
Alexander et al. (2018)	Treatment	Observed		p=0.0548, p=0.0319, p=0.3434 for Paired t-test for change in availability within 3 intervention groups from baseline to average follow up. [1]	p=0.0037, p=0.0165, p=0.0030 for Paired t-test for change in availability within 3 intervention groups from baseline to average follow up. [1]	No comparative statistic reported.
		Reported		p=1.000, p=0.6637, p=0.3938 for Paired t-test for change in availability within 3 intervention groups from baseline to average follow up. [1]	p<0.001, p=0.0296, p<0.001 for Paired t-test for change in availability within 3 intervention groups from baseline to average follow up. [1]	p=0.0550, p=0.0710, p=0.2647 for Paired t-test for change in availability within 3 intervention groups from baseline to average follow up. [1]
				p=0.1679, p=0.3434, p=0.1679 for Paired t-test for change in availability within 3 intervention groups	p=0.0017, p=0.0009, p=0.0011 for Paired t-test for change in availability within 3 intervention	p=0.0852, p=0.0343, p=0.0629 for Paired t-test for change in availability within 3 intervention

				from baseline to average follow up. [1]	groups from baseline to average follow up. [1]	groups from baseline to average follow up. [1]
Bohnert et al. (2016)	Treatment	Observed				No comparative statistic reported.
Booysen, MJ (2019)	Treatment	NA				
Buxton et al. (2019)	Treatment	NA				
Caruso et al. (2014)	Latrine cleaning and handwashing	Observed		p=0.17 for comparison of intervention (average of 4 follow up visits) versus control arms adjusting for baseline values.	p<0.01 for comparison of intervention (average of 4 follow up visits) versus control arms adjusting for baseline values.	
	Handwashing	Observed		p=0.27 for comparison of intervention (average of 4 follow up visits) versus control arms adjusting for baseline values.	p<0.01 for comparison of intervention (average of 4 follow up visits) versus control arms adjusting for baseline values.	
Saboori et al. (2013)	Latrine cleaning and handwashing	Observed		p=0.07 for comparison of intervention (average of 4 follow up visits) versus control arm using logistic regression.	p<0.0001 for comparison of intervention (average of 4 follow up visits) versus control arm using logistic regression.	
		Reported		p=0.32 for comparison of intervention (at final follow up visit) versus control arm using a student t-test.	p<0.0001 for comparison of intervention (at final follow up visit) versus control arm using a student t-test.	

	Handwashing	Observed		p=0.31 for comparison of intervention (average of 4 follow up visits) versus control arm using logistic regression.	p<0.0001 for comparison of intervention (average of 4 follow up visits) versus control arm using logistic regression.	
		Reported		p=0.03 for comparison of intervention (at final follow up visit) versus control arm using a student t-test.	p<0.0001 for comparison of intervention (at final follow up visit) versus control arm using a student t-test.	
Karon et al. (2017)	Treatment	Observed		p=0.002 for comparison of intervention versus non-intervention arm.	p=0.003 for comparison of intervention versus non-intervention arm.	
Kochurani et al. (2009)	Treatment	Observed	Soap was not more available in intervention schools compared to non-intervention schools[2].		Soap was more available in intervention schools than non-intervention schools, but rare in all schools (1-8% of schools)[3].	

[1] P-values from intervention groups are reported together since intervention arms (provision of different types of menstrual hygiene products) are not expected to impact Infrastructure Maintenance or WASH Consumables outcomes.

[2] Study reports p=0.23, however, authors caution that reported p-values are only indicative.

[3] Study reports p=0.01, however, authors caution that reported p-values are only indicative.