SUPPLEMENT

Supplementary Table 1. MRI Characteristics

Study	MRI Characteristics								Reader	
	Vendor	Scanner	Magnetic Strength (T)	Median Time Interval (Hours)*	Slice Thickness (mm)	Intersplice Gap (mm)	b Value	Image Plane	No. of Readers	Years of Experience
Higashida et al. [1]	NA	NA	1.5 or 3	NA†	2-8§	NA	1000 or 2000	Axial	2	NA
Shimizu et al. [2]	Siemens	MAGNETOM Skyra	3	96†	2	NA	1000 or 2000	NA	2	10
Szabo et al. [3]	Siemens	Vision, Sonata/TRIO, SKYRA	1.5 or3	NA (range: < 144)	2	NA	1000 or 2000	Axial	NA	NA
Tynas and Panegyres [4]	NA	NA	NA	96 (IQR: 48–576)	NA	NA	NA	NA	NA	NA
Jain et al. [5]	Philips	Ingenia	1.5 or 3	47.5 (range: 25–114)	NA	1	1000 or 2000	Axial	NA	NA
Lanzone et al. [6]	NA	NA	1.5	NA	NA	NA	NA	NA	NA	NA
Güngör Tunçer et al. [7]	NA	NA	NA	NA (mean: 11)	NA	NA	NA	NA	NA	NA
Park et al. [8]	Philips	AchievaTx	3	7 (range: 2–480)	5	NA	2000	NA	NA	NA
Döhring et al. [9]	Philips	Intera Achieva	3	NA (range: 24–72)	3	NA	NA	NA	NA	NA
Kim et al. [10]	Philips	Intera Achieva	1.5 or 3	8.5 (range: 1–72)	3, or 5	NA	1000, 2000, or 3000	NA	2	NA
Scheel et al. [11]	GE	GE Centricity RA 1000	1, 1.5, or 3	NA	3, 5, or 6§	NA	NA	NA	2	NA
Uttner et al. [12]	Siemens	Symphony	1.5	29.5 (range: 16.5–41)‡	3	NA	1000	Axial, coronal	NA	NA
Ahn et al. [13]	GE	Signa	1.5	6 (IQR: 4–9)	5	NA	1000	NA	1	NA
Auyeung et al. [14]	NA	NA	1.5	NA (mean: 44.4 ± 16.3)	NA	NA	NA	NA	2	NA
Ueno et al. [15]	Siemens	Magnetom Symphony	1.5	range: 24-72‡	5	1.5	1000	Axial	NA	NA
Lee et al. [16]	Siemens/GE	Vision/Signa	1.5 or 3	NA (mean: 37.5 ± 20.4)	4 or 5	NA	1000	Axial	1	NA
Alberici et al. [17]	Philips	NA	1.5	NA (range: 8–48)	4	NA	NA	Axial	2	NA
Enzinger et al. [18]	Siemens/Philips	Symphony/Gyroscan NT, Gyroscan ACS	1.5	66.6 ± 54.6	5	1	1000	Axial	2	NA
Winbeck et al. [19]	Siemens	Magnetom Symphony	1.5	21.7 (range: < 48)	4 or 6	NA	1000	Axial, coronal, sagittal	1	NA
Sedlaczek et al. [20]	Siemens	Magnetom Vision or Magnetom Sonata	1.5	NA (range: 2–48)‡	5	0	1000 or 2000	Axial, coronal	NA	NA
Huber et al. [21]	Siemens	Magnetom Symphony	1.5	18 (range: 3–60)	5	NA	1000	Axial	2	NA
Strupp et al. [22]	Siemens	Vision	1.5	26 (range: 2–144)†	6	NA	NA	Coronal	2	NA

*Median time interval between symptom onset and MRI scan when the highest detection rates were observed, †Detection rates of subgroups for different time intervals were available, ‡Detection rates on follow-up DWI scans were available, §Detection rates of subgroups for different slice thickness were available. DWI = diffusion-weighted imaging, IQR = interquartile range, MRI = magnetic resonance imaging, NA = not available

REFERENCES

1. Higashida K, Okazaki S, Todo K, Sasaki T, Ohara N, Kohara N, et al. A multicenter study of transient global amnesia for the better detection of magnetic resonance imaging abnormalities. *Eur J Neurol* 2020;27:2117-2124
2. Shimizu K, Hara S, Hori M, Tanaka Y, Maehara T, Aoki S, et al. Transient global amnesia: a diffusion and perfusion MRI study. *J Neuroimaging* 2020;30:828-832
3. Szabo K, Hoyer C, Caplan LR, Grassl R, Griebe M, Ebert A, et al. Diffusion-weighted MRI in transient global amnesia and its diagnostic implications. *Neurology* 2020;95:e206-e212
4. Tynas R, Panegyres PK. Factors determining recurrence in transient global amnesia. *BMC Neurol* 2020;20:83
5. Jain TP, Patel R, Gawarikar Y. Transient global amnesia: diffusion MRI findings. *Indian J Radiol Imaging* 2018;28:6-9
6. Lanzone J, Ricci L, Assenza G, Ulivi M, Di Lazzaro V, Tombini M. Transient epileptic and global amnesia: real-life differential diagnosis. *Epilepsy Behav* 2018;88:205-211
7. Güngör Tunçer Ö, Aksay Koyuncu B, Vildan Okudan Z, Altindag˘ E, Tolun R, Krespi Y. Vascular ischemia as a cause of transient global amnesia: a patient series. *Noro Psikiyatr Ars* 2015;52:59-63
8. Park KM, Han YH, Kim TH, Mun CW, Shin KJ, Ha SY, et al. Pre-existing structural abnormalities of the limbic system in transient global amnesia. *J Clin Neurosci* 2015;22:843-847
9. Döhring J, Schmuck A, Bartsch T. Stress-related factors in the emergence of transient global amnesia with hippocampal lesions. *Front Behav Neurosci* 2014;8:287
10. Kim J, Kwon Y, Yang Y, Jang IM, Chang Y, Park YH, et al. Clinical experience of modified diffusion-weighted imaging protocol for lesion detection in transient global amnesia: an 8-year large-scale clinical study. *J Neuroimaging* 2014;24:331-337
11. Scheel M, Malkowsky C, Klingebiel R, Schreiber SJ, Bohner G. Magnetic resonance imaging in transient global amnesia: lessons learned from 198 cases. *Clin Neuroradiol* 2012;22:335-340
12. Uttner I, Prexl S, Freund W, Unrath A, Bengel D, Huber R. Long-term outcome in transient global amnesia patients with and without focal hyperintensities in the CA1 region of the hippocampus. *Eur Neurol* 2012;67:155-160
13. Ahn S, Kim W, Lee YS, Kim WY, Lee JH, Oh BJ, et al. Transient global amnesia: seven years of experience with diffusion weighted imaging in an emergency department. *Eur Neurol* 2011;65:123-128
14. Auyeung M, Tsoi TH, Cheung CM, Fong DY, Li R, Chan JK, et al. Association of diffusion weighted imaging abnormalities and recurrence in transient global amnesia. *J Clin Neurosci* 2011;18:531-534
15. Ueno H, Naka H, Ohshita T, Wakabayashi S, Matsumoto M. Serial changes in delayed focal hippocampal lesions in patients with transient global amnesia. *Hiroshima J Med Sci* 2010;59:77-81
16. Lee SY, Kim WJ, Suh SH, Oh SH, Lee KY. Higher lesion detection by 3.0T MRI in patient with transient global amnesia. *Yonsei Med* J 2009;50:211-214
17. Alberici E, Pichiecchio A, Caverzasi E, Farina LM, Persico A, Cavallini A, et al. Transient global amnesia: hippocampal magnetic resonance imaging abnormalities. *Funct Neurol* 2008;23:149-152
18. Enzinger C, Thimary F, Kapeller P, Ropele S, Schmidt R, Ebner F, et al. Transient global amnesia: diffusion-weighted imaging lesions and cerebrovascular disease. *Stroke* 2008;39:2219-2225
19. Winbeck K, Etgen T, von Einsiedel HG, Röttinger M, Sander D. DWI in transient global amnesia and TIA: proposal for an ischaemic origin of TGA. *J Neurol Neurosurg Psychiatry* 2005;76:438-441
20. Sedlaczek O, Hirsch JG, Grips E, Peters CN, Gass A, Wöhrle J, et al. Detection of delayed focal MR changes in the lateral hippocampus in transient global amnesia. *Neurology* 2004;62:2165-2170
21. Huber R, Aschoff AJ, Ludolph AC, Riepe MW. Transient global amnesia. Evidence against vascular ischemic etiology from diffusion weighted imaging. *J Neurol* 2002;249:1520-1524
22. Strupp M, Brüning R, Wu RH, Deimling M, Reiser M, Brandt T. Diffusion-weighted MRI in transient global amnesia: elevated signal intensity in the left mesial temporal lobe in 7 of 10 patients. *Ann Neurol* 1998;43:164-170