PEER REVIEW HISTORY

BMJ Open publishes all reviews undertaken for accepted manuscripts. Reviewers are asked to complete a checklist review form (**http://bmjopen.bmj.com/site/about/resources/checklist.pdf**) and are provided with free text boxes to elaborate on their assessment. These free text comments are reproduced below.

ARTICLE DETAILS

TITLE (PROVISIONAL)	Neoadjuvant therapy or upfront surgery in advanced endometrial cancer: A systematic review protocol
AUTHORS	McCarthy, Amy; Balfour, Katharine; El Sayed, Iman; Edmondson, Richard; Wan, Yee-Loi

VERSION 1 – REVIEW

REVIEWER	John, Denny Amrita Institute of Medical Sciences and Research Centre
REVIEW RETURNED	03-Sep-2021

GENERAL COMMENTS	The manuscript is acceptable in the current form.

REVIEWER	Rundle, Stuart Gateshead Health, Northern Gynaecological Oncology Cente
REVIEW RETURNED	09-Sep-2021

GENERAL COMMENTS	Last paragraph of intro: ...sequence ofof treatment in advnaced endometrial cancer by providing... In comparator groups would it be reasonable to consider cytoreductive surgery alone or followed by hormonal treatment as well as adjuvant cytotoxic chemotherpy? No other comments

VERSION 1 – AUTHOR RESPONSE

Many thanks for the comments from the reviewers

In response to reviewer 2, we have corrected the last paragraph of the intro to read 'In doing so we hope to examine the case for undertaking a randomised trial of the optimal sequence of treatment in advanced endometrial by providing an evaluation of its effectiveness and a summary of the limitations of the available studies in this area.'

We welcome the reviewer's suggestions for considering other comparator groups but feel that as neither cytoreductive surgery alone nor followed by hormonal treatment are within current clinical practice guidelines for advanced endometrial cancers that these will not add to the impact of the study. We recognise these regimens may be used in individualized circumstances for which there may be case reports and anecdotal evidence but feel there is insufficient support amongst the clinical community for these approaches to warrant asking this in our comparisons.