Supplementary file 9 Table S5 Subgroup analysis of outcomes except renal failure events.

Subgroup	No. of trials	n	RR/MD/SMD (95%CI)	*P* for RR/MD/SMD	I^2	*P* for heterogeneity test[a]
Change in eGFR（mL/min/1.73m²）						
Study type						
RCT	5	881	-0.75 (-7.58, 6.08)	0.83	96.2%	0.28
Cohort studies	7	2406	2.48 (0.32, 4.64)	0.02	98.1%	
Baseline eGFR (mL/min/1.73m²)						
60-89	4	804	-1.00 (-7.41, 5.42)	0.76	99.2%	0.48
30-59	6	2109	2.58 (-0.44, 5.61)	0.10	95.4%	
<15	2	374	1.24 (-0.88, 3.37)	0.25	66.2%	
Mean age (years)						
<58	5	106	0.44 (-4.40, 5.27)	0.86	98.5%	0.70
≥58	6	2669	1.68 (-1.83, 5.18)	0.35	97.8%	
Baseline BP (mmHg)						
<140/90	6	2420	-0.52 (-4.36, 3.32)	0.32	97.8%	0.41
SBP≥140 或 DBP≥90	4	469	2.54 (-2.47, 7.55)	0.55	70.1%	
Comparator of control group						
Usual care or usual diet	6	1052	-0.38 (-5.14, 4.39)	0.88	95.8%	0.85
Different levels of sodium intake	6	2235	2.62 (0.31, 4.94)	0.03	98.2%	
Proteinuria or Albuminuria						
Study type						
RCT	7	997	-1.03 (-2.28, 0.21)	0.11	98.4%	0.45
Cohort studies	6	1925	-0.28 (-0.59, 0.02)	0.07	80.8%	
Baseline eGFR (mL/min/1.73m²)						
60-89	7	821	-1.06 (-2.32, 0.20)	0.10	98.2%	0.42
30-59	6	1201	-0.23 (-0.50, 0.04)	0.10	79.6%	
Mean age (years)						
<58	7	846	-1.12 (-2.35, 0.12)	0.08	98.2%	0.34
≥58	6	2076	-0.13 (-0.34, 0.09)	0.24	64.2%	
Baseline BP (mmHg)						
<140/90	8	200	-0.82 (-1.55, -0.10)	0.01	98.3%	0.65
SBP≥140 或 DBP≥90	4	2665	-0.39 (-0.68, -0.11)	0.03	0.0%	
Comparator of control group						
Usual care or usual diet	7	1083	-1.01 (-2.18, 0.15)	0.09	98.4%	0.12
Different levels of sodium intake	6	1839	-0.34 (-0.69, 0.00)	0.05	81.7%	
All-cause mortality						
Study type						
RCT	2	539	0.96 (0.31, 3.02)	0.95	0.0%	0.99
Cohort studies	7	95458	0.90 (0.50, 1.59)	0.71	98.5%	
Baseline eGFR (mL/min/1.73m²)						
30-59	4	5438	0.65 (0.41, 1.02)	0.06	85.2%	0.008
15-29	2	2126	1.14 (0.88, 1.48)	0.33	0.0%	
<15	3	88433	1.34 (0.41, 4.39)	0.62	43.4%	

Mean age (years)						
<58	3	704	0.61 (0.37, 1.00)	0.05	90.1%	0.08
≥58	6	90977	1.34 (0.76, 2.37)	0.32	84.2%	
Baseline BP (mmHg)						
<140/90	5	5438	0.64 (0.41, 0.99)	0.05	80.6%	0.86
SBP≥140 或 DBP≥90	1	121	0.51 (0.05, 5.46)	0.58	-	
Comparator of control group						
Usual care or usual diet	2	539	0.96 (0.31, 3.02)	0.95	0.0%	0.90
Different levels of sodium intake	7	95458	0.90 (0.50, 1.59)	0.71	98.5%	
CV events						
Study type						
RCT	1	418	0.47 (0.09, 2.53)	0.88	-	0.51
Cohort studies	4	93155	1.05 (0.51, 2.18)	0.89	98.2%	
Baseline eGFR (mL/min/1.73m^2)						
30-59	3	5352	0.78 (0.63, 0.95)	0.02	57.1%	**0.02**
<15	2	88221	1.50 (0.39, 5.85)	0.56	58.3%	
Mean age (years)						
<58	2	3863	0.71 (0.62, 0.80)	<0.001	0.0%	0.40
≥58	3	89710	1.20 (0.49, 2.91)	0.69	97.3%	
Baseline BP (mmHg)						
<140/90	2	4175	0.71 (0.62, 0.80)	<0.001	0.0%	0.62
SBP≥140 或 DBP≥90	2	1283	0.88 (0.74, 1.05)	0.17	0.0%	
Comparator of control group						
Usual care or usual diet	2	524	0.49 (0.14, 1.73)	0.27	0.0%	0.52
Different levels of sodium intake	3	93049	1.14 (0.53, 2.46)	0.74	98.8%	
SBP（mmHg）						
Study type						
RCT	11	1246	-3.54 (-6.40, -0.68)	0.02	65.6%	0.43
Cohort studies	8	2362	-7.22 (-11.69, -2.76)	0.002	93.6%	
Baseline eGFR (mL/min/1.73m^2)						
60-89	7	1017	-3.30 (-5.26, -1.34)	0.001	31.7%	0.64
30-59	7	2210	-6.10 (-11.62, -0.59)	0.03	92.8%	
< 15	4	349	-5.66 (-11.62, 0.30)	0.06	76.0%	
Mean age (years)						
< 58	12	1215	-7.45 (-11.76, -3.14)	0.001	85.6%	0.25
≥ 58	7	2393	-2.39 (-5.65, 0.87)	0.15	87.3%	
Baseline BP (mmHg)						
< 140/90	10	2786	-0.97 (-2.75, 0.81)	0.28	46.9%	**0.02**
SBP ≥ 140 or DBP ≥ 90	9	822	-10.24 (-15.37, -5.12)	< 0.001	88.4%	
Comparator of control group						
Usual care or usual diet	13	1546	-4.28 (-7.20, -1.36)	0.004	74.3%	0.05
Different levels of sodium intake	6	2062	-7.55 (-12.83, -2.28)	0.005	94.6%	
DBP（mmHg）						
Study type						
RCT	7	1246	-2.41 (-4.04, -0.77)	0.004	66.0%	0.85
Cohort studies	11	2278	-2.13 (-3.90, -0.37)	0.02	74.6%	

Baseline eGFR (mL/min/1.73m²)						
30-59	7	1017	-1.74 (-3.40, -0.08)	0.04	56.7%	0.13
15-29	6	2126	-0.53 (-1.25, 0.20)	0.16	11.9%	
< 15	4	349	-4.94 (-7.08, -2.80)	<0.001	23.2%	
Mean age (years)						
< 58	11	1131	-2.78 (-4.35, -1.22)	<0.001	50.8%	0.34
≥ 58	7	2393	-0.86 (-1.77, 0.04)	0.06	62.2%	
Baseline BP (mmHg)						
< 140/90	9	2747	-0.94 (-1.90, 0.02)	0.056	5.6%	**0.04**
SBP ≥ 140 或 DBP ≥ 90	9	777	-4.05 (-6.61, -1.49)	0.002	0.2%	
Comparator of control group						
Usual care or usual diet	11	1246	-2.41 (-4.04, -0.77)	0.004	66.0%	0.12
Different levels of sodium intake	7	2278	-2.13 (-3.90, -0.37)	0.02	74.6%	

Note: [a] *P* value calculated by meta regression. BP, blood pressure; CI, confidence interval; CV, cardiovascular; DBP, diastolic blood pressure; eGFR, estimated glomerular filtration rate; MD, mean difference; MAP, mean arterial pressure; n, number of patients; RCT, randomised parallel-group controlled trial; RR, relative risk; SMD, standardized mean difference; SBP, systolic blood pressure.