Supplementary file 12 Table S6 Sensitivity analysis of outcomes.

Sensitivity analysis	Renal composite outcomes			The rate of change in eGFR			Proteinuria or albuminuria			All-cause mortaity		
	No./n	RR (95% CI)	τ^2	No./n	MD (95%CI)	τ^2	No./n	MD/SMD (95%CI)	τ^2	No./n	RR (95% CI)	τ^2
Omit studies of follow-up time <12 months	-	-	-	7/2740	1.60 (-1.10, 4.31)	12.78	-	-	-	8/95876	0.92 (0.53, 1.59)	0.48
Omit studies of sample size <50	-	-	-	10/3206	1.07 (-1.26, 3.41)	12.86	9/2787	-0.89 (-1.56, -0.23)	0.99	-	-	-
Omit studies with low quality	-	-	-	-	-	-	-	-	-	-		
Omit studies with extreme outliers	-	-	-	11/3042	**2.43 (0.63, 4.23)**	8.01	12/2677	-0.24 (-0.43, -0.05)	0.06	8/7882	0.75 (0.56, 1.00)	0.08
Different statistical estimators												
DL	10/9578	0.71 (0.57, 0.88)	0.09	12/3287	1.18 (-0.94, 3.29)	12.57	13/2922	-0.70 (-1.24, -0.1)	0.89	9/95997	0.90 (0.52, 1.53)	0.48
EB	10/9578	0.72 (0.58, 0.89)	0.06	12/3287	1.16 (-2.02, 4.33)	23.4	13/2922	**-0.71(-1.66, 0.24)**	2.39	9/95997	0.92 (0.53, 1.46)	0.23
REML	10/9578	0.72 (0.58, 0.89)	0.03	12/3287	1.16 (-2.04, 4.35)	23.73	13/2922	**-0.71(-1.66, 0.24)**	2.38	9/95997	0.91 (0.53, 1.55)	0.33

Sensitivity analysis	CV events			SBP			DBP		
	No./n	RR (95% CI)	τ^2	No./n	MD (95%CI)	τ^2	No./n	MD (95%CI)	τ^2
Omit studies of follow-up time <12 months	4/93467	1.03 (0.50, 2.12)	0.45	-	-	-	-	-	-
Omit studies of sample size <50	-	-	-	13/3402	-4.05 (-6.93, -1.17)	22.57	12/3318	-1.45 (-2.34, -0.56)	1.05
Omit studies with low quality	-	-	-	-	-	-	-	-	
Different statistical estimators									
DL	5/93573	0.97 (0.49, 1.93)	0.45	19/3608	-5.34 (-7.93, -2.75)	22.80	15/3524	-2.00(-2.93, -1.06)	1.56
EB	5/93573	1.01 (0.46, 2.22)	0.28	19/3608	-5.81(-9.63, -1.99)	14.19	15/3524	-2.30(-3.61, -0.98)	3.78
REML	5/93573	0.99 (0.41, 2.37)	0.35	19/3608	-5.78(-9.59, -1.97)	52.37	15/3524	-2.27(-3.57, -0.96)	3.41

Note: CI, confidence interval; CV, cardiovascular; DL, DerSimonian-Laird; EB, empirical Bayes; DBP, diastolic blood pressure; eGFR, estimated glomerular filtration rate; MD, mean difference; MAP, mean arterial pressure; n, number of patients; No., number of trials; NOS, Newcastle-Ottawa Scale; REML, restricted maximum likelihood; RR, relative risk; SMD, standardized mean difference; SBP, systolic blood pressure.