

Supplementary file 6: Forest plots

Comparison 1: Integrated models of care vs. usual care
Outcome: Mortality

Study or Subgroup	log[Risk Ratio]	SE	Weight	Risk Ratio IV, Fixed, 95% CI
Havlir 2019	-0.1054	0.0665	100.0%	0.90 [0.79, 1.03]
Total (95% CI)			**100.0%**	**0.90 [0.79, 1.03]**


Heterogeneity: Not applicable
Test for overall effect: Z = 1.58 (P = 0.11)





Risk of bias legend
(A) Random sequence generation (selection bias)
(B) Allocation concealment (selection bias)
(C) Baseline outcome measurements similar
(D) Baseline characteristics similar
(E) Incomplete outcome data (attrition bias)
(F) Blinding of participants and personnel (performance bias)
(G) Blinding of outcome assessment (detection bias)
(H) Protection against contamination
(I) Selective reporting (reporting bias)
(J) Recruitment bias
(K) Baseline differences (clusters)
(L) Loss of clusters
(M) Incorrect analysis
(N) Compatibility with RCTs randomised by individuals
(O) Other bias


Comparison 1: Integrated models of care vs. usual care
Outcome: BP control



Footnotes
(1) Among people living with HIV (PLHIV)
(2) Among people living with HIV (PLHIV)

Risk of bias legend
(**A**) Random sequence generation (selection bias)
(**B**) Allocation concealment (selection bias)
(**C**) Baseline outcome measurements similar
(**D**) Baseline characteristics similar
(**E**) Incomplete outcome data (attrition bias)
(**F**) Blinding of participants and personnel (performance bias)
(**G**) Blinding of outcome assessment (detection bias)
(**H**) Protection against contamination
(**I**) Selective reporting (reporting bias)
(**J**) Recruitment bias
(**K**) Baseline differences (clusters)
(**L**) Loss of clusters
(**M**) Incorrect analysis
(**N**) Compatibility with RCTs randomised by individuals
(**O**) Other bias


Comparison 1: Integrated models of care vs. usual care
Outcome: BP and HIV control




Risk of bias legend
(A) Random sequence generation (selection bias)
(B) Allocation concealment (selection bias)
(C) Baseline outcome measurements similar
(D) Baseline characteristics similar
(E) Incomplete outcome data (attrition bias)
(F) Blinding of participants and personnel (performance bias)
(G) Blinding of outcome assessment (detection bias)
(H) Protection against contamination
(I) Selective reporting (reporting bias)
(J) Recruitment bias
(K) Baseline differences (clusters)
(L) Loss of clusters
(M) Incorrect analysis
(N) Compatibility with RCTs randomised by individuals
(O) Other bias

Comparison 1: Integrated models of care vs. usual care
Outcome: NCD control

Study or Subgroup	log[Risk Ratio]	SE	Risk Ratio IV, Fixed, 95% CI
1.4.1 Prevalent NCD at baseline			
Havlir 2019 (1)	0.0583	0.095	1.06 [0.88, 1.28]
1.4.2 Prevalent NCD at follow-up			
Havlir 2019 (2)	0.1222	0.0779	1.13 [0.97, 1.32]






Footnotes
(1) Among PLHIV
(2) Among PLHIV


Risk of bias legend
(A) Random sequence generation (selection bias)
(B) Allocation concealment (selection bias)
(C) Baseline outcome measurements similar
(D) Baseline characteristics similar
(E) Incomplete outcome data (attrition bias)
(F) Blinding of participants and personnel (performance bias)
(G) Blinding of outcome assessment (detection bias)
(H) Protection against contamination
(I) Selective reporting (reporting bias)
(J) Recruitment bias
(K) Baseline differences (clusters)
(L) Loss of clusters
(M) Incorrect analysis
(N) Compatibility with RCTs randomised by individuals
(O) Other bias

Comparison 1: Integrated models of care vs. usual care
Outcome: NCD and HIV control




Risk of bias legend
(A) Random sequence generation (selection bias)
(B) Allocation concealment (selection bias)
(C) Baseline outcome measurements similar
(D) Baseline characteristics similar
(E) Incomplete outcome data (attrition bias)
(F) Blinding of participants and personnel (performance bias)
(G) Blinding of outcome assessment (detection bias)
(H) Protection against contamination
(I) Selective reporting (reporting bias)
(J) Recruitment bias
(K) Baseline differences (clusters)
(L) Loss of clusters
(M) Incorrect analysis
(N) Compatibility with RCTs randomised by individuals
(O) Other bias

Comparison 2: Strategies to promote integrated models of care vs. usual care
Outcome: Mortality



Risk of bias legend
(A) Random sequence generation (selection bias)
(B) Allocation concealment (selection bias)
(C) Baseline outcome measurements similar
(D) Baseline characteristics similar
(E) Incomplete outcome data (attrition bias)
(F) Blinding of participants and personnel (performance bias)
(G) Blinding of outcome assessment (detection bias)
(H) Protection against contamination
(I) Selective reporting (reporting bias)
(J) Recruitment bias
(K) Baseline differences (clusters)
(L) Loss of clusters
(M) Incorrect analysis
(N) Compatibility with RCTs randomised by individuals
(O) Other bias

Comparison 2: Strategies to promote integrated models of care vs. usual care
Outcome: Depression



Footnotes

(1) Change from baseline to follow-up; 10-item Center for EpidemiologicStudies...

(2) Value at follow-up; Patient Health Questionnaire-9

Risk of bias legend

(**A**) Random sequence generation (selection bias)

(**B**) Allocation concealment (selection bias)

(**C**) Baseline outcome measurements similar

(**D**) Baseline characteristics similar

(**E**) Incomplete outcome data (attrition bias)

(**F**) Blinding of participants and personnel (performance bias)

(**G**) Blinding of outcome assessment (detection bias)

(**H**) Protection against contamination

(**I**) Selective reporting (reporting bias)

(**J**) Recruitment bias

(**K**) Baseline differences (clusters)

(**L**) Loss of clusters

(**M**) Incorrect analysis


(**N**) Compatibility with RCTs randomised by individuals

(**O**) Other bias

Comparison 2: Strategies to promote integrated models of care vs. usual care
Outcome: Quality of life

Study or Subgroup	Mean Difference	SE	Weight	Mean Difference IV, Fixed, 95% CI
Fairall 2016 (1)	6.06	4.7501	100.0%	6.06 [-3.25, 15.37]
Total (95% CI)			**100.0%**	**6.06 [-3.25, 15.37]**


Heterogeneity: Not applicable
Test for overall effect: Z = 1.28 (P = 0.20)



Footnotes
(1) Euro-Qol-5D visual analogue scale: 0=worst imaginable state of health,...


Risk of bias legend
(**A**) Random sequence generation (selection bias)
(**B**) Allocation concealment (selection bias)
(**C**) Baseline outcome measurements similar
(**D**) Baseline characteristics similar
(**E**) Incomplete outcome data (attrition bias)
(**F**) Blinding of participants and personnel (performance bias)
(**G**) Blinding of outcome assessment (detection bias)
(**H**) Protection against contamination
(**I**) Selective reporting (reporting bias)
(**J**) Recruitment bias
(**K**) Baseline differences (clusters)
(**L**) Loss of clusters
(**M**) Incorrect analysis
(**N**) Compatibility with RCTs randomised by individuals
(**O**) Other bias

Comparison 2: Strategies to promote integrated models of care vs. usual care
Outcome: Change in systolic BP




Risk of bias legend
(A) Random sequence generation (selection bias)
(B) Allocation concealment (selection bias)
(C) Baseline outcome measurements similar
(D) Baseline characteristics similar
(E) Incomplete outcome data (attrition bias)
(F) Blinding of participants and personnel (performance bias)
(G) Blinding of outcome assessment (detection bias)
(H) Protection against contamination
(I) Selective reporting (reporting bias)
(J) Recruitment bias
(K) Baseline differences (clusters)
(L) Loss of clusters
(M) Incorrect analysis
(N) Compatibility with RCTs randomised by individuals
(O) Other bias

Comparison 2: Strategies to promote integrated models of care vs. usual care
Outcome: Change in HbA1c



Risk of bias legend
(A) Random sequence generation (selection bias)
(B) Allocation concealment (selection bias)
(C) Baseline outcome measurements similar
(D) Baseline characteristics similar
(E) Incomplete outcome data (attrition bias)
(F) Blinding of participants and personnel (performance bias)
(G) Blinding of outcome assessment (detection bias)
(H) Protection against contamination
(I) Selective reporting (reporting bias)
(J) Recruitment bias
(K) Baseline differences (clusters)
(L) Loss of clusters
(M) Incorrect analysis
(N) Compatibility with RCTs randomised by individuals
(O) Other bias

Comparison 2: Strategies to promote integrated models of care vs. usual care
Outcome: Change in total cholesterol



Risk of bias legend
(A) Random sequence generation (selection bias)
(B) Allocation concealment (selection bias)
(C) Baseline outcome measurements similar
(D) Baseline characteristics similar
(E) Incomplete outcome data (attrition bias)
(F) Blinding of participants and personnel (performance bias)
(G) Blinding of outcome assessment (detection bias)
(H) Protection against contamination
(I) Selective reporting (reporting bias)
(J) Recruitment bias
(K) Baseline differences (clusters)
(L) Loss of clusters
(M) Incorrect analysis
(N) Compatibility with RCTs randomised by individuals
(O) Other bias