Supplement Table 2: Findings of studies examining the association between occupational sitting and all-cause mortality[1]

Nr.	Study	Sample	Major occupational groups	Follow-up length/ duration	Assessment of occupational sitting (as analysed)	Outcomes	Findings[2]	Study quality[3]	Sex-stratified analysis
	First author, year, country, cohort	n (% female) age at baseline	General population or specification		Categories of measurement and standardized instrument (if applicable)	(Data basis)	Statistical results and reference categories (e.g. risk estimates with 95 % confidence interval)	(++) (+) (0)	Yes/no
1.	Chau 2015 Norway HUNT3	n=36817 (45 % F); (for occupational sitting) 18–>60 y	General population (incl. working)	(2006–2008) mean 3.3 y	Self-reported OPA in 4 categories: 'work that mostly involves sitting'; 'work that requires much walking', 'work that requires much walking and lifting'; 'heavy physical labour'	All-cause mortality (death register)	Reference: mostly sitting much walk and lift: HR=0.65 (0.44–0.97) **Compared to mostly sitting, those with jobs with much walking/lifting had 35 % lower risk of all-cause mortality.** much walking: HR=0.73 (0.51–1.03) heavy labour: HR=0.95 (0.64–1.40) **No difference in risk of mortality, when mostly sitting OPA was compared with more OPA.**	++	No
2.	Pulsford 2015 Great Britain Whitehall II cohort study	n=5132; (27.5 % F); mean age low sitting: 46.6 y; mean age high sitting: 41.5 y	Clerical and office support, executive, senior administra-tives grades	Mean 17.7 y (+/-2.2 y)	Self-reported work sitting time/week summed up into 4 categories: ≥0–<8 h/wk; ≥8–<25 h/wk; ≥25–<40 h/wk; ≥40 h/wk	All-cause mortality (national register)	Reference: ≥0–<8 h/wk work sitting time work sitting ≥8–<25 h/wk HR=0.93 (0.73–1.19) work sitting ≥25–<40 h/wk HR=0.83 (0.66–1.25) work sitting ≥40 h/wk HR=0.81 (0.57–1.14) **No associations between any work sitting time category (h/wk) and all-cause mortality.**	++	No
3.	Sakaue 2018 Japan Seven Countries Study	n=1680 (58.8 % F); 40–95 y mean age: 61.1 y	General population (incl. working)	Mean 15.9 y (+/-3.8 y)	Baecke physical activity questionnaire: self-reported work sitting time on a five-point Likert scale: never, seldom, sometimes, often, always	All-cause mortality (obituaries, medical records, registries, and other)	Index per unit HR=1.11 (1.01–1.22)[4] **Occupational sitting time was a risk factor for all-cause death.** M: HR=1.28 (1.13–1.46) F: HR=1.02 (0.88–1.17) **Stratified by sex, occupational sitting time was significantly associated with all-cause death in males of, but not in females.**	++	Yes
4.	van der Ploeg 2015 Denmark Danish Work Environment Cohort Study (DWECS)	2000: n=5926 (47.3 % F); 2005: n=8769 (50.9 % F); 2010: n=10624 (52.4 % F); 21–≥70 y	Working population	(1990–2010) mean 12.61 y	Self-reported occupational sitting time, analysed in 2 categories work sitting time <24 h/wk and ≥24 h/wk	All-cause mortality (death register)	Reference: <24 h/wk work sitting ≥24 h/wk work sitting: HR=1.25 (0.90–1.74) *Occ. sitting time per 10 h/wk increase:* HR=1.09 (0.97–1.23) **No statistically significant association between occupational sitting time and all-cause mortality**	++	No
5.	Kikuchi 2015 Japan	n=36516; (44.4 % F); mean age: 58.7 y 50–74 y	Agriculture, forestry, fishing, salaried workers, home business, profess-ionals	(2000–2011) mean 10.1 y	Self-reported occupational sitting time analysed in 3 categories: short <1 h/d; middle 1–<3 h/d; longer ≥3 h/d	All-cause mortality (death certificates/ national register)	Reference: short work sitting <1 h/d[4] M longer work sitting ≥3 h/d HR=0.96 (0.86–1.09) F longer work sitting ≥3 h/d HR=1.15 (0.92–1.43) **No association between occupational sitting duration and mortality was found neither for the overall sample or stratified by gender:** PI M longer work sitting ≥3 h/d HR=1.23 (1.00–1.51) PI F longer work sitting ≥3 h/d HR=1.34 (0.97–1.84) **Among male workers in primary industry, longer duration of occupational sitting time was borderline significantly associated with higher mortality.**	+	Yes

Nr.	Study	Sample	Major occupational groups	Follow-up length/ duration	Assessment of occupational sitting (as analysed)	Outcomes	Findings[2]	Study quality[3]	Sex-stratified analysis
	First author, year, country, cohort	n (% female) age at baseline	General population or specification		Categories of measurement and standardized instrument (if applicable)	(Data basis)	Statistical results and reference categories (e.g. risk estimates with 95 % confidence interval)	(++) (+) (0)	Yes/no
6.	Kim 2013 USA Multiethnic Cohort Study (MEC)	n=134596 (54.4 % F); 45–75 y	General population (incl. working)	13.7 y median	Self-reported work sitting time h/d analysed in 3 categories: <1 h/d; 1–4 h/d; ≥5 h/d	All-cause mortality (death certificates/ national register)	Reference: work sitting <1 h/d M long work sitting ≥5 h/d HR=0.94 (0.88–1.02) F long work sitting ≥5 h/d HR=0.98 (0.90–1.08) **Sitting at work not associated with risk of all-cause mortality**	+	Yes
7.	Moe 2013 Norway HUNT2	n=37300; (49.2 % F); mean 44.1 y	General population (incl. working)	(1995–2008) 12.4 y median	Self-reported OPA, analysed in 3 categories: 'mostly sedentary', 'much walking/lifting', 'heavy physical work'	All-cause and cardiovascular mortality (death register)	Reference: much walking/lifting without MS mostly sedentary HR=0.97 (0.83–1.14) **No associations between OPA and all-cause mortality for participants without metabolic syndrome (MS), but different associations for persons with MS:** mostly sedentary with MS HR=1.63 (1.31–2.02)	+	No
8.	Stamatakis 2013 Great Britain Health Survey for England (HSE) Scottish Health Survey (SHS)	n=10834 (48.1 % F); >40 y	General population (incl. working)	Mean 12.9 y (+/-3.3 y)	Self-reported predominant OPA categorized into predominant sitting activity at work versus predominant standing/walking activity at work	All-cause mortality (national register)	Reference: sitting OPA F standing/walking OPA HR=0.68 (0.52–0.89) M standing/walking OPA HR=0.97 (0.78–1.19) **In women, a standing/walking occupation was associated with a lower risk of all-cause mortality by 32 % compared to sitting occupations (reference), but not in men.**	+	Yes

Legend:
n (% F): number of participants (percent female)
Female=F; Male=M
OPA: occupational physical activity
PI: primary industry
1 In alphabetical order and after methodological assessment rating (high to low quality)
2 In model adjusted for age, sex, leisure time physical activity (LPTA), health behaviour (alcohol, smoking, diet), and socioeconomic status (education, household income) if available, unless reported otherwise
3 Methodological Assessment incl. risk of bias: assessment after the SIGN-checklists resulting in overall ratings of 'high quality' (++), 'acceptable quality' (+) and 'unacceptable/low quality' (0), synonymic with 'high quality' (++) / low risk of bias, 'acceptable quality' (+) / accepable risk of bias, and 'unacceptable/low quality' (0) / unacceptable risk of bias.
4 Not adjusted for socioeconomic status