Supplemental Table 3: Reasons for Exclusion of 90 Articles

#	Article (Year)	Full Citation	Inclusion criteria not fulfilled or other reasons for exclusion
1.	Dammaschke (2003)	Dammaschke T, Steven D, Kaup M, Ott KH. Long-term survival of root-canal-treated teeth: a retrospective study over 10 years. *J Endod*. 2003;29(10):638-643.	[5]Success based on clinical and/ or radiographic (strict, absence of apical radiolucency; loose, reduction in size of radiolucency) criteria
2.	Fouad (2003)	Fouad AF, Burleson J. The effect of diabetes mellitus on endodontic treatment outcome: data from an electronic patient record. *J Am Dent Assoc*. 2003;134(1):43-118.	[2]Stratified analysis of primary root canal therapy available, if re-treatment cases had been included
3.	Tani-Ishii (2003)	Tani-Ishii N, Teranaka T. Clinical and radiographic evaluation of root-canal obturation with obtura II. *J Endod*. 2003;29(11):739-742.	[7]Other- Clinical Protocol, not standard endodontic procedure
4.	Travassos (2003)	Travassos RM, Caldas Ade F, de Albuquerque DS. Cohort study of endodontic therapy success. *Braz Dent J*. 2003;14(2):109-113.	[7]Other- Excluded due to quality of the study
5.	Field (2004)	Field JW, Gutmann JL, Solomon ES, Rakusin H. A clinical radiographic retrospective assessment of the success rate of single-visit root canal treatment. *Int Endod J*. 2004;37(1):70-82.	[4] At least 12 months post-operative review
6.	Salehrabi (2004)	Salehrabi R, Rotstein I. Endodontic treatment outcomes in a large patient population in the USA: an epidemiological study. *J Endod*. 2004;30(12):846-850.	[5]Success based on clinical and/ or radiographic (strict, absence of apical radiolucency; loose, reduction in size of radiolucency) criteria.
7.	Tilashalski (2004)	Tilashalski KR, Gilbert GH, Boykin MJ, Shelton BJ. Root canal treatment in a population-based adult sample: status of teeth after endodontic treatment. *J Endod*. 2004;30(8):577-581.	[5]Success based on clinical and/ or radiographic (strict, absence of apical radiolucency; loose, reduction in size of radiolucency) criteria.
8.	Peters (2004)	Peters OA, Barbakow F, Peters CI. An analysis of endodontic treatment with three nickel-titanium rotary root canal preparation techniques. *Int Endod J*. 2004;37(12):849-859.	[2]Stratified analysis of primary root canal therapy available, if re-treatment cases had been included
9.	Marending (2005)	Marending M, Peters OA, Zehnder M. Factors affecting the outcome of orthograde root canal therapy in a general dentistry hospital practice. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 2005;99(1):119-124.	[6]Overall success rates given or could be calculated from the raw data at the tooth level.

10.	Stoll (2005)	Stoll R, Betke K, Stachniss V. The influence of different factors on the survival of root canal fillings: a 10-year retrospective study. *J Endod*. 2005;31(11):783-790.	[5]Success based on clinical and/ or radiographic (strict, absence of apical radiolucency; loose, reduction in size of radiolucency) criteria.
11.	Waltimo (2005)	Waltimo T, Trope M, Haapasalo M, Ørstavik D. Clinical efficacy of treatment procedures in endodontic infection control and one-year follow-up of periapical healing. *J Endod*. 2005;31(12):863-866.	[5]Success based on clinical and/ or radiographic (strict, absence of apical radiolucency; loose, reduction in size of radiolucency) criteria.
12.	DeQuadros (2005)	De Quadros I, Gomes BP, Zaia AA, Ferraz CC, Souza-Filho FJ. Evaluation of endodontic treatments performed by students in a Brazilian Dental School. *J Dent Educ*. 2005;69(10):1161-1170.	[2]Stratified analysis of primary root canal therapy available, if re-treatment cases had been included
13.	Quesnell (2005)	Quesnell BT, Alves M, Hawkinson RW Jr, Johnson BR, Wenckus CS, BeGole EA. The effect of human immunodeficiency virus on endodontic treatment outcome. *J Endod*. 2005;31(9):633-636.	[6]Overall success rates given or could be calculated from the raw data at the tooth level
14.	Gesi (2006)	Gesi A, Hakeberg M, Warfvinge J, Bergenholtz G. Incidence of periapical lesions and clinical symptoms after pulpectomy--a clinical and radiographic evaluation of 1- versus 2-session treatment. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 2006;101(3):379-388.	[6]Overall success rates given or could be calculated from the raw data at the tooth level.
15.	Shetty (2006)	Shetty K, Garcia J, Leigh J. Success of root canal therapy in HIV-positive patients. *Gen Dent*. 2006;54(6):397-402.	[4] At least 12 months post-operative review
16.	Suchina (2006)	Suchina JA, Levine D, Flaitz CM, Nichols CM, Hicks MJ. Retrospective clinical and radiologic evaluation of nonsurgical endodontic treatment in human immunodeficiency virus (HIV) infection. *J Contemp Dent Pract*. 2006;7(1):1-8.	[5]Success based on clinical and/ or radiographic (strict, absence of apical radiolucency; loose, reduction in size of radiolucency) criteria; [6] Overall success rate given or could be calculated from the raw data at the tooth level.
17.	Chen (2007)	Chen SC, Chueh LH, Hsiao CK, Tsai MY, Ho SC, Chiang CP. An epidemiologic study of tooth retention after nonsurgical endodontic treatment in a large population in Taiwan. *J Endod*. 2007;33(3):226-229.	[5]Success based on clinical and/ or radiographic (strict, absence of apical radiolucency; loose, reduction in size of radiolucency) criteria
18.	Chugal (2007)	Chugal NM, Clive JM, Spångberg LS. Endodontic treatment outcome: effect of the permanent restoration. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 2007;104(4):576-582.	[7]The data for this study comes from Chugal's 2001 study, which was already included in Ng's 2007 study

19.	Liang (2007)	Liang YH, Zhang G, Wang JD. Clinical Effects of Calcium Hydroxide-based sealer on chronic apical periodontitis. *Zhonghua Kou Qiang Yi Xue Za Zhi*. 2007;42(11):658-662.	[7] Other- cannot be translated into English
20.	Sari (2007)	Sari S, Duruturk L. Radiographic evaluation of periapical healing of permanent teeth with periapical lesions after extrusion of AH Plus sealer. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 2007;104(3):e54-e59.	[6]Overall success rate given or could be calculated from the raw data at the tooth level.
21.	Zmener (2007)	Zmener O, Pameijer CH. Clinical and radiographical evaluation of a resin-based root canal sealer: a 5-year follow-up. *J Endod*. 2007;33(6):676-679.	[7] Other- Same patient pool as Zmener 2004 study, 2004 study chosen due to higher recall rate
22.	Alley (2008)	Alley BS, Buchanan TH, Eleazer PD. Comparison of the success of root canal therapy in HIV/AIDS patients and non-infected controls. *Gen Dent*. 2008;56(2):155-157.	[5]Success based on clinical and/ or radiographic (strict, absence of apical radiolucency; loose, reduction in size of radiolucency) criteria.
23.	Chen (2008)	Chen SC, Chueh LH, Hsiao CK, Wu HP, Chiang CP. First untoward events and reasons for tooth extraction after nonsurgical endodontic treatment in Taiwan. *J Endod*. 2008;34(6):671-674.	[5]Success based on clinical and/ or radiographic (strict, absence of apical radiolucency; loose, reduction in size of radiolucency) criteria.
24.	deChevigny (2008)	de Chevigny C, Dao TT, Basrani BR, et al. Treatment outcome in endodontics: the Toronto study--phases 3 and 4: orthograde retreatment. *J Endod*. 2008;34(2):131-137.	[5]Success based on clinical and/ or radiographic (strict, absence of apical radiolucency; loose, reduction in size of radiolucency) criteria
25.	Hannahan (2008)	Hannahan JP, Eleazer PD. Comparison of success of implants versus endodontically treated teeth. *J Endod*. 2008;34(11):1302-1305.	[7] Other- Clinical Protocol not described
26.	Burke (2009)	Burke FM, Lynch CD, Ní Ríordáin R, Hannigan A. Technical quality of root canal fillings performed in a dental school and the associated retention of root-filled teeth: a clinical follow-up study over a 5-year period. *J Oral Rehabil*. 2009;36(7):508-515.	[5]Success based on clinical and/ or radiographic (strict, absence of apical radiolucency; loose, reduction in size of radiolucency) criteria
27.	Cheung (2009)	Cheung GS, Liu CS. A retrospective study of endodontic treatment outcome between nickel-titanium rotary and stainless-steel hand filing techniques. *J Endod*. 2009;35(7):938-943.	[4] At least 12 months post-operative review

28.	Fonzar (2009)	Fonzar F, Fonzar A, Buttolo P, Worthington HV, Esposito M. The prognosis of root canal therapy: a 10-year retrospective cohort study on 411 patients with 1175 endodontically treated teeth. *Eur J Oral Implantol*. 2009;2(3):201-208.	[2]Stratified analysis of primary root canal therapy available, if re-treatment cases had been included
29.	Hsiao (2009)	Hsiao A, Glickman G, He J. A retrospective clinical and radiographic study on healing of periradicular lesions in patients taking oral bisphosphonates. *J Endod*. 2009;35(11):1525-1528.	[4] At least 12 months post-operative review
30.	Suter (2009)	Suter B, Portmann P, Lussi A. A 5-year review of teeth filled with the noninstrumentation technology. *Int Endod J*. 2009;42(7):639-648.	[1]Clinical study on primary root canal therapy: randomized control trials, cohort studies, retrospective observational studies - Case Study/ Case series
31.	Mohan (2009)	Mohan SM, Kaushik SK. Root Canal Treatment Using Thermoplasticized Carrier Condensation Technique. *Med J Armed Forces India*. 2009;65(4):336-341.	[7]Other- Excluded due to quality of the study
32.	Asgary (2010)	Asgary S, Eghbal MJ. A clinical trial of pulpotomy vs. root canal therapy of mature molars *J Dent Res*. 2010;89(10):1080-1085.	[4] At least 12 months post-operative review
33.	Fleming (2010)	Fleming CH, Litaker MS, Alley LW, Eleazer PD. Comparison of classic endodontic techniques versus contemporary techniques on endodontic treatment success. *J Endod*. 2010;36(3):414-418.	[5]Success based on clinical and/ or radiographic (strict, absence of apical radiolucency; loose, reduction in size of radiolucency) criteria
34.	Gilbert (2010)	Gilbert GH, Tilashalski KR, Litaker MS, et al. Outcomes of root canal treatment in Dental Practice-Based Research Network practices. *Gen Dent*. 2010;58(1):28-36.	[7] Other- Clinical Protocol not described
35.	Moazami (2011)	Moazami F, Sahebi S, Sobhnamayan F, Alipour A. Success rate of nonsurgical endodontic treatment of nonvital teeth with variable periradicular lesions. *Iran Endod J*. 2011;6(3):119-124.	[2] Stratified analysis of primary root canal therapy available, if re-treatment cases had been included
36.	Ng (2011)	Ng YL, Mann V, Gulabivala K. A prospective study of the factors affecting outcomes of non-surgical root canal treatment: part 2: tooth survival. *Int Endod J*. 2011;44(7):610-625.	[5]Success based on clinical and/ or radiographic (strict, absence of apical radiolucency; loose, reduction in size of radiolucency) criteria

37.	Riccitiello (2011)	Riccitiello F, Stabile P, Amato M, Rengo S, D'Ambrosio C. The treatment of the large periradicular endodontic injury. *Minerva Stomatol*. 2011;60(9):417-426.	[1]Clinical study on primary root canal therapy: randomized control trials, cohort studies, retrospective observational studies
38.	Koral (2011)	Koral SM. Calcium oxide as a root filling material: a three-year prospective clinical outcome study. *Open Dent J*. 2011;5:13-17.	[7]Other- Clinical Protocol, not standard endodontic procedure
39.	Liang (2011)	Liang YH, Li G, Wesselink PR, Wu MK. Endodontic outcome predictors identified with periapical radiographs and cone-beam computed tomography scans. *J Endod*. 2011;37(3):326-331.	[6]Overall success rates given or could be calculated from the raw data at the tooth level
40.	Bernstein (2012)	Bernstein SD, Horowitz AJ, Man M, et al. Outcomes of endodontic therapy in general practice: a study by the Practitioners Engaged in Applied Research and Learning Network. *J Am Dent Assoc*. 2012;143(5):478-487.	[7] Other- Clinical Protocol not described
41.	Lee (2012)	Lee AH, Cheung GS, Wong MC. Long-term outcome of primary non-surgical root canal treatment. *Clin Oral Investig*. 2012;16(6):1607-1617.	[5]Success based on clinical and/ or radiographic (strict, absence of apical radiolucency; loose, reduction in size of radiolucency) criteria
42.	Tootla (2012)	Tootla S, Owen CP. A comparison of endodontic treatment outcomes between HIV-positive and HIV-negative patients. *SADJ*. 2012;67(7):322-325.	[5]Success based on clinical and/ or radiographic (strict, absence of apical radiolucency; loose, reduction in size of radiolucency) criteria
43.	Liang (2012)	Liang YH, Li G, Shemesh H, Wesselink PR, Wu MK. The association between complete absence of post-treatment periapical lesion and quality of root canal filling. *Clin Oral Investig*. 2012;16(6):1619-1626.	[6]Overall success rates given or could be calculated from the raw data at the tooth level
44.	Patel (2012)	Patel S, Wilson R, Dawood A, Foschi F, Mannocci F. The detection of periapical pathosis using digital periapical radiography and cone beam computed tomography - part 2: a 1-year post-treatment follow-up. *Int Endod J*. 2012;45(8):711-723.	[6]Overall success rates given or could be calculated from the raw data at the tooth level
45.	Zmener (2012)	Zmener O, Pameijer CH. Clinical and radiographic evaluation of a resin-based root canal sealer: 10-year recall data. *Int J Dent*. 2012;2012:763248.	[7] Other- Same patient pool as Zmener 2004 study, 2004 study chosen due to higher recall rate
46.	Abbaszadegan (2013)	Abbaszadegan A, Nabavizadeh M, Hoseini Yekani A, Khayat A. Comparison of Endodontic Treatment Results Yielded from Using Normal Saline with IKI Final Rinse or	[6]Overall success rates given or could be calculated from the raw data at the tooth level.

		NaOCl Irrigation: A 30-Month Follow-up Study. *Iran Endod J*. 2013;8(4):171-176.	
47.	Castelot-Enkel (2013)	Castelot-Enkel B, Nguyen JM, Armengol V, et al. A recall program for the outcome of conventional root canal treatment performed in a teaching hospital. *Acta Odontol Scand*. 2013;71(6):1399-1409.	[5]Success based on clinical and/ or radiographic (strict, absence of apical radiolucency; loose, reduction in size of radiolucency) criteria
48.	Fernandez (2013)	Fernández R, Cadavid D, Zapata SM, Alvarez LG, Restrepo FA. Impact of three radiographic methods in the outcome of nonsurgical endodontic treatment: a five-year follow-up. *J Endod*. 2013;39(9):1097-1103.	[6]Overall success rate given or could be calculated from the raw data at the tooth level.
49.	Ilgüy (2013)	Ilgüy D, Ilgüy M, Fisekçioglu E, Ersan N, Tanalp J, Dölekoglu S. Assessment of root canal treatment outcomes performed by Turkish dental students: results after two years. *J Dent Educ*. 2013;77(4):502-509.	[5]Success based on clinical and/ or radiographic (strict, absence of apical radiolucency; loose, reduction in size of radiolucency) criteria
50.	Liang (2013)	Liang YH, Jiang LM, Jiang L, et al. Radiographic healing after a root canal treatment performed in single-rooted teeth with and without ultrasonic activation of the irrigant: a randomized controlled trial. *J Endod*. 2013;39(10):1218-1225.	[4] At least 12 months post-operative review
51.	Balto (2013)	Balto K. Trial suggests no difference between single-visit and two-visit root canal treatment. *Evid Based Dent*. 2013;14(2):48.	[1]Clinical study on primary root canal therapy: randomized control trials, cohort studies, retrospective observational studies
52.	Chen (2014)	Chen ZH. Application of sonic endo-instruments in combination with Mtwo Ni-Ti files in root canal preparation of molars.Shanghai Journal of Stomatology.2014;23(5):624-627.	[7] Other- cannot be translated into English
53.	Cousson (2014)	Cousson PY, Nicolas E, Hennequin M. A follow-up study of pulpotomies and root canal treatments performed under general anaesthesia. *Clin Oral Investig*. 2014;18(4):1155-1163.	[2]Stratified analysis of primary root canal therapy available, if other treatments had been included
54.	Edionwe (2014)	Edionwe JI, Shaba OP, Umesi DC. Single visit root canal treatment: a prospective study. *Niger J Clin Pract*. 2014;17(3):276-281.	[4] At least 12 months post-operative review
55.	Estrela (2014)	Estrela C, Silva JA, Decurcio DA, et al. Monitoring nonsurgical and surgical root canal treatment of teeth	[4] At least 12 months post-operative review

		with primary and secondary infections. *Braz Dent J*. 2014;25(6):494-501.	
56.	Ferreira (2014)	Ferreira MM, Carrilho E, Carrilho F. Diabetes mellitus and its influence on the success of endodontic treatment: a retrospective clinical study. *Acta Med Port*. 2014;27(1):15-22.	[7] Other- cannot be translated into English
57.	Jordal (2014)	Jordal K, Valen A, Örstavik D. Periapical status of root-filled teeth in Norwegian children and adolescents. *Acta Odontol Scand*. 2014;72(8):801-805.	[7] Other- Clinical Protocol not described
58.	Jordan (2014)	Jordan RA, Holzner AL, Markovic L, Brueckner I, Zimmer S. Clinical effectiveness of basic root canal treatment after 24 months: a randomized controlled trial. *J Endod*. 2014;40(4):465-470.	[5]Success based on clinical and/ or radiographic (strict, absence of apical radiolucency; loose, reduction in size of radiolucency) criteria
59.	Lin (2014)	Lin PY, Huang SH, Chang HJ, Chi LY. The effect of rubber dam usage on the survival rate of teeth receiving initial root canal treatment: a nationwide population-based study. *J Endod*. 2014;40(11):1733-1737.	[5]Success based on clinical and/ or radiographic (strict, absence of apical radiolucency; loose, reduction in size of radiolucency) criteria
60.	Martins (2014)	Martins MR, Carvalho MF, Pina-Vaz I, Capelas JA, Martins MA, Gutknecht N. Outcome of Er,Cr:YSGG laser-assisted treatment of teeth with apical periodontitis: a blind randomized clinical trial. *Photomed Laser Surg*. 2014;32(1):3-9.	[6]Overall success rates given or could be calculated from the raw data at the tooth level.
61.	Touboul (2014)	Touboul V, Germa A, Lasfargues JJ, Bonte E. Outcome of endodontic treatments made by postgraduate students in the dental clinic of bretonneau hospital. *Int J Dent*. 2014;2014:684979.	[5]Success based on clinical and/ or radiographic (strict, absence of apical radiolucency; loose, reduction in size of radiolucency) criteria
62.	Menakaya (2015)	Menakaya IN, Adegbulugbe IC, Oderinu OH, Shaba OP. The Efficacy of Calcium Hydroxide Powder mixed with 0.2% Chlorhexidine Digluconate or mixed with Normal Saline as Intracanal Medicament in the Treatment of Apical Periodontitis. *J Contemp Dent Pract*. 2015;16(8):657-664.	[7]Other- Excluded due to quality of the study
63.	Raedel (2015)	Raedel M, Hartmann A, Bohm S, Walter MH. Three-year outcomes of root canal treatment: Mining an insurance database. *J Dent*. 2015;43(4):412-417.	[5]Success based on clinical and/ or radiographic (strict, absence of apical radiolucency; loose, reduction in size of radiolucency) criteria

64.	Saidi (2015)	Saidi A, Naaman A, Zogheib C. Accuracy of Cone-beam Computed Tomography and Periapical Radiography in Endodontically Treated Teeth Evaluation: A Five-Year Retrospective Study. *J Int Oral Health*. 2015;7(3):15-19.	[7] Other- Clinical Protocol not described
65.	Tang (2015)	Tang Z, Wang H, Jiang S. Clinical study of single-visit root canal treatment with a nickel-titanium (Ni-Ti) rotary instrument combined with different ultrasonic irrigation solutions for elderly patients with chronic apical periodontitis. *Biomed Mater Eng*. 2015;26 Suppl 1:S311-S318.	[6]Overall success rates given or could be calculated from the raw data at the tooth level.
66.	Yousuf (2015)	Yousuf W, Khan M, Sheikh A. Success Rate of Overfilled Root Canal Treatment. *J Ayub Med Coll Abbottabad*. 2015;27(4):780-783.	[7]Other- Study only included treatments with technical errors
67.	Almohareb (2016)	Almohareb R, Ahlquist M, Englund GS. The association between root canal filling and treatment outcome. *Swed Dent J*. 2016;40(2):133-142.	[2]Stratified analysis of primary root canal therapy available, if re-treatment cases had been included
68.	Azim (2016)	Azim AA, Griggs JA, Huang GT. The Tennessee study: factors affecting treatment outcome and healing time following nonsurgical root canal treatment. *Int Endod J*. 2016;49(1):6-16.	[6]Overall success rates given or could be calculated from the raw data at the tooth level
69.	deCastroRizzi-Maia (2016)	de Castro Rizzi-Maia C, Maia-Filho EM, Nelson-Filho P, et al. Single vs Two-session Root Canal Treatment: A Preliminary Randomized Clinical Study using Cone Beam Computed Tomography. *J Contemp Dent Pract*. 2016;17(7):515-521.	[6]Overall success rate given or could be calculated from the raw data at the tooth level
70.	Fransson (2016)	Fransson H, Dawson VS, Frisk F, Bjørndal L; EndoReCo, Kvist T. Survival of Root-filled Teeth in the Swedish Adult Population. *J Endod*. 2016;42(2):216-220.	[5]Success based on clinical and/ or radiographic (strict, absence of apical radiolucency; loose, reduction in size of radiolucency) criteria
71.	Hertel (2016)	Hertel M, Sommer K, Kostka E, Imiolczyk SM, Ballout H, Preissner S. Outcomes of Endodontic Therapy Comparing Conventional Sodium Hypochlorite Irrigation with Passive Ultrasonic Irrigation Using Sodium Hypochlorite and Ethylenediaminetetraacetate. A Retrospective Analysis. *Open Dent J*. 2016;10:375-381.	[5]Success based on clinical and/ or radiographic (strict, absence of apical radiolucency; loose, reduction in size of radiolucency) criteria; [4]At least 12-months post-operative review

72.	Pirani (2016)	Pirani, Chiara et al. Prognosis of root canal treatments filled with Thermafil system: a 5-year retrospective study.*Giornale Italiano di Endodonzia* 30 (2016): 46-51.	[7]Other- Overlapping data w/ Pirani 2018 study, which was included in this review
73.	Ricucci (2016)	Ricucci D, Rôças IN, Alves FR, Loghin S, Siqueira JF Jr. Apically Extruded Sealers: Fate and Influence on Treatment Outcome. *J Endod*. 2016;42(2):243-249.	[7]Other- Overlapping data w/ Ricucci 2011 study, which was included in this review
74.	Sigurdsson (2016)	Sigurdsson A, Le KT, Woo SM, et al. Six-month healing success rates after endodontic treatment using the novel GentleWave™ System: The pure prospective multi-center clinical study. *J Clin Exp Dent*. 2016;8(3):e290-e298.	[7]Other- Overlapping data w another Sigurdsson 2016 study which was included in this review
75.	Almeida (2017)	Almeida DO, Chaves SC, Souza RA, Soares FF. Outcome of Single- vs Multiple-visit Endodontic Therapy of Nonvital Teeth: A Meta-analysis. *J Contemp Dent Pract*. 2017;18(4):330-336.	[1]Clinical study on primary root canal therapy: randomized control trials, cohort studies, retrospective observational studies -Systematic Review/ Meta-Analysis
76.	Barborka (2017)	Barborka BJ, Woodmansey KF, Glickman GN, Schneiderman E, He J. Long-term Clinical Outcome of Teeth Obturated with Resilon. *J Endod*. 2017;43(4):556-560.	[7]Other- Excluded due to quality of the study (Resilon)
77.	Bansal (2017)	Bansal A, Parihar AS, Sethi A, Majety KK, Panjabi J, Choudhury BK. Retrospective Assessment of Healing Outcome of Endodontic Treatment for Mandibular Molars with C-shaped Root Canal. *J Contemp Dent Pract*. 2017;18(7):591-595.	[7]Other- only teeth with c-shaped morphology included
78.	Fernandez (2017)	Fernández R, Cardona JA, Cadavid D, Álvarez LG, Restrepo FA. Survival of Endodontically Treated Roots/Teeth Based on Periapical Health and Retention: A 10-year Retrospective Cohort Study. *J Endod*. 2017;43(12):2001-2008.	[6]Overall success rate given or could be calculated from the raw data at the tooth level.
79.	Fonzar (2017)	Fonzar F, Mollo A, Venturi M, et al. Single versus two visits with 1-week intracanal calcium hydroxide medication for endodontic treatment: One-year post-treatment results from a multicentre randomised controlled trial. *Eur J Oral Implantol*. 2017;10(1):29-41.	[2]Stratified analysis of primary root canal therapy available, if re-treatment cases had been included
80.	Ramey (2017)	Ramey K, Yaccino J, Wealleans J. A Retrospective, Radiographic Outcomes Assessment of 1960 Initial	[6]Overall success rate given or could be calculated from the raw data at the tooth level

		Posterior Root Canal Treatments Performed by Endodontists and Dentists. *J Endod*. 2017;43(8):1250-1254.	
81.	Alghofaily (2018)	Alghofaily M, Tordik P, Romberg E, Martinho F, Fouad AF. Healing of Apical Periodontitis after Nonsurgical Root Canal Treatment: The Role of Statin Intake. *J Endod*. 2018;44(9):1355-1360.	[3]Sample size provided- discrepancy in sample size between preoperative and follow-up.
82.	Berrezouga (2018)	Berrezouga L, Bouguezzi A, Belkhir MS. Outcome of Initial Endodontic Treatment Performed, by One Specialist, in 122 Tunisian Patients: A Retrospective Study. *Int J Dent*. 2018;2018:3504245.	[4] At least 12 months post-operative review
83.	Brignardello-Petersen (2018)	Brignardello-Petersen R. High rate of tooth retention and periapical health 10 years after nonsurgical root canal treatment but uncertainty about the factors that may influence these outcomes. *J Am Dent Assoc*. 2018;149(3):e58	[1]Clinical study on primary root canal therapy: randomized control trials, cohort studies, retrospective observational studies -Review/ Systematic Review/ Meta-analysis
84.	Tajonar (2018)	Tajonar RGSY, Sánchez-Mendieta KP, Martínez-Martínez RE, Domínguez-Pérez RA. Periapical Healing of Endodontically Treated Teeth Filled Only in the Apical Third: A Randomized Controlled Trial. *Eur Endod J*. 2017;3(1):24-30.	[7]Other- only teeth filled in the apical 1/3
85.	Laukkanen (2019)	Laukkanen E, Vehkalahti MM, Kotiranta AK. Impact of type of tooth on outcome of non-surgical root canal treatment. *Clin Oral Investig*. 2019;23(11):4011-4018.	[4] At least 12 months post-operative review
86.	Bardini (2020)	Bardini G, Casula L, Ambu E, Musu D, Mercadè M, Cotti E. A 12-month follow-up of primary and secondary root canal treatment in teeth obturated with a hydraulic sealer.*Clin Oral Investig*. 2021;25(5):2757-2764.	[2]Stratified analysis of primary root canal therapy available, if re-treatment cases had been included
87.	Goldberg (2020)	Goldberg F, Cantarini C, Alfie D, Macchi RL, Arias A. Relationship between unintentional canal overfilling and the long-term outcome of primary root canal treatments and nonsurgical retreatments: a retrospective radiographic assessment. *Int Endod J*. 2020;53(1):19-26.	[5]Success based on clinical and/ or radiographic (strict, absence of apical radiolucency; loose, reduction in size of radiolucency) criteria
88.	Garcia-Guerrero (2020)	García-Guerrero C, Delgado-Rodríguez CE, Molano-González N, et al. Predicting the outcome of initial non-surgical endodontic procedures by periapical status and	[5]Success based on clinical and/ or radiographic (strict, absence of apical radiolucency; loose, reduction in size of radiolucency) criteria

		quality of root canal filling: a cohort study. *Odontology*. 2020;108(4):697-703.	
89.	Koli (2020)	Koli B, Chawla A, Logani A, Kumar V, Sharma S. Combination of Nonsurgical Endodontic and Vital Pulp Therapy for Management of Mature Permanent Mandibular Molar Teeth with Symptomatic Irreversible Pulpitis and Apical Periodontitis. *J Endod*. 2021;47(3):374-381.	[1]Clinical study on primary root canal therapy: randomized control trials, cohort studies, retrospective observational studies
90.	Stenhagen (2020)	Stenhagen S, Skeie H, Bårdsen A, Laegreid T. Influence of the coronal restoration on the outcome of endodontically treated teeth. *Acta Odontol Scand*. 2020;78(2):81-86.	[1]Clinical study on primary root canal therapy: randomized control trials, cohort studies, retrospective observational studies