Characteristics of studies conducted on drug related problems in Ethiopia that are excluded from the systematic review and Meta-analysis

| **Studies** | **Year of publication** | **Study area** | **Study Setting** | **Study Design** | **Study subject** | **Sample size** | **Prevalence of DRPs (%)** | **Total DRPs** | **Mean DRP per patient** | **Source of Data** |
|---|---|---|---|---|---|---|---|---|---|---|
| Abdela et al | 2017 | Gondar, Northern Ethiopia | Cardiology unit of a single hospital | CS | CVS patients | 227 | 63.4 | 265 | 1.17±1.1 | Used different classification |
| Hussein et al | 2014 | Adama, Central Ethiopia | Ambulatory unit of a single hospital | CS | HTN patients | 192 | 80.7 | 452 | 2.36 | Incomplete information |
| Tegegne et al | 2014 | Bahir Dar, Northern Ethiopia | Single Hospital | CS | CVS patients | 76 | 96.1 | 105 | 1.38±0.8 | Incomplete information Poor quality |
| Tegegne et al | 2015 | Ambo, Central Ethiopia | Ambulatory clinic of a single hospital | CS | HTN patients | 151 | Not reported | 200 | 1.32±0.5 | Incomplete information |
| Ayele et al | 2018 | Harar, Eastern Ethiopia | Ambulatory clinic of a single hospital | CS | DM with HTN patients | 203 | Not reported | 364 | 1.8 | Incomplete information |

*DRPs-Drug related problems, CS- Cross-sectional, HTN-Hypertension, T2DM-Type two diabetes, CVS-Cardiovascular disease*