



Iteration — Number of studies = 3
Model: Random-effects — observed = 3
Method: DerSimonian-Laird — imputed = 0

Pooling
Model: Random-effects
Method: DerSimonian-Laird


Studies	Hedges' g	[95% Conf.	Interval]
Observed	-1.043	-1.665	-0.421
Observed + Imputed	-1.043	-1.665	-0.421



Iteration — Number of studies = 3
Model: Random-effects — observed = 3
Method: DerSimonian-Laird — imputed = 0

Pooling
Model: Random-effects
Method: DerSimonian-Laird

Studies	Hedges' g	[95% Conf.	Interval]
Observed	-1.043	-1.665	-0.421
Observed + Imputed	-1.043	-1.665	-0.421



Iteration — Number of studies = 3
Model: Random-effects — observed = 3
Method: DerSimonian-Laird — imputed = 0

Pooling
Model: Random-effects
Method: DerSimonian-Laird

Studies	Hedges' g	[95% Conf.	Interval]
Observed	-1.043	-1.665	-0.421
Observed + Imputed	-1.043	-1.665	-0.421

Supplementary Figure 10. Publication bias for trait anxiety: 1 month after psilocybin treatment. A: Linear estimator. B: Run estimator. C: Quadratic estimator.