



A

Iteration — Number of studies = 4
Model: Random-effects — observed = 3
Method: DerSimonian-Laird — imputed = 1

Pooling
Model: Random-effects
Method: DerSimonian-Laird

Studies	Hedges' g	[95% Conf.	Interval]
Observed	-1.030	-1.544	-0.516
Observed + Imputed	-0.857	-1.375	-0.340




B

Iteration — Number of studies = 3
Model: Random-effects — observed = 3
Method: DerSimonian-Laird — imputed = 0

Pooling
Model: Random-effects
Method: DerSimonian-Laird

Studies	Hedges' g	[95% Conf.	Interval]
Observed	-1.030	-1.544	-0.516
Observed + Imputed	-1.030	-1.544	-0.516



C

Iteration — Number of studies = 4
Model: Random-effects — observed = 3
Method: DerSimonian-Laird — imputed = 1

Pooling
Model: Random-effects
Method: DerSimonian-Laird

Studies	Hedges' g	[95% Conf.	Interval]
Observed	-1.030	-1.544	-0.516
Observed + Imputed	-0.857	-1.375	-0.340

Supplementary Figure 12. Publication bias for trait anxiety: 6 months after psilocybin treatment. A: Linear estimator. B: Run estimator. C: Quadratic estimator.