SUPPLEMENTAL MATERIAL

Table S1. Type of dual antiplatelet agents and stents used in the trials.

Trials	Case		Control	
	Dual Antiplatelet Agent	Stent	Dual Antiplatelet Agent	Stent
EBC MAIN	Clopidogrel (67%) Ticagrelor (20%) Prasugrel (6%)	Onyx ZES	Clopidogrel (66%) Ticagrelor (22%) Prasugrel (5%)	Onyx ZES
NBBS IV	Clopidogrel (100%)	Cypher SES	Clopidogrel (99.5%)	Cypher SES
DEFINITION II	Clopidogrel (100%)	Firebird-2 SES, Excel SES, BuMA SES, Partner SES, Xience EES, Endeavor ZES	Clopidogrel (100%)	Firebird-2 SES, Excel SES, BuMA SES, Partner SES, Xience EES, Endeavor ZES
DKCRUSH-V	Clopidogrel (100%)	Xience Prime EES, Resolute ZES	Clopidogrel (100%)	Xience Prime EES, Resolute ZES
COBRA	Clopidogrel (100%)	Axxess BES, Biomatrix BES	Clopidogrel (100%)	Xience EES
DKCRUSH-II	Clopidogrel (100%)	Excel SES	Clopidogrel (100%)	Excel SES
BBK II	Clopidogrel, ticagrelor, or prasugrel	Cypher SES (8%), Xience pro EES (8.7%), Promus EES (28.7%), Resolute ZES (36.0%), Orsiro SES (11.3%), Synergy SES (7.3%)	Clopidogrel, ticagrelor, or prasugrel	Cypher SES (10.0%), Xience pro EES (10.0%), Promus EES (32.7%), Resolute ZES (30.7%), Orsiro SES (14.0%), Synergy SES (3.3%)
POLBOS II	-	BiOSS LIM SES	-	EES (46%), SES (24%), EES (22%), BES (6%), ZES (2%)
EBC TWO	Clopidogrel (100%)	Nobori BES	Clopidogrel (100%)	Nobori BES
SMART-STRATEGY	-	SES (48%), EES (27%), others (25%)	-	SES (47%), EES (31%), others (22%)
***Zhang et al.* (2016)**	Clopidogrel (100%)	SES (79%), ZES (12%), EES (10%)	Clopidogrel (100%)	SES (62%), EES (19%), ZES (15%), PES (4%)
***Zheng et al.* (2016)**	Clopidogrel (100%)	SES	Clopidogrel (100%)	SES
DKCRUSH-III	Clopidogrel (100%)	Firebird-2 SES, Xience V EES	Clopidogrel (100%)	Firebird-2 SES, Xience V EES
BBK I	Clopidogrel (100%)	Cypher SES	Clopidogrel (100%)	Cypher SES
TRYTON	Clopidogrel (100%)	Tryton side branch stent (100%), Xience EES (59.0%), Promus EES (27.5%), Resolute Integrity ZES (6.0%), Endeavor ZES (4.3%), SES (3.2%)	Clopidogrel (100%)	Xience EES (59.2%), Promus EES (30.6%), Resolute Integrity (5.2%), Endeavor EES (2.6%), SES (2.3%), others (0.3%)
POLBOS I	Clopidogrel (100%)	BiOSS Expert PES	Clopidogrel (100%)	EES (41.5%), PES (38.2%), BES (9.8%), SES (8.9%), ZES (0.8%), TES (0.8%)
PERFECT	Clopidogrel (100%)	SES (60%), EES (28%), ZES (9%), PES (1%), others (3%)	Clopidogrel (100%)	SES (57%), EES (29%), ZES (9%), PES (2%), others (3%)
NSTS	Clopidogrel (100%)	Cypher SES	Clopidogrel (100%)	Cypher SES
NBS	Clopidogrel (99.5%)	Cypher SES	Clopidogrel (100%)	Cypher SES
***Ruiz et al.* (2013)**	-	Xience Prime EES	-	Xience Prime EES
***Ye et al.* (2012)**	Clopidogrel (100%)	Cypher SES, Firebird-2 SES	Clopidogrel (100%)	Cypher SES, Firebird-2 SES
BBC ONE	Clopidogrel (100%)	Taxus PES	Clopidogrel (100%)	Taxus PES
***Lin et al.* (2010)**	Clopidogrel or ticloplidine	SES (57%), PES (43%)	Clopidogrel or ticloplidine	SES (65%), PES (35%)
***Ye et al.* (2010)**	Clopidogrel (100%)	SES (57.4%)	Clopidogrel (100%)	SES (64.8%)
CACTUS	Clopidogrel or ticloplidine	Cypher SES	Clopidogrel or ticloplidine	Cypher SES
***Cervinka et al.* (2008)**	Clopidogrel (100%)	Twin-Rail BMS	Clopidogrel (100%)	Liberte BMS
DKCRUSH-I	Clopidogrel (100%)	PES or SES	Clopidogrel (100%)	PES or SES
***Colombo et al.* (2004)**	Clopidogrel or ticloplidine	Cypher SES	Clopidogrel or ticloplidine	Cypher SES
***Pan et al.* (2004)**	Clopidogrel (100%)	SES	Clopidogrel (100%)	SES

Abbreviations: BES = biolimus-eluting stent; BMS = bare metal stent; EES = everolimus-eluting stent; PES = paclitaxel-eluting stent; SES = sirolimus-eluting stent; TES = tacrolimus-eluting stent; ZES = zirolimus-eluting stent

Table S2. Inclusion and exclusion criteria.

Case/Control	Inclusion Criteria	Exclusion Criteria
EBC MAIN	Age ≥18 years Medina type 1,1,1 or 0,1,1 Both MV and SB >50% narrowed Both MV and SB RVD ≥2.75mm Ischemic symptoms, positive non-invasive imaging, positive fractional flow reserve, or LM stem IVUS-derived minimum luminal area <6mm^2	Acute STEMI Cardiogenic shock Chronic total occlusion of either vessel LM trifurcation with all three vessels ≥2.75mm diameter LM stem diameter >5.75mm Life expectancy <12 months Known relevant allergies
NBBS IV	Stable angina pectoris, unstable angina, or silent ischemia Medina 1,1,1, 1,0,1, or 0,1,1 MV diameter ≥3.0mm SB diameter ≥2.75mm	STEMI within 24 hours SB lesion length >15mm Life expectancy <1 year Serum creatinine >200μmol/L Allergy to aspirin, clopidogrel, ticlopidine, sirolimus, or everolimus
DEFINITION II	Age ≥18 years Medina 1,1,1 or 0,1,1 SB RVD ≥2.5mm Native coronary lesion Complex bifurcation lesions based on DEFINITION study	STEMI within 24 hours Pregnancy or breastfeeding Life expectancy <50% at 12 months Scheduled major surgery in 12 months Allergy to aspirin, clopidogrel, or ticagrelor History of major hemorrhage Chronic total occlusion in either LAD, LCx, or RAD Severe calcification needing rotational atherectomy
DKCRUSH-V	Ages 18 to 80 years Stable angina, unstable angina, or myocardial infarction Medina 1,1,1 or 0,1,1 De novo unprotected LM lesion Unprotected LM lesion with chronic total occlusion in LAD, LCx, or RCA after recanalization Diameter stenosis in LAD/LM and LCx ≥50% by visual estimation	Acute myocardial infarction within 24 hours Severe calcification needing rotational atherectomy Restenotic lesion Allergies to study or protocol-required medications Intolerance to dual anti-platelet therapy Life expectancy <12 months Pregnancy or breastfeeding Distal LM coronary restenosis
COBRA	Age >18 years Stable angina, unstable angina, or positive functional study De novo true native coronary bifurcation lesion Medina 1,1,1, 1,0,1, or 0,1,1 Proximal MV RVD 2.75-3.75mm SB RVD >2.25mm	LVEF <30% Serum creatinine >2.0mg/dL Previous or planned brachytherapy of target vessel Unprotected lesion of the LM trunk 50% Intraluminal thrombus in the target vessel Moderate to severe calcification or tortuosity Allergies to protocol stent or medications Pregnancy or breastfeeding Life expectancy <12 months
DKCRUSH-II	Age ≥18 years Silent ischemia, angina, or acute myocardial infarction Chronic total occlusion in MV or SB immediately after successful recanalization Unprotected distal LM bifurcation lesions involving both ostia of LAD and LCx without chronic total occlusion in the RCA	Liver dysfunction Life expectancy <12 months Heavy calcification requiring rotational atherectomy Pregnancy Contraindication to one of the study drugs

	Only one coronary bifurcation lesion Medina 1,1,1 and 0,1,1 Diameter stenosis >50% in both vessels with RVD between 2.5 and 4.0mm	
BBK II	De novo bifurcation lesion MV RVD 2.5 to 4.0mm SB RVD ≥2.25mm and ≤1.0mm smaller than that of RV	Acute STEMI Hemodynamic instability History of bleeding diathesis or coagulopathy Intraluminal thrombus Heavy calcification Severe tortuosity Contraindication to study drugs
POLBOS II	Age ≥18 years Stable coronary artery disease or non-STEMI De novo coronary bifurcation lesion MV RVD ≥2.5mm SB RVD ≥2.0mm	STEMI LVEF ≤30% Medina 0,0,1 Serum creatinine ≥2.0mg/dL Inability to take dual antiplatelet therapy for 12 months
EBC TWO	Age ≥18 years Both MV and SV >50% narrowed Both MV and SV RVD ≥2.5mm SB ostial disease ≥5mm SB lesion causing angina and a potential target for intervention	Acute STEMI Cardiogenic shock Unprotected LM stem narrowing ≥50% Chronic total occlusion of either vessel Additional type C lesion requiring PCI LVEF ≤20% Platelet count ≤50,000/μL Life expectancy < 12 months Known relevant allergies
SMART-STRATEGY	Stable coronary artery disease or non-STEMI De novo coronary bifurcation lesion, including unprotected LM MV RVD ≥2.5mm SB RVD ≥2.3mm	Hemodynamic instability LVEF <25% Undergoing primary PCI
***Zhang et al.* (2016)**	Age ≥18 years De novo true coronary bifurcation lesion Medina 1,1,1, 0,1,1, and 1,0,1 Located either in LM stem, LAD, and LCx MV RVD ≥2.5mm SB RVD ≥2.0mm	Acute STEMI within 24 hours Life expectancy <12 months Allergies to any drugs used in the study
***Zheng et al.* (2016)**	De novo coronary true bifurcation lesion SB RVD ≥2.25mm	Acute STEMI within 24 hours Liver or renal dysfunction LVEF ≤30% Life expectancy <1 year Platelet count ≤10,000/μL Suspected intolerance to any of the drugs used
DKCRUSH-III	Age ≥18 years Ischemic symptoms or myocardial ischemia LMV RVD ≤5mm Difference in RVD between LAC and LCX ≤1mm Medina 1,1,1 or 0,1,1 De novo LM distal brain lesion Chronic total occlusion after successful recanalization	Acute myocardial infarction within 2 weeks Liver dysfunction Lung cancer LVEF <30% eGFR <40mL/min/1.73m^2 Trifurcation lesions or heavy calcification Pregnancy

		Platelet count ≤10,000/μL Suspected intolerance to any of the study drugs
BBK I	Stable angina or positive stress test De novo bifurcation lesion Native coronary artery >50% diameter stenosis MV RVD 2.5-4mm SB RVD ≥2.25mm	LM stenosis Intraluminal thrombus Heavy calcification Severe tortuosity Contraindication to drug or stent used in the study History of bleeding diathesis or coagulopathy
TRYTON	Symptoms or objective evidence of ischemia True bifurcation lesion ≥50% narrowing Medina 1,1,1, 1,0,1, and 0,1,1 MV RVD 2.5-4mm SB RVD 2.5-3.5mm Lesion length in MV ≤28mm Lesion length in SB ≤5mm	STEMI within 72 hours Non-STEMI within 7 days LVEF <30% Serum creatinine >2.5mg/dL LM disease Trifurcation lesion Total occlusion of target vessel Severe calcification Excessive tortuosity Angiographic evidence of thrombus
POLBOS I	Age ≥18 years Stable coronary artery disease or non-STEMI De novo coronary bifurcation lesion MV RVD ≥2.5mm SB RVD ≥2.0mm	STEMI Medina 0,0,1 Serum creatinine ≥2.0mg/dL Inability to receive dual antiplatelet therapy for 12 months LVEF ≤30%
PERFECT	Age 18-75 years Angina with bifurcation coronary disease requiring protection MV RVD ≥2.5mm MV lesion length ≤50mm SB RVD ≥2.0mm	LM disease In-stent restenosis Graft lesions Chronic total occlusion STEMI within 2 weeks Decreased SB flow Renal failure LVEF ≤35% Life expectancy <12 months
NSTS	Age ≥18 years Stable angina, unstable angina, or silent ischemia De novo coronary bifurcation lesion MV RVD ≥3.0mm SB RVD ≥2.5mm	STEMI within 24 hours Life expectancy <12 months Serum creatinine ≥200μmol/L Allergy to any of the drugs used
NBS	Age ≥18 years Stable angina, unstable angina, or silent ischemia MV RVD ≥2.5mm SB RVD ≥2.0mm	STEMI within 24 hours Life expectancy <12 months Serum creatinine ≥200μmol/L Allergy of any of the drugs used LM bifurcation in a left dominant system
***Ruiz et al.* (2013)**	True bifurcation lesions MV RVD 2.5-4mm SB RVD >2mm	LM disease Thrombotic lesions Acute coronary syndrome within 48 hours LVEF <30% Serum creatinine >3mg/dL

***Ye et al.* (2012)**	Age 18-75 years Medina 1,1,1, 1,0,1, or 0,1,1 SB RVD ≥2.25mm	LVEF ≤30% Life expectancy <12 months Pregnancy Severe diffuse or calcified MV or SB lesions Platelet count ≤10,000/μL Serum creatinine >3mg/dL Cerebrovascular events within 6 months STEMI or other condition with refractory hypotension Intolerance to injection of adenosine
BBC ONE	Age ≥18 years MV RVD ≥2.5mm SB RVD ≥2.25mm	STEMI Cardiogenic shock Chronic total occlusion of either vessel LM stem narrowing ≥50% Additional type C or bifurcation lesion that required PCI LVEF ≤20% Platelet count ≤50,000/μL Life expectancy <12 months Known relevant allergies
***Lin et al.* (2010)**	Age ≥18 years Stable angina, unstable angina, or silent ischemia De novo coronary true bifurcation Bifurcation angle ≤60 degrees MV RVD ≥2.5mm SB RVD ≥2.22mm	Myocardial infarction within 24 hours Life expectancy <12 months Serum creatinine >3.0mg/dL Allergy to any of the drugs used LM bifurcation in a left dominant system
***Ye et al.* (2010)**	Age 18-75 years Single vessel disease with de novo bifurcation lesion Medina 1,1,1, 1,0,1, or 0,1,1 MV RVD ≥2.0mm SB RVD ≥2.0mm	STEMI LVEF ≤30% Life expectancy <1 year Pregnancy Severely diffuse or calcified lesions Platelet count ≤100,000/μL Cerebrovascular events within 6 months Allergy to aspirin, clopidogrel, or sirolimus
CACTUS	Age ≥18 years Stable angina, unstable angina, or silent ischemia De novo true coronary bifurcation lesion Stenosis in both MB and ostium of the SB >50% Both branches with TIMI flow ≥1 Treatable lesion length ≤28mm MV RVD 2.5-3.5mm SB RVD 2.25-3.5mm	Myocardial infarction within 24 hours LM trunk unprotected by a graft Visible thrombus within target lesion Chronic total occlusion LVEF <35% Serum creatinine ≥2.65μmol/L Contraindication to one of the study drugs
***Cervinka et al.* (2008)**	Symptoms or signs of angina All types but Medina 0,1,0 Lesion length <15mm MV RVD 2.7-4.0mm SB RVD >2.2mm	Not reported
DKCRUSH-I	MV RVD ≥2.5mm SB RVD ≥2.0mm	Life expectancy <12 months Liver dysfunction

	Two or more bifurcation lesions if there was one bifurcation lesion per vessel Another single lesion in a different target vessel that could be covered by a single DES were also included	Plasma creatinine ≥200μmol/L Cerebrovascular within 6 months History of coronary artery bypass grafting Allergy to aspirin, clopidogrel, and stent
***Colombo et al.* (2004)**	Age ≥18 years Stable angina, unstable angina, or silent ischemia De novo true coronary bifurcation lesion Stenosis in both MB and ostium of the SB >50% Both branches with TIMI flow ≥1 Treatable lesion length ≤24mm Both MV and SB RVD 2.5-3.5mm	Myocardial infarction within 24 hours Stenosis of LM unprotected by graft Thrombus within the target lesion LVEF ≤35% Serum creatinine ≥3.0mg/dL Suspected intolerance to one of the study drugs
***Pan et al.* (2004)**	Lesion located in major bifurcation regardless of morphology and angulation MV RVD ≥2.5mm SB RVD ≥2.25mm Significant stenosis in both main vessel and SB origin	Diffuse SB lesions

Abbreviations: IVUS = intravascular ultrasound; LAD = left anterior descending coronary artery; LCx = left circumflex coronary artery; LM = left main coronary artery; LVEF = left ventricular ejection fraction; MV = main vessel; RCA = right coronary artery; RVD = reference vessel diameter; SB = side branch (vessel); STEMI = ST elevation myocardial infarction

Table S3. Anatomical characteristics of bifurcation lesions included in each trial.

Trial	Left Main Included	True Bifurcation	Medina Classification	Main Vessel		Side Branch		Lesion Length, mm
				Diameter, mm	Stenosis, %	Diameter, mm	Stenosis, %	
EBC MAIN	Yes	Yes	1,1,1 0,1,1	≥2.75	>50%	≥2.75	>50%	-
NBBS IV	Yes	Yes	1,1,1 1,0,1 0,1,1	≥3.0	≥50%	≥2.75	≥50%	SB ≤15
DEFINITION II	Yes	Yes	1,1,1 0,1,1	<2.5*	-	≥2.5	Distal LM ≥70%* Non-LM ≥90%*	MV ≥25* SB ≥10*
DKCRUSH-V	Yes	Yes	1,1,1 0,1,1	-	≥50%	-	≥50%	-
COBRA	No	Yes	1,1,1 1,0,1 0,1,1	2.75-3.75	-	2.25	-	-
DKCRUSH-II	Yes	Yes	1,1,1 0,1,1	2.5-4.0	>50%	2.5-4.0	>50%	≤2 stents
BBK II	Yes	No	1,0,0 0,1,0 1,1,0 1,1,1 0,0,1 1,0,1 0,1,1	2.5-4.0	-	2.25	-	-
POLBOS II	Yes	No	1,0,0 0,1,0 1,1,0 1,1,1 1,0,1 0,1,1	≥2.5	-	≥2.0	-	-
EBC TWO	No	Yes	1,1,1 1,0,1 0,1,1	≥2.5	>50%	≥2.5	>50%	-
SMART-STRATEGY	Yes	No	1,0,0 0,1,0 1,1,0 1,1,1 0,0,1 1,0,1 0,1,1	≥2.5	-	≥2.3	-	-
***Zhang et al.* (2016)**	Yes	Yes	1,1,1 1,0,1 0,1,1	≥2.5	-	≥2.0	-	-

***Zheng et al.* (2016)**	Yes	Yes	1,1,1 1,0,1 0,1,1	-	-	≥2.25	-	-
DKCRUSH-III	Yes	Yes	1,1,1 0,1,1	≤5	-	-	-	-
BBK I	No	No	1,0,0 0,1,0 1,1,0 1,1,1 0,0,1 1,0,1 0,1,1	2.5-4	>50%	≥2.25	>50%	-
TRYTON	No	Yes	1,1,1 1,0,1 0,1,1	2.5-4	≥50%	2.5-3.5	≥50%	MV ≤28 SB ≤5
POLBOS I	Yes	No	1,0,0 0,1,0 1,1,0 1,1,1 1,0,1 0,1,1	≥2.5	-	≥2.0	-	-
PERFECT	No	No	1,0,0 0,1,0 1,1,0 1,1,1 0,0,1 1,0,1 0,1,1	≥2.5	-	≥2.0	≥50%	MV ≤50
NSTS	Yes	No	-	≥3.0	-	≥2.5	-	-
NBS	Yes	No	-	≥2.5	-	≥2.0	-	-
***Ruiz et al.* (2013)**	No	No	1,0,0 0,1,0 1,1,0 1,1,1 1,0,1 0,1,1	2.5-4	-	>2	-	-
***Ye et al.* (2012)**	No	Yes	1,1,1 1,0,1 0,1,1	-	-	≥2.25	-	≤2 stents
BBC ONE	No	No	1,0,0 0,1,0 1,1,0 1,1,1 0,0,1	≥2.5	-	≥2.25	-	-

			1,0,1 0,1,1					
***Lin et al.* (2010)**	No	Yes	1,1,1 1,0,1 0,1,1	≥2.5	-	≥2.2	-	-
***Ye et al.* (2010)**	No	Yes	1,1,1 1,0,1 0,1,1	≥2.0	-	≥2.0	-	-
CACTUS	No	Yes	1,1,1 1,0,1 0,1,1	2.5-3.5	>50%	2.25-3.5	>50%	≤28
***Cervinka et al.* (2008)**	No	No	1,1,1 1,0,0 0,1,0	2.7-4.0	-	>2.2	-	<15
DKCRUSH-I	Yes	Yes	-	≥2.5	-	≥2.0	-	≤2 stents
***Colombo et al.* (2004)**	No	Yes	-	2.5-3.5	>50%	2.5-3.5	>50%	≤24
***Pan et al.* (2004)**	Yes	Yes	1,1,1 0,1,1	≥2.5	Significant	≥2.25	Significant	-

*One of the criteria of complex bifurcation lesions included in the trial

Abbreviations: LM = left main coronary artery; MV = main vessel; SB = side branch

Table S4. Procedural characteristics.

Trial	Radial Approach, %	Intravascular Ultrasound, %	Main Vessel		Side Branch		POT, %	FKBI, %
Case/Control			Stent Diameter, mm ± SD	Stent Length, mm ± SD	Stent Diameter, mm ± SD	Stent Length, mm ± SD		
EBC MAIN	70/71	31/36	3.6±0.6/3.8±0.5	21.8±7.0/22.1±7.0	3.6±0.6/3.5±0.6	19.3±6.7/17.6±6.9	87/85	93/22
NBBS IV	-	-	-	24.3±9.6/25.0±9.5	-	9(6:13)/13([8:15]*	-	91.2/36.1
DEFINITION II	80.6/78.7	24.4/31.1	3.1±0.3/3.0±0.3	46.3±19.3/46.5±19.7	2.6±0.3/2.8±0.4	25.6±11.3/26.5±12.3	64.6/100	99.3/95.9
DKCRUSH-V	-	42.9/40.5	-	-	-	-	99.2/98.8	99.6/78.9
COBRA	-	-	3.4±0.2/3.0±0.3	49.2±14.1/49.1±17.8**	2.7±0.2/2.7±0.3	-	-	100/100
DKCRUSH-II	-	46.2/47.8	-	28.6±12.4/28.8±13.5	-	16.2±9.1/16.7±8.6	-	100/79.2
BBK II	35.3/37.3	-	3.3±0.4/3.3±0.4	26.7±9.4/26.4±11.7	2.9±0.4/2.8±0.4	21.6±7.7/18.5±8.9	-	100/100
POLBOS II	63.7/81	-	3.7±0.5/3.35±0.5	17.8±2.7/19.9±6.3	-	-	37.3/68	32.7/49.0
EBC TWO	57/63	-	3.0±0.3/3.1±0.3	22.9±5.1/23.4±4.8	2.7±0.3/2.6±0.3	20.7±5.5/19.9±6.8	-	96/94
SMART-STRATEGY	-	98.5/96.9	3.3±0.4/3.3±0.4	25.1±5.3/24.9±5.6	2.9±0.4/2.8±0.2	17.7±5.6/18.4±7.8	-	68.5/25.8
***Zhang et al.* (2016)**	-	-	-	-	-	-	-	92.3/82.7
***Zheng et al.* (2016)**	-	-	-	22.8±7.5/24.6±6.7	-	10.4±5.6/10.2±5.8	-	71.3±86.0
DKCRUSH-III	58.1/58.9	69.0/73.7	3.4±0.3/3.3±0.4	33.5±14.0/35.7±16.0	3.0±0.4/3.0±0.4	25.9±13.8/26.7±11.9	-	99.5/99.5
BBK I	-	-	-	-	-	-	-	100/100
TRYTON	34.6/35.5	-	3.1±0.4/3.1±0.4	14.8±4.0/14.4±3.5	2.6±0.4/2.4±0.4	13.4±3.3/13.8±3.5	-	89.3/88.8
POLBOS I	80.8/82.1	-	3.7±0.3/3.2±0.5	17.4±2.5/20.7±6.8	-	-	37.5/69.1	30.8/49.6
PERFECT	11.7/12.1	91.5/79.6	3.3±0.3/3.3±0.3	37.3±14.7/36.9±15.3	2.7±0.2/2.7±0.2	21.4±6.7/21.5±6.9	-	95.8/79.1
NSTS	-	-	-	23.5±9.3/23.6±9.1	-	10.6±5.6/10.6±5.8	-	85/92
NBS	-	-	-	23.2±8.6/23.4±8.6	-	10.3±5.0/2.8±6.1	-	74/32
***Ruiz et al.* (2013)**	-	-	-	22±11/25±11	-	-	-	64/42
***Ye et al.* (2012)**	-	-	3.1±0.4/3.2±0.4	33.5±12.4/30.7±16.3	2.7±0.3/2.8±0.3	20.7±7.4/18.0±8.2	-	100/86.7
BBC ONE	29/34	-	3.2±0.3/3.0±0.3	22±6/21±6	2.6±0.3/NR	16±5/NR	-	90/31
***Lin et al.* (2010)**	-	-	3.5±0.1/3.6±0.2	23.6±2.1/23.8±2.6	2.9±0.1/2.9±0.2	12.7±2.8/12.9±3.1	-	90.7/94.4
***Ye et al.* (2010)**	-	-	-	-	-	-	-	-
CACTUS	-	3.4/4.1	-	23.8±5.9/22.2±5.7	-	17.9±5.0/18.1±6.2	-	92.1/90.2
***Cervinka et al.* (2008)**	-	-	-	-	-	-	-	97/97
DKCRUSH-I	-	-	3.4±0.4/3.4±0.5	22.2±12.5/25.0±12.7	2.7±0.4/2.8±0.4	17.4±5.6/17.9±7.0	-	100/76.1
***Colombo et al.* (2004)**	-	100100	3.1±0.3/3.2±0.3	-	2.7±0.3/2.6±0.4	-	-	90.5/81.8
***Pan et al.* (2004)**	-	-	2.9±0.3/2.9±0.3	26±9/25±12	-	-	-	77/60

*Median with interquartile range

**Includes stent length in the side branch

Abbreviation: FKBI = final kissing balloon inflation; NR = not reported; POT = proximal optimization technique; SD = standard deviation

Table S5. Definition of outcomes.

Trial	Cardiac Death	Myocardial Infarction	Target Lesion Revascularization	Target Vessel Revascularization	Stent Thrombosis	Major Adverse Cardiac Events
EBC MAIN	Not specified	Third Universal Definition of Myocardial Infarction except for PCI- and CABG-related MI, for which Society for Cardiovascular Angiography and Interventions (SCAI) consensus definition was used	Repeat revascularization of MV or SB within treated vessel or 5mm adjacent area	Not specified	ARC definition	All-cause death, MI, TLR
NBBS IV	Death from coronary artery disease, heart failure, or cardiac procedure within 28 days	Rise or fall or cardiac biomarkers with at least one value above the 99th percentile of ULN and evidence of ischemia in the myocardium documented by either ischemic symptoms, ECG changes, evidence of new loss of viable myocardium, or new RMWA Sudden and unexpected cardiac death Pathological findings suggestive of acute MI	Repeat revascularization by PCI or CABG of the target lesion in stented segments or their 5mm margins	Not specified	ARC definition	Cardiac death, non-procedural MI, clinically indicated TLR, definite stent thrombosis within 6 months
DEFINITION II	Death from sudden cardiac death, acute MI, arrhythmia, heart failure, stroke, other cardiovascular causes, or bleeding	Post-procedure (within 48 hours after PCI): CK-MB ≥10 times ULN or ≥5 times with new pathologic Q waves in at least 2 contiguous leads or new LBBB, rise in CK-MB to aforementioned increment from the most recent pre-procedure level Spontaneous MI (48 hours after PCI): rise of CK-MB or troponin >1 times ULN with evidence of prolonged ischemia, ischemic ST segment changes, new pathological Q waves, angiographic evidence of flow-limiting complication, imaging evidence of new loss of viable myocardium, or new RWMA	Repeat revascularization for target lesions in the presence of symptoms or objective signs of ischemia	Repeat revascularization for target vessels in the presence of symptoms or objective signs of ischemia	ARC definition	Cardiac death, target-vessel MI, clinically driven TLR
DKCRUSH-V	Any death without a clear non-cardiac cause	Peri-procedural MI: CK-MB >10 times ULN or >5 times ULN with either new pathological Q waves in 2 contiguous leads; angiographically documented graft or coronary artery occlusion or new severe stenosis with thrombosis; or imaging evidence of new loss of viable myocardium or new RWMA Spontaneous MI 72 hours after PCI: clinical syndrome consistent with MI with CK-MB or troponin >1 times ULN and new ST segment elevation or depression or equivalent	Angina or ischemic referable to the target lesion requiring clinically driven PCI or CABG	Not specified	ARC definition	Cardiac death, target-vessel MI (all MI unless clearly attributable to non-target vessel), TLR
COBRA	All deaths considered cardiac unless unequivocal non-cardiac cause demonstrated	Evidence of myocardial necrosis consistent with myocardial ischemia, typically including detection of cardiac biomarkers with at least one value above the 99th percentile of ULN together with ischemic symptoms, pathognomonic ECG, or imaging evidence of ischemia By convention, increase of cardiac biomarkers above 3 times ULN was used to defined PCI-related MI	Revascularization of significant angiographic restenosis ≥70% anywhere in the target lesion in combination with angina or FFR <0.80 in MV or SB subtending a large myocardial territory	Not specified	Not specified	Cardiac death, MI (CK-MB based), TVR
DKCRUSH-II	All deaths considered cardiac	Rise of biochemical markers above 99th percentile with at least one of the following ischemic symptoms:	Repeat revascularization by	Repeat revascularization by	ARC definition	Cardiac death, MI, TVR

	unless other cause documented	ECG changes indicative of ischemia, development of pathological Q waves, and no relation to PCI procedure	PCI or surgery of the target lesion	PCI or surgery of the target vessel		
BBK II	Not specified	During hospitalization: presence of new Q waves in 2 or more contiguous ECG leads or an elevation of CK or CK-MB to at least 3 times of ULN in 2 samples After discharge: ESC/ACC guidelines (2000) and based on new rise in troponin T associated with typical symptoms, ECG changes, and/or angiographic findings	CABG or repeat PCI involving the stented segmented and performed for symptoms or signs of ischemia in the presence of angiographic restenosis	Not specified	ARC definition	All-cause death, MI, TLR
POLBOS II	All deaths deemed cardiac unless proven otherwise	Third Universal Definition of Myocardial Infarction	Re-intervention of target lesion because of symptomatic stenosis ≥50% of diameter	Revascularization of any segment of the index coronary artery	Not specified	Cardiac death, MI TLR
EBC TWO	Not specified	Typical rise and fall of biochemical markers of myocardial necrosis with ischemic symptoms or ECG changes as per ESC/ACC 2000 Peri-procedural MI: elevation of troponin >5 times the 99th percentile in patients with normal baseline values or rise >20% if the baseline values are elevated and are stable or falling In addition, either symptoms suggestive of myocardial ischemia, new ischemic ECG changes, angiographic findings consistent with procedural complication, or imaging demonstrating of new loss of viable myocardium or new RWMA	Not specified	PCI or CABG of either the MV or SB and/or TIMI flow <3 in either the MV or SB after vasodilators	ARC definition	All-cause death, MI, TVR during index admission
SMART-STRATEGY	All deaths considered cardiac unless definite non-cardiac cause established	Elevated troponin or CK-MB greater than ULN with ischemic symptoms or ECG findings indicative of ischemia unrelated to the index procedure	Repeat PCI of lesion within 5 mm of stent or CABG of the target vessel	Repeat revascularization of target vessel by PCI or CABG	ARC definition	Target-vessel failure, target bifurcation revascularization, TVL, TVR
***Zhang et al.* (2016)**	Not specified	NQWMI: CK-MB or troponin-T/I increase ≥3 times ULN with clinical signs of MI but without Q waves and not related to intervention QWMI: New pathological Q waves in 2 or more contiguous leads with clinical signs of MI	Repeat target lesion therapy either by PCI or surgery	Repeat target vessel therapy either by PCI or surgery	ARC definition	Cardiac death, MI, TLR, TVR, stent thrombosis
***Zheng et al.* (2016)**	ARC definition	ARC definition	ARC definition	ARC definition	ARC definition	Cardiac death, MI, TVR, stent thrombosis
DKCRUSH-III	ARC definition	ARC definition	Any repeat PCI of the stented segment, including 5mm proximal and distal margins	Not specified	ARC definition	Cardiac death, MI, TVR
BBK I	Not specified	During hospitalization: presence of new Q waves in 2 or more contiguous ECG leads or an elevation of CK or CK-MB to at least 3 times of ULN in 2 samples	CABG or repeat PCI involving the stented segmented and	Not specified	ARC definition	All-cause death, MI, TLR

		After discharge: ESC/ACC guidelines (2000) and based on new rise in troponin T associated with typical symptoms, ECG changes, and/or angiographic findings	performed for symptoms or signs of ischemia in the presence of angiographic restenosis			
TRYTON	Not specified	CK-MB >3 times ULN	Clinically or ischemia-driven revascularization of target lesion	Clinically or ischemia-driven revascularization of target vessel	ARC definition	All-cause death, MI, CABG, TLR
POLBOS I	All deaths deemed cardiac unless proven otherwise	Third Universal Definition of Myocardial Infarction	Re-intervention of target lesion because of symptomatic stenosis ≥50% of diameter	Revascularization of any segment of the index coronary artery	Not specified	Cardiac death, MI TLR
PERFECT	All deaths considered cardiac unless unequivocal non-cardiac cause established	Increase in CK-MB >3 times ULN with ischemic symptoms or new ischemic ECG changes	Repeat revascularization with PCI or CABG within the stent or adjacent 5mm margins	Repeat revascularization in treated vessel with ≥50% stenosis and ischemic signs or symptoms	Not specified	All-cause death, MI, TVR
NSTS	All deaths considered cardiac unless other cause documented	Rise of biochemical markers above the 99th percentile of ULN with ischemic symptoms, ischemic ECG changes, pathological Q waves, or no relation to PCI procedure	Repeat revascularization by PCI or surgery of the target lesion	Repeat revascularization by PCI of surgery of the target vessel	ARC definition	All-cause death, MI, TVR, stent thrombosis
NBS	All deaths considered cardiac unless other cause documented	NQWMI: CK-MB or troponin-T/I increase ≥3 times ULN with clinical signs of MI and not related to intervention QWMI: New pathological Q waves in 2 or more contiguous leads with clinical signs of MI	Repeat revascularization by PCI or surgery of the target lesion	Repeat revascularization by PCI of surgery of the target vessel	Angiographically verified	Cardiac death, MI, TVR, stent thrombosis
***Ruiz et al.* (2013)**	Not specified	Hospital admission with diagnosis of acute coronary syndrome with or without ST segment elevation	Not specified	Repeat revascularization of target vessel	ARC definition	Cardiac death, MI, TVR
***Ye et al.* (2012)**	All deaths considered cardiac unless otherwise documented	Rise of CK-MB 3 times ULN	Repeat revascularization for a stenosis >50% in target lesion	Clinically driven revascularization of target vessel	ARC definition	Cardiac death, MI, TLR, TVR
BBC ONE	Not specified	First 24 hours after PCI: CK 3 times ULN, CK rise to 50% of the previous value for patient who already had a diagnosis of MI After 24 hours from PCI: ESC/ACC guidelines (2000)	Not specified	PCI or CABG of either the MV or SB and/or TIMI flow <3 in either the MV or SB after vasodilators	ARC definition	All-cause death, MI, target-vessel failure
***Lin et al.* (2010)**	Not specified	Not specified	Repeat revascularization with stenosis diameter ≥50% within the stent or adjacent 5mm margins	Repeat revascularization within the treated vessel	ARC definition	Cardiac death, MI, TVR, stent thrombosis
***Ye et al.* (2010)**	All deaths considered cardiac	Rise of CK-MB 3 times ULN	Repeat revascularization for a	Repeat revascularization	ARC definition	Cardiac death, MI, TLR, TVR

	unless otherwise documented		stenosis >50% in target lesion	within the treated vessel		
CACTUS	Not specified	QWMI: New Q waves in 2 or more contiguous leads with post-procedure CK or CK-MB above ULN NQWMI: Postprocedural CK levels 2 times ULN with elevated CK-MB in the absence of pathological Q waves	Not specified	Not specified	ARC definition	Cardiac death, MI, TVR
***Cervinka et al.* (2008)**	Not specified	QWMI: New Q waves in 2 or more ECG leads with post-procedural elevation of CK 3 times ULN and CK-MB >10% of CK level NQWMI: Post-procedural elevation of CK 3 times ULN and CK-MB above normal	Revascularization within 5mm of the stent edges	Revascularization in the treated vessel	ARC definition	All-cause death, MI, CABG, PCI
DKCRUSH-I	Not specified	CK-MB enzyme elevation 3 times ULN either with or without Q waves in at least two contiguous leads on ECG	Repeat revascularization with a diameter stenosis ≥50% within the stent or in the 5mm distal or proximal margins	Repeat revascularization within the treated vessel	ARC definition	Cardiac death, MI, TLR
***Colombo et al.* (2004)**	Not specified	Not specified	Not specified	Not specified	In-stent occlusion or thrombus, or death or MI within 30 days	Cardiac death, MI, TVR
***Pan et al.* (2004)**	Not specified	Increase in CK level 3 times ULN	Not specified	Not specified	Not specified	Cardiac death, MI, TVR

Abbreviations: CABG = coronary artery bypass graft; CK = creatine kinase; ECG = electrocardiogram; FFR = fractional flow reserve; MI = myocardial infarction; MV = main vessel; NQWMI = non-Q-wave myocardial infarction; PCI = percutaneous coronary intervention; QWMI = Q-wave myocardial infarction; RWMA = regional wall motion abnormality; SB = side branch; TLR = target lesion revascularization; TVR = target vessel revascularization; ULN = upper limit of normal

Table S6. Quantitative coronary angiography at baseline.

Trial	Proximal Main Vessel				Distal Main Vessel				Side Branch			
Case/Control	RVD (mm)	MLD (mm)	DS (%)	Length (mm)	RVD (mm)	MLD (mm)	DS (%)	Length (mm)	RVD (mm)	MLD (mm)	DS (%)	Length (mm)
EBC MAIN	3.8 ± 0.7 3.8 ± 0.8	1.6 ± 0.8 1.8 ± 0.9	56 ± 21 53 ± 21	6.3 ± 2.8 6.4 ± 3.2	2.9 ± 0.6 2.8 ± 0.4	1.3 ± 0.5 1.1 ± 0.4	56 ± 17 59 ± 14	8.0 ± 5.1 8.4 ± 6.1	2.7 ± 0.5 2.7 ± 0.6	1.19 ± 0.5 1.29 ± 0.6	55 ± 16 52 ± 19	7.9 ± 5.7 5.8 ± 4.0
NBBS IV	3.2 ± 0.6 3.1 ± 0.5	1.41 ± 0.6 1.29 ± 0.6	57 ± 17 59 ± 16	19.5 ± 8.9 20.8 ± 9.9	2.61 ± 0.5 2.57 ± 0.5	1.43 ± 0.6 1.43 ± 0.6	43 ± 22 40 ± 20	-	2.40 ± 0.5 2.33 ± 0.5	1.21 ± 0.5 1.43 ± 0.7	49 ± 17 43 ± 18	7.7 ± 4.9 6.4 ± 4.1
DEFINITION II	3.5 ± 0.5 3.4 ± 0.5	2.0 ± 1.0 2.0 ± 1.0	43 ± 26 43 ± 25	41 ± 13* 42 ± 15*	2.7 ± 0.4 2.7 ± 0.4	1.12 ± 0.5 1.15 ± 0.5	58 ± 19 57 ± 18	-	2.38 ± 0.4 2.37 ± 0.5	0.91 ± 0.4 0.91 ± 0.4	62 ± 16 62 ± 15	20.7 ± 10 19.9 ± 9
DKCRUSH-V	3.1 ± 0.5 3.1 ± 0.5	1.2 ± 0.6 1.2 ± 0.5	61 ± 7 62 ± 8	22.4 ± 13 23.5 ± 13	-	-	-	-	2.7 ± 0.4 2.7 ± 0.4	1.03 ± 0.5 1.02 ± 0.4	66 ± 7 65 ± 8	16.2 ± 14 16.6 ± 12
COBRA	3.4 ± 0.5 3.4 ± 0.5	1.86 ± 0.7 1.46 ± 0.7	57 ± 13 62 ± 14	7.35 ± 3.7 10.1 ± 3.7	2.56 ± 0.5 2.49 ± 0.6	1.32 ± 0.5 1.51 ± 0.6	48 ± 16 39 ± 17	10.6 ± 8.0 7.6 ± 5.8	2.33 ± 0.3 2.21 ± 0.4	0.94 ± 0.4 1.19 ± 0.6	59 ± 18 46 ± 23	9.2 ± 5.9 7.1 ± 5.3
DKCRUSH-II	2.9 ± 0.3 2.8 ± 0.4	0.94 ± 0.4 0.86 ± 0.4	67 ± 14 69 ± 17	28.4 ± 13 28.7 ± 16	-	-	-	-	2.38 ± 0.3 2.29 ± 0.4	0.89 ± 0.3 0.84 ± 0.3	63 ± 15 63 ± 14	15.4 ± 11 14.9 ± 13
BBK II	3.1 ± 0.5 3.2 ± 0.5	1.28 ± 0.8 1.29 ± 0.8	59 ± 23 59 ± 24	23.9 ± 7.6 22.7 ± 7.3	2.9 ± 0.5 3.0 ± 0.5	1.24 ± 0.8 1.19 ± 0.7	58 ± 23 60 ± 23	-	2.68 ± 0.5 2.62 ± 0.5	0.85 ± 0.5 0.82 ± 0.6	68 ± 19 69 ± 19	13.8 ± 6.6 15.5 ± 6.9
POLBOS II	3.6 ± 0.2 3.5 ± 0.3	1.40 ± 0.2 1.43 ± 0.3	61 ± 14 59 ± 16	9.3 ± 3.4 9.7 ±3.7	3.01 ± 0.1 3.07 ± 0.2	1.56 ± 0.2 1.57 ± 0.2	48 ± 16 49 ± 16	8.3 ±2.9 9.8 ±3.4	2.45 ± 0.4 2.34 ± 0.3	1.13 ± 0.2 1.29 ± 0.2	54 ± 24 45 ± 18	4.2 ± 2.1 3.9 ± 2.8
EBC TWO	-	1.1 ± 0.5 1.1 ± 0.5	48 ± 21 51 ± 21	18 ± 8.8 18 ± 6.7	-	-	-	-	-	0.93 ± 0.3 0.96 ± 0.4	55 ± 14 54 ± 16	10.8 ± 7.3 9.7 ± 7.1
SMART-STRATEGY	3.1 ± 0.6 3.0 ± 0.5	0.82 ± 0.5 0.77 ± 0.5	73 ± 15 75 ± 15	13.7 ± 8.1 13.2 ± 6.9	-	-	-	-	2.49 ± 0.5 2.46 ± 0.6	1.27 ± 0.7 1.29 ± 0.7	50 ± 23 49 ± 22	4.9 ± 4.0 4.4 ± 4.5
***Zhang et al.* (2016)**	3.3 ± 0.5 3.4 ± 0.6	1.16 ± 0.8 1.29 ± 0.9	65 ± 24 61 ± 27	8.1 ± 4.5 7.6 ± 4.9	3.0 ± 0.4 3.1 ± 0.5	0.85 ± 0.6 0.76 ± 0.6	72 ± 20 75 ± 21	15.8 ± 8.6 14.7 ± 7.7	2.56 ± 0.3 2.44 ± 0.4	0.56 ± 0.3 0.59 ± 0.3	78 ± 13 76 ± 11	14.1 ± 7.1 12.8 ± 4.9
***Zheng et al.* (2016)**	3.4 ± 0.4 3.3 ± 0.5	1.85 ± 0.5 1.87 ± 0.5	56 ± 9 57 ± 10	16.1 ± 6.3 18.5 ± 7.6	-	1.42 ± 0.5 1.47 ± 0.4	64 ± 8 66 ± 9	-	2.6 ± 0.3 2.7 ± 0.4	1.23 ± 0.3 1.32 ± 0.3	56 ± 10 56 ± 12	7.9 ± 4.1 7.4 ± 4.3
DKCRUSH-III	-	1.47 ± 0.4 1.49 ± 0.4	60 ± 9 59 ± 8	6.36 ± 3.7 6.97 ± 3.9	-	1.09 ± 0.4 1.07 ± 0.4	65 ± 6 66 ± 6	16.7 ± 9 18.7 ± 12	-	1.01 ± 0.4 1.07 ± 0.5	65 ± 7 63 ± 8	16.5 ± 11 17.0 ± 13
BBK I	3.1 ± 0.4 3.1 ± 0.4	1.63 ± 0.9 1.53 ± 0.8	47 ± 26 50 ± 27	20.9 ± 8.2 21.7 ± 7.5	-	1.20 ± 0.7 1.28 ± 0.7	55 ± 24 53 ± 24	-	2.38 ± 0.3 2.39 ± 0.3	2.30 ± 0.4 1.97 ± 0.5	54 ± 22 53 ± 23	9.9 ± 4.2 10.4 ± 4.1
TRYTON	2.9 ± 0.4 2.9 ± 0.4	0.99 ± 0.4 1.01 ± 0.4	66 ± 12 65 ± 11	16.8 ± 7.3 16.0 ± 6.8	-	-	-	-	2.25 ± 0.3 2.21 ± 0.3	0.95 ± 0.3 1.02 ± 0.3	58 ± 14 54 ± 14	4.8 ± 1.6 4.4 ± 1.1
POLBOS I	3.7 ± 0.2 3.6 ± 0.3	-	56 ± 16 62 ± 17	8.5 ± 2.4 8.9 ± 2.7	3.1 ± 0.4 3.2 ± 0.3	-	69 ± 13 64 ± 11	5.8 ± 1.9 7.5 ± 3.4	2.65 ± 0.3 2.43 ± 0.4	-	61 ± 17 54 ± 21	3.7 ± 1.3 2.9 ± 1.8
PERFECT	3.6 ± 0.4 3.7 ± 0.5	1.1 ± 0.4 1.1 ± 0.4	64 ± 12 66 ± 12	28.9 ± 15 27.8 ± 13	2.6 ± 0.4 2.6 ± 0.4	-	-	-	2.2 ± 0.4 2.2 ± 0.4	1.1 ± 0.4 1.1 ± 0.4	57 ± 14 53 ± 17	10.3 ± 8.2 8.3 ± 7.3
NSTS	3.0 ± 0.7 3.0 ± 0.7	1.9 ± 0.8 1.8 ± 0.8	36 ± 22 40 ± 23	17.4 ± 10 17.4 ± 10	2.59 ± 0.6 2.49 ± 0.6	1.53 ± 0.7 1.48 ± 0.7	40 ± 23;41 ± 24	-	2.39 ± 0.6 2.38 ± 0.6	1.45 ± 0.6 1.39 ± 0.7	39 ± 23 42 ± 23	7.3 ± 5.8 7.5 ± 6.0
NBS	3.0 ± 0.7 2.9 ± 0.7	1.62 ± 0.9 1.43 ± 0.8	46 ± 27 40 ± 27	23.2 ± 8.6 23.4 ± 8.6	2.63 ± 0.6 2.41 ± 0.6	1.32 ± 0.7 1.18 ± 0.7	50 ± 25 52 ± 24	-	2.28 ± 0.5 2.24 ± 0.5	1.22 ± 0.6 1.21 ± 0.6	47 ± 26 46 ± 26	6.4 ± 4.7 6.0 ± 4.8
***Ruiz et al.* (2013)**	-	1.10 ± 0.7 1.15 ± 0.8	62 ± 22 62 ± 23	-	-	1.35 ± 0.6 1.26 ± 0.7	45 ± 24 47 ± 21	-	-	1.23 ± 0.6 1.10 ± 0.4	42 ± 24 45 ± 20	-
***Ye et al.* (2012)**	2.9 ± 0.5 2.8 ± 0.5	0.97 ± 0.3 0.95 ± 0.4	64 ± 7 65 ± 9	28.9 ± 11 25.5 ± 10	-	-	-	-	2.27 ± 0.3 2.39 ± 0.5	0.97 ± 0.3 1.01 ± 0.3	70 ± 7 61 ± 8	16.9 ± 8.2 10.2 ± 8.4
BBC ONE	-	-	85 ± 11	-	-	-	-	-	-	-	68 ± 29	-

			87 ± 10								63 ± 31	
***Lin et al.* (2010)**	3.9 ± 0.4 4.0 ± 0.4	1.6 ± 0.4 1.7 ± 0.4	60 ± 8 60 ± 8	23.6 ± 2.1 23.8 ± 2.6	3.82 ± 0.5 3.91 ± 0.6	1.43 ± 0.2 1.45 ± 0.3	62 ± 5 63 ± 4	-	2.79 ± 0.2 2.82 ± 0.3	0.84 ± 0.1 0.85 ± 0.2	70 ± 5 70 ± 5	12.7 ± 2.8 12.9 ± 3.1
***Ye et al.* (2010)**	3.1 ± 0.6 3.3 ± 0.7	2.0 ± 0.7 2.1 ± 0.7	-	12.6 ± 7.2 12.3 ± 5.1	2.4 ± 0.5 2.8 ± 0.7	1.4 ± 0.6 1.6 ± 0.6	-	29.3 ± 8.3 25.3 ± 7.7	2.2 ± 0.4 2.4 ± 0.7	1.2 ± 0.4 1.3 ± 0.5	-	17.1 ± 8.0 11.5 ± 6.9
CACTUS	2.9 ± 0.3 2.7 ± 0.4	0.90 ± 0.4 0.83 ± 0.3	68 ± 12 69 ± 12	15.8 ± 8.7 14.7 ± 8.2	-	-	-	-	2.30 ± 0.3 2.16 ± 0.3	0.84 ± 0.3 0.83 ± 0.3	63 ± 12 61 ± 13	5.9 ± 4.7 5.7 ± 4.2
***Cervinka et al.* (2008)**	3.4 ± 0.4 3.5 ± 0.4	-	-	-	-	-	-	-	2.59 ± 0.4 2.53 ± 0.4	-	-	-
DKCRUSH-I	2.85 ± 0.5 2.86 ± 0.6	0.93 ± 0.5 0.94 ± 0.6	65 ± 13 64 ± 14	21.3 ± 11 20.0 ± 10	2.53 ± 0.4 2.56 ± 0.5	0.65 ± 0.3 0.62 ± 0.3	66 ± 20 62 ± 12	-	2.46 ± 0.5 2.45 ± 0.5	0.84 ± 0.6 0.84 ± 0.5	65 ± 20 66 ± 19	10.3 ± 6.3 10.5 ± 7.5
***Colombo et al.* (2004)**	2.6 ± 0.4 2.6 ± 0.5	0.99 ± 0.4 0.92 ± 0.3	62 ± 12 65 ± 11	10.8 ± 4.8 12.2 ± 5.6	-	-	-	-	2.1 ± 0.3 2.1 ± 0.3	0.88 ± 0.4 1.14 ± 0.5	57 ± 17 46 ± 22	5.5 ± 4.1 5.1 ± 4.4
***Pan et al.* (2004)**	-	0.76 ± 0.4 0,74 ± 0.5	74 ± 11 77 ± 14	-	-	-	-	-	-	0.85 ± 0.4 0.93 ± 0.4	65 ± 14 64 ± 13	-

*Lesion length including both proximal and distal main vessel

Abbreviations: DS = diameter stenosis; MLD = minimum lumen diameter; RVD = reference vessel diameter

Table S7. Risk of bias in the selected trials using the Cochrane Risk Assessment Tool.

Trial	Random Sequence Generation (Selection Bias)	Allocation Concealment (Selection Bias)	Blinding of Participants and Personnel* (Performance Bias)	Blinding of Outcome Assessment (Detection Bias)	Incomplete Outcome Data (Attrition Bias)	Selective Reporting (Reporting Bias)	Other Bias
EBC MAIN	Low	Low	Low	Low	Low	Low	Low
NBBS IV	High	High	High	High	Low	Low	Low
DEFINITION II	Low	Low	Low	Low	Low	Low	Low
DKCRUSH-V	Low	Low	Unclear	Low	Low	Low	Low
COBRA	Low	Low	Unclear	Low	Low	Low	Low
DKCRUSH-II	Unclear	Low	Unclear	Low	Low	Low	Low
BBK II	Unclear	Low	High	Low	Low	Low	Low
POLBOS II	Low	Low	High	Unclear	Low	Low	Low
EBC TWO	Low	Low	Unclear	Unclear	Low	Low	Low
SMART-STRATEGY	Unclear	Low	High	Low	Low	Low	Low
***Zhang et al.* (2016)**	Low	Low	Unclear	Unclear	Low	Low	Low
***Zheng et al.* (2016)**	Unclear	Low	Unclear	Unclear	Low	Low	Low
DKCRUSH-III	Unclear	Low	Unclear	Unclear	Low	Low	Low
BBK I	Low	Low	High	Unclear	Low	Unclear	Low
TRYTON	Low	Low	Low	Low	High	Low	Low
POLBOS I	Low	Low	High	Low	Low	Low	High
PERFECT	Low	Low	High	Unclear	Low	Low	Low
NSTS	Low	Low	High	Low	Low	Low	Low
NBS	Low	Low	High	Unclear	Low	Low	Low
***Ruiz et al.* (2013)**	Unclear	Low	Unclear	High	Unclear	Low	High
***Ye et al.* (2012)**	High	High	Unclear	Unclear	Low	Unclear	Low
BBC ONE	Low	Low	Unclear	Unclear	Low	Low	Low
***Lin et al.* (2010)**	Low	Low	Unclear	Unclear	Low	Unclear	Low
***Ye et al.* (2010)**	High	High	Unclear	Unclear	Low	Unclear	Low
CACTUS	Unclear	Low	Unclear	Low	Low	Low	Low
***Cervinka et al.* (2008)**	Low	Low	Unclear	Low	Low	Low	Low
DKCRUSH-I	Low	Low	Unclear	Low	Low	Unclear	Low
***Colombo et al.* (2004)**	Unclear	Low	Unclear	Low	Low	Unclear	High
***Pan et al.* (2004)**	High	High	Unclear	Unclear	Low	Unclear	High

*Single-blinded trials were assessed to have low risk of bias as operators could not be completely blinded due to the nature of the study

Table S8. Assessment of publication bias for each outcome in conventional meta-analysis.

	Begg and Mazumdar' Test, *P*-value	Egger's Test, *P*-value
All-cause mortality	0.93	0.44
Cardiac Death	0.09	0.40
Major Adverse Cardiac Events	0.32	0.39
Myocardial Infarction	0.43	0.29
Stent Thrombosis	0.37	0.02
Target Lesion Revascularization	0.71	0.93
Target Vessel Revascularization	0.09	0.40

Table S9. Assessment of heterogeneities in the network model.

Outcome	τ^2	I^2	Total			Within Designs			Between Designs		
			Q	**DF**	***P*-value**	***Q***	**DF**	***P*-value**	***Q***	**DF**	***P*-value**
All-cause mortality	0	0%	4.80	14	0.9884	2.59	11	0.9951	2.21	3	0.5292
Cardiac Death	0	0%	4.92	18	0.9990	4.27	13	0.9880	0.65	5	0.9856
Major Adverse Cardiac Events	0.0172	14.8%	24.64	21	0.2632	20.02	16	0.2195	4.62	5	0.4637
Myocardial Infarction	0.0366	11.4%	27.10	24	0.2997	22.79	19	0.2469	4.32	5	0.5048
Stent Thrombosis	0	0%	11.54	23	0.9770	8.48	18	0.9705	3.06	5	0.6915
Target Lesion Revascularization	0	0%	14.27	21	0.8578	10.29	16	0.8510	3.98	5	0.5526
Target Vessel Revascularization	0	0%	14.38	18	0.7042	9.38	14	0.8058	5.00	4	0.2877

Abbreviations: DF = degrees of freedom

Table S10. P-scores of each bifurcation technique for every outcome.

	Provisional	Culotte	DK Crush	DBS	T/TAP	Crush
All-cause mortality	0.6064	0.2766	0.7387	0.6744	0.4972	0.2067
Cardiac Death	0.4960	0.3649	0.8473	0.8622	0.1751	0.2545
Major Adverse Cardiac Events	0.6867	0.5076	1.0000	0.3854	0.1755	0.2448
Myocardial Infarction	0.5580	0.4343	0.9522	0.3304	0.5796	0.1455
Stent Thrombosis	0.6430	0.3731	0.9809	0.3513	0.1125	0.5392
Target Lesion Revascularization	0.6451	0.5208	1.0000	0.2306	0.1571	0.4465
Target Vessel Revascularization	0.6205	0.5899	0.9997	0.1762	0.2957	0.3180

Abbreviation: DBS = dedicated bifurcation stent; DK crush = double kissing crush; T/TAP = T-stenting or T and protrusion

Table S11. SUCRA scores of each bifurcation technique for every outcome.

	Provisional	Culotte	DK Crush	DBS	T/TAP	Crush
All-cause mortality	0.5826	0.3111	0.7048	0.6655	0.5119	0.2242
Cardiac Death	0.4923	0.3676	0.8292	0.8639	0.1853	0.2627
Major Adverse Cardiac Events	0.6759	0.5046	0.9998	0.3968	0.1721	0.2508
Myocardial Infarction	0.5278	0.4266	0.9324	0.41019	0.5663	0.1450
Stent Thrombosis	0.6474	0.3854	0.9750	0.3596	0.1264	0.5061
Target Lesion Revascularization	0.6271	0.5055	0.9999	0.2486	0.1576	0.4613
Target Vessel Revascularization	0.5822	0.5730	0.9987	0.1864	0.3228	0.3369

Abbreviation: DBS = dedicated bifurcation stent; DK crush = double kissing crush; T/TAP = T-stenting or T and protrusion

Table S12. Best rank analyses with hierarchical Bayesian model for each outcome.

		1st	2nd	3rd	4th	5th	6th
All-cause Mortality	Culotte	0.03300	0.07775	0.11763	0.20628	0.32578	0.23958
	DK Crush	0.33365	0.26843	0.15513	0.11625	0.07445	0.05210
	DBS	0.37618	0.18980	0.12650	0.11200	0.09328	0.10225
	Crush	0.01923	0.04540	0.07813	0.15545	0.28853	0.41328
	T/TAP	0.18905	0.17388	0.14315	0.17163	0.14450	0.17780
	PS	0.04890	0.24475	0.37948	0.23840	0.07348	0.01500

		1st	2nd	3rd	4th	5th	6th
Cardiac Death	Culotte	0.01343	0.06648	0.18848	0.31595	0.30783	0.10785
	DK Crush	0.28798	0.59903	0.09003	0.01778	0.00440	0.00080
	DBS	0.66385	0.17045	0.06743	0.03983	0.03675	0.02170
	Crush	0.00420	0.02993	0.09915	0.23995	0.39528	0.23150
	T/TAP	0.02810	0.05340	0.08155	0.09883	0.12485	0.61328
	PS	0.00245	0.08073	0.47338	0.28768	0.13090	0.02488

		1st	2nd	3rd	4th	5th	6th
Major Adverse Cardiac Events	Culotte	0.00003	0.22493	0.29793	0.28718	0.15508	0.03488
	DK Crush	0.99903	0.00098	0.00000	0.00000	0.00000	0.00000
	DBS	0.00088	0.17345	0.19593	0.22605	0.24593	0.15778
	Crush	0.00000	0.01795	0.08110	0.26393	0.41103	0.22600
	T/TAP	0.00008	0.06313	0.06823	0.11473	0.17348	0.58038
	PS	0.00000	0.51958	0.35683	0.10813	0.01450	0.00098

		1st	2nd	3rd	4th	5th	6th
Myocardial Infarction	Culotte	0.01085	0.14490	0.23765	0.27080	0.24463	0.09118
	DK Crush	0.72735	0.22533	0.03380	0.00978	0.00315	0.00060
	DBS	0.05663	0.16495	0.16580	0.17635	0.21653	0.21975
	Crush	0.00060	0.01100	0.04065	0.11825	0.31935	0.51015
	T/TAP	0.20100	0.28068	0.12290	0.10935	0.11648	0.16960
	PS	0.00358	0.17315	0.39920	0.31548	0.09988	0.00873

		1st	2nd	3rd	4th	5th	6th
Stent Thrombosis	Culotte	0.00573	0.07410	0.19285	0.36505	0.29340	0.06888
	DK Crush	0.89528	0.08993	0.01055	0.00333	0.00085	0.00008
	DBS	0.07113	0.15840	0.10823	0.11203	0.26023	0.29000
	Crush	0.01613	0.21670	0.28818	0.27925	0.15993	0.03983
	T/TAP	0.00220	0.01653	0.04500	0.08358	0.25273	0.59998
	PS	0.00955	0.44435	0.35520	0.15678	0.03288	0.00125

		1st	2nd	3rd	4th	5th	6th
Target Lesion Revascularization	Culotte	0.00003	0.26973	0.24548	0.27343	0.16533	0.04603
	DK Crush	0.99973	0.00028	0.00000	0.00000	0.00000	0.00000
	DBS	0.00023	0.08858	0.09708	0.14105	0.31400	0.35908
	Crush	0.00000	0.20553	0.23448	0.29223	0.19633	0.07145
	T/TAP	0.00003	0.03478	0.05028	0.10430	0.28953	0.52110
	PS	0.00000	0.40113	0.37270	0.18900	0.03483	0.00235

		1st	2nd	3rd	4th	5th	6th
Target Vessel Revascularization	Culotte	0.00090	0.41400	0.24023	0.18490	0.11398	0.04600
	DK Crush	0.99393	0.00580	0.00023	0.00005	0.00000	0.00000
	DBS	0.00053	0.04838	0.07040	0.12575	0.27323	0.48173
	Crush	0.00008	0.06155	0.16600	0.31193	0.31615	0.14430
	T/TAP	0.00450	0.18350	0.11535	0.13800	0.23550	0.32315
	PS	0.00008	0.28678	0.40780	0.23938	0.06115	0.00483

Table S13. Follow up coronary angiography and time at outcome assessment.

Trial	Angiographic Follow Up	Clinical Follow Up
EBC MAIN	N/A	1 year
NBBS IV	8 months	2 years
DEFINITION II	13 months	1 year
DKCRUSH-V	13 months	3 years
COBRA	9 months	5 years
DKCRUSH-II	8 months	5 years
BBK II	9 months	1 year
POLBOS II	12 months	1 year
EBC TWO	N/A	1 year
SMART-STRATEGY	9 months	3 years
***Zhang et al.* (2016)**	9 months	9 months
***Zheng et al.* (2016)**	12 months	1 year
DKCRUSH-III	8 months	3 years
BBK I	9 months	5 years
TRYTON	9 months	9 months
POLBOS I	12 months	1 year
PERFECT	8 months	1 year
NSTS	8 months	3 years
NBS	8 months	5 years
***Ruiz et al.* (2013)**	9 months	9 months
***Ye et al.* (2012)**	8 months	1 year
BBC ONE	N/A	9 months
***Lin et al.* (2010)**	8 months	8 months
***Ye et al.* (2010)**	8 months	8 months
CACTUS	6 months	6 months
***Cervinka et al.* (2008)**	12 months	1 year
DKCRUSH-I	8 months	8 months
***Colombo et al.* (2004)**	6 months	6 months
***Pan et al.* (2004)**	6 months	6 months

Table S14. Node-splitting analysis for each outcome.

Outcome	Comparison	*K*	Prop	*NMA*	Direct	Indirect	Difference	*Z*-value	*P*-value
All-cause Mortality	Culotte vs Crush	2	0.59	-0.0670	0.2098	-0.4617	0.6714	1.13	0.2593
	Culotte vs DBS	1	0.18	0.4225	-0.6931	0.6727	-1.3658	-1.04	0.2968
	Culotte vs PS	3	0.57	0.2922	0.1136	0.5272	-0.4136	-0.74	0.4566
	Culotte vs T/TAP	1	0.24	0.2070	0.4055	0.1443	0.2612	0.25	0.8015
	DBS vs PS	4	0.86	-0.1303	-0.3172	1.0486	-1.3658	-1.04	0.2968
	PS vs Crush	3	0.72	-0.3592	-0.5460	0.1254	-0.6714	-1.13	0.2593
	PS vs T/TAP	3	0.84	-0.0852	-0.1281	0.1331	-0.2612	-0.25	0.8015
Cardiac Death	Culotte vs Crush	2	0.66	-0.1545	-0.1324	-0.1977	0.0652	0.08	0.9374
	Culotte vs DBS	1	0.19	1.1243	0.0000	1.3844	-1.3844	-0.63	0.5278
	Culotte vs DK Crush	1	0.36	0.7791	0.6980	0.8242	-0.1262	-0.14	0.8851
	Culotte vs PS	2	0.23	0.2097	0.2682	0.1923	0.0759	0.08	0.9374
	Culotte vs T/TAP	1	0.36	-0.5517	0.0000	-0.8612	0.8612	0.49	0.6248
	DBS vs PS	4	0.85	-0.9146	-1.1254	0.2589	-1.3844	-0.63	0.5278
	DK Crush vs Crush	1	0.12	-0.9337	-1.0922	-0.9111	-0.1811	-0.15	0.8828
	DK Crush vs PS	6	0.92	-0.5694	-0.5715	-0.5445	-0.0270	-0.04	0.9721
	PS vs Crush	3	0.61	-0.3642	-0.3582	-0.3737	0.0155	0.02	0.9841
	PS vs T/TAP	2	0.70	-0.7614	1.0224	-0.1612	-0.8612	-0.49	0.6248
Major Adverse Cardiac Events	Culotte vs Crush	2	0.45	-0.1336	-0.2095	-0.0707	-0.1388	-0.47	0.6350
	Culotte vs DBS	1	0.02	-0.0641	1.6094	-0.0967	1.7062	1.11	0.2675
	Culotte vs DK Crush	1	0.30	0.7825	1.0634	0.6611	0.4023	1.14	0.2539
	Culotte vs PS	4	0.46	0.0832	0.0785	0.0872	-0.0086	-0.03	0.9745
	Culotte vs T/TAP	1	0.38	-0.2345	-0.5878	-0.0169	-0.5709	-1.12	0.2607
	DBS vs PS	4	0.99	0.1472	0.1667	-1.5395	1.7062	1.11	0.2675
	DK Crush vs Crush	1	0.26	-0.9161	-0.7408	-0.9779	0.2371	0.69	0.4875
	DK Crush vs PS	6	0.72	-0.6993	-0.6729	-0.7658	0.0928	0.33	0.7382
	PS vs Crush	4	0.72	-0.2168	-0.2239	-0.1982	-0.0258	-0.10	0.9181
	PS vs T/TAP	2	0.70	-0.3177	-0.1438	-0.7147	0.5709	1.12	0.2607
Myocardial Infarction	Culotte vs Crush	2	0.36	-0.2197	-0.2961	-0.1758	-0.1203	-0.27	0.7902
	Culotte vs DBS	1	0.04	-0.0859	1.0986	-0.1319	1.2305	0.75	0.4547
	Culotte vs DK Crush	1	0.28	0.5944	0.8921	0.4775	0.4146	0.73	0.4625
	Culotte vs PS	4	0.61	0.0767	-0.0161	0.2245	-0.2406	-0.60	0.5467
	Culotte vs T/TAP	1	0.16	0.1723	0.6931	0.0736	0.6196	0.46	0.6453
	DBS vs PS	4	0.98	0.1627	0.1912	-1.0393	1.2305	0.75	0.4547

	DK Crush vs Crush	1	0.40	-0.8141	-0.3565	-1.1225	0.7660	1.64	0.1002
	DK Crush vs PS	6	0.57	-0.5176	-0.6955	-0.2840	-0.4114	-0.96	0.3386
	PS vs Crush	4	0.73	-0.2964	-0.4020	-0.0056	-0.3965	-1.08	0.2804
	PS vs T/TAP	5	0.86	0.0956	0.0100	0.6296	-0.6196	-0.46	0.6453
Stent Thrombosis	Culotte vs Crush	2	0.73	0.1854	0.2310	0.0647	0.1663	0.25	0.7998
	Culotte vs DBS	1	0.14	-0.1501	0.0000	-0.1752	0.1752	0.08	0.9346
	Culotte vs DK Crush	1	0.14	1.0032	2.0843	0.8325	1.2518	1.10	0.2707
	Culotte vs PS	4	0.54	0.3083	-0.0551	0.7361	-0.7912	-1.26	0.2065
	Culotte vs T/TAP	1	0.10	-0.5529	1.0986	-0.7400	1.8386	1.07	0.2847
	DBS vs PS	4	0.88	0.4584	0.4797	0.3045	0.1752	0.08	0.9346
	DK Crush vs Crush	1	0.23	-0.8178	-0.9099	-0.7910	-0.1189	-0.13	0.8996
	DK Crush vs PS	5	0.84	-0.6949	-0.5874	-1.2717	0.6843	0.87	0.3849
	PS vs Crush	4	0.47	-0.1229	-0.1794	-0.0723	-0.1071	-0.16	0.8694
	PS vs T/TAP	5	0.93	-0.8612	-0.9855	0.8530	-1.8386	-1.07	0.2847
Target Lesion Revascularization	Culotte vs Crush	2	0.39	-0.0418	-0.1169	0.0066	-0.1236	-0.31	0.7547
	Culotte vs DBS	1	0.03	-0.2152	1.0986	-0.2540	1.3526	0.83	0.4058
	Culotte vs DK Crush	1	0.26	0.8872	1.2927	0.7442	0.5485	1.22	0.2231
	Culotte vs PS	3	0.51	0.0604	0.0776	0.0425	0.0351	0.11	0.9150
	Culotte vs T/TAP	1	0.36	-0.2713	-0.6931	-0.0343	-0.6589	-1.35	0.1782
	DBS vs PS	4	0.98	0.2756	0.3005	-1.0521	1.3526	0.83	0.4058
	DK Crush vs Crush	1	0.37	-0.9290	-0.7218	-1.0531	0.3313	0.86	0.3904
	DK Crush vs PS	6	0.71	-0.8268	-0.8113	-0.8645	0.0531	0.16	0.8731
	PS vs Crush	3	0.61	-0.1022	-0.1616	-0.0085	-0.1531	-0.47	0.6391
	PS vs T/TAP	4	0.77	-0.3317	-0.1800	0.8389	0.6589	1.35	0.1782
Target Vessel Revascularization	Culotte vs Crush	2	0.53	-0.1634	-0.2146	-0.1060	-0.1086	-0.31	0.7588
	Culotte vs DBS	1	0.05	-0.3027	0.6931	-0.3560	1.0491	0.86	0.3877
	Culotte vs DK Crush	1	0.40	0.7409	1.1835	0.4510	0.7325	1.81	0.0708
	Culotte vs PS	3	0.37	-0.0014	-0.3762	0.2234	-0.5995	-1.61	0.1084
	DBS vs PS	4	0.97	0.3013	0.3308	-0.7183	1.0491	0.86	0.3877
	DK Crush vs Crush	1	0.39	-0.9042	-0.7473	-1.0052	0.2578	0.72	0.4744
	DK Crush vs PS	3	0.58	-0.7422	-0.6233	-0.9040	0.2807	0.83	0.4049
	PS vs Crush	4	0.71	-0.1620	-0.1919	-0.0885	-0.1034	-0.34	0.7340

Table S15. Pooled estimates of network meta-analysis for each outcome.

All-cause Mortality (19 trials)	Culotte	0.65 (0.31-1.37)	0.66 (0.24-1.77)	0.81 (0.34-1.94)	1.07 (0.60-1.90)	0.75 (0.44-1.28)
	1.55 (0.73-3.28)	DK Crush	1.02 (0.36-2.83)	1.26 (0.50-3.15)	1.66 (0.79-3.46)	1.16 (0.69-1.95)
	1.53 (0.57-4.12)	0.98 (0.35-2.74)	DBS	1.24 (0.39-3.93)	1.63 (0.59-4.47)	1.14 (0.47-2.75)
	1.23 (0.52-2.94)	0.79 (0.32-1.98)	0.81 (0.25-2.56)	T/TAP	1.32 (0.54-3.22)	0.92 (0.43-1.95)
	0.94 (0.53-1.66)	0.60 (0.29-1.26)	0.61 (0.22-1.68)	0.76 (0.31-1.86)	Crush	0.70 (0.41-1.18)
	1.34 (0.78-2.30)	0.86 (0.51-1.45)	0.88 (0.36-2.12)	1.09 (0.51-2.32)	1.43 (0.85-2.42)	Provisional
Cardiac Death (23 trials)	Culotte	0.46 (0.20-1.04)	0.32 (0.06-1.74)	1.74 (0.33-9.10)	1.17 (0.54-2.52)	0.81 (0.37-1.80)
	2.18 (0.96-4.95)	DK Crush	0.71 (0.14-3.47)	3.78 (0.75-19.20)	2.54 (1.15-5.64)	1.77 (1.19-2.63)
	3.08 (0.57-16.50)	1.41 (0.29-6.92)	DBS	5.34 (0.59-48.20)	3.59 (0.66-19.41)	2.50 (0.53-11.69)
	0.58 (0.11-3.02)	0.26 (0.05-1.34)	0.19 (0.02-1.69)	T/TAP	0.67 (0.12-3.67)	0.47 (0.10-2.28)
	0.86 (0.40-1.85)	0.39 (0.18-0.87)	0.28 (0.05-1.50)	1.49 (0.27-8.12)	Crush	0.69 (0.33-1.46)
	1.23 (0.56-2.73)	0.57 (0.38-0.84)	0.40 (0.09-1.88)	2.14 (0.44-10.46)	1.44 (0.68-3.03)	Provisional
Major Adverse Cardiac Events (26 trials)	Culotte	0.46 (0.33-0.63)	1.07 (0.71-1.61)	1.26 (0.78-2.05)	1.14 (0.86-1.52)	0.92 (0.71-1.20)
	2.19 (1.59-3.00)	DK Crush	2.33 (1.56-3.49)	2.76 (1.66-3.49)	2.50 (1.86-3.35)	2.01 (1.57-2.57)
	0.94 (0.62-1.42)	0.43 (0.29-0.64)	DBS	1.19 (0.68-2.07)	1.07 (0.73-1.58)	0.86 (0.63-1.19)
	0.79 (0.49-1.28)	0.36 (0.22-0.60)	0.84 (0.48-1.47)	T/TAP	0.90 (0.55-1.48)	0.73 (0.46-1.15)
	0.87 (0.66-1.16)	0.40 (0.30-0.54)	0.93 (0.63-1.38)	1.11 (0.67-1.82)	Crush	0.81 (0.65-1.00)
	1.09 (0.83-1.42)	0.50 (0.39-0.64)	1.16 (0.84-1.60)	1.37 (0.87-2.17)	1.24 (1.00-1.55)	Provisional
Myocardial Infarction (29 trials)	Culotte	0.55 (0.39-0.91)	1.09 (0.59-2.01)	0.84 (0.32-2.21)	1.25 (0.81-1.91)	0.93 (0.63-1.36)
	1.81 (1.10-2.98)	DK Crush	1.97 (1.04-3.74)	1.53 (0.56-4.13)	2.26 (1.44-3.53)	1.68 (1.11-2.55)
	0.92 (0.50-1.69)	0.51 (0.27-0.96)	DBS	0.77 (0.28-2.17)	1.14 (0.64-2.04)	0.85 (0.52-1.38)
	1.19 (0.45-3.12)	0.66 (0.24-1.78)	1.29 (0.46-3.63)	T/TAP	1.48 (0.57-3.86)	1.10 (0.44-2.73)
	0.80 (0.52-1.23)	0.44 (0.28-0.69)	0.87 (0.49-1.56)	0.68 (0.26-1.76)	Crush	0.74 (0.54-1.02)
	1.08 (0.74-1.58)	0.60 (0.39-0.90)	1.18 (0.72-1.91)	0.91 (0.37-2.26)	1.35 (0.98-1.85)	Provisional
Stent Thrombosis (28 trials)	Culotte	0.37 (0.17-0.79)	1.16 (0.27-5.04)	1.74 (0.63-4.81)	0.83 (0.47-1.47)	0.73 (0.40-1.35)
	2.73 (1.27-5.86)	DK Crush	3.17 (0.73-13.3)	4.74 (1.72-13.04)	2.27 (1.05-4.90)	2.00 (1.14-3.51)
	0.86 (0.20-3.73)	0.32 (0.07-1.38)	DBS	1.50 (0.30-7.45)	0.71 (0.16-3.17)	0.63 (0.16-2.48)
	0.58 (0.21-1.59)	0.21 (0.08-0.58)	0.67 (0.13-3.33)	T/TAP	0.48 (0.17-1.36)	0.42 (0.18-0.98)
	1.20 (0.68-2.14)	0.44 (0.20-0.95)	1.40 (0.32-6.20)	2.09 (0.74-5.94)	Crush	0.88 (0.47-1.67)
	1.36 (0.74-2.51)	0.50 (0.28-0.88)	1.58 (0.40-6.21)	2.37 (1.02-5.51)	1.13 (0.60-2.14)	Provisional
Target Lesion Revascularization (26 trials)	Culotte	0.41 (0.28-0.61)	1.24 (0.73-2.11)	1.31 (0.83-2.08)	1.04 (0.71-1.52)	0.94 (0.68-1.30)
	2.43 (1.65-3.58)	DK Crush	3.01 (1.79-5.07)	3.19 (1.95-5.20)	2.53 (1.76-3.65)	2.29 (1.70-3.07)
	0.81 (0.47-1.37)	0.33 (0.20-0.56)	DBS	1.06 (0.59-1.90)	0.84 (0.50-1.43)	0.76 (0.49-1.17)

	0.76 (0.48-1.21)	0.31 (0.19-0.51)	0.95 (0.53-1.70)	T/TAP	0.79 (0.48-1.30)	0.72 (0.48-1.07)
	0.96 (0.66-1.40)	0.39 (0.27-0.57)	1.19 (0.70-2.02)	1.26 (0.77-2.06)	Crush	0.90 (0.66-1.23)
	1.06 (0.77-1.47)	0.44 (0.33-0.59)	1.32 (0.86-2.02)	1.39 (0.93-2.09)	1.11 (0.81-1.51)	Provisional
Target Vessel Revascularization (23 trials)	Culotte	0.48 (0.32-0.70)	1.35 (0.80-2.28)	1.25 (0.66-2.35)	1.18 (0.83-1.66)	1.00 (0.70-1.43)
	2.10 (1.42-3.09)	DK Crush	2.84 (1.71-4.72)	2.61 (1.41-4.86)	2.47 (1.75-3.49)	2.10 (1.52-2.91)
	0.74 (0.44-1.25)	0.35 (0.21-0.59)	DBS	0.92 (0.48-1.78)	0.87 (0.54-1.40)	0.74 (0.50-1.10)
	0.80 (0.43-1.51)	0.38 (0.21-0.71)	1.09 (0.56-2.09)	T/TAP	0.94 (0.52-1.71)	0.80 (0.47-1.36)
	0.85 (0.60-1.20)	0.40 (0.29-0.57)	1.15 (0.71-1.85)	1.06 (0.59-1.91)	Crush	0.85 (0.65-1.11)
	1.00 (0.70-1.42)	0.48 (0.34-0.66)	1.35 (0.91-2.00)	1.24 (0.74-2.11)	1.18 (0.90-1.54)	Provisional

Table S16. Sensitivity analysis of trials that only included true bifurcations.

All-cause Mortality (10 trials)	Culotte	0.77 (0.32-1.87)	0.88 (0.20-3.87)	0.55 (0.05-5.82)	2.00 (0.18-21.82)	0.89 (0.44-1.83)
	1.30 (0.54-3.14)	DK Crush	1.14 (0.28-4.62)	0.72 (0.07-7.16)	2.59 (0.20-33.13)	1.16 (0.69-1.95)
	1.14 (0.26-5.01)	0.88 (0.22-3.56)	DBS	0.63 (0.05-8.40)	2.27 (0.14-37.88)	1.02 (0.28-3.72)
	1.81 (0.17-18.99)	1.39 (0.14-13.92)	1.59 (0.12-21.18)	T/TAP	3.61 (0.13-103.4)	1.61 (0.17-15.17)
	0.50 (0.05-5.46)	0.41 (0.03-5.18)	0.56 (0.04-8.28)	0.29 (0.01-8.33)	Crush	0.47 (0.04-5.68)
	1.12 (0.55-2.29)	0.86 (0.51-1.45)	0.98 (0.27-3.61)	0.62 (0.07-5.83)	2.24 (0.18-27.15)	Provisional
Cardiac Death (12 trials)	Culotte	0.52 (0.19-1.40)	0.95 (0.09-9.87)	N/A	1.21 (0.25-5.89)	0.93 (0.34-2.54)
	1.93 (0.72-5.21)	DK Crush	1.83 (0.18-18.13)	N/A	2.34 (0.52-10.58)	1.79 (1.20-2.69)
	1.06 (0.10-11.01)	0.55 (0.06-5.43)	DBS	N/A	1.28 (0.09-18.75)	0.98 (0.10-9.52)
	N/A	N/A	N/A	T/TAP	N/A	N/A
	0.83 (0.17-4.02)	0.43 (0.09-1.94)	0.78 (0.05-11.47)	N/A	Crush	0.77 (0.17-3.52)
	1.08 (0.39-2.94)	0.56 (0.37-0.84)	1.02 (0.11-9.90)	N/A	1.30 (0.28-5.99)	Provisional
Major Adverse Cardiac Events (14 trials)	Culotte	0.72 (0.33-1.57)	1.29 (0.47-3.56)	N/A	1.13 (0.47-2.70)	1.00 (0.55-1.81)
	1.40 (0.64-3.05)	DK Crush	1.80 (0.68-4.74)	N/A	1.57 (0.59-4.17)	1.40 (0.84-2.33)
	0.77 (0.28-2.14)	0.56 (0.21-1.46)	DBS	N/A	0.87 (0.27-2.81)	0.78 (0.34-1.76)
	N/A	N/A	N/A	T/TAP	N/A	N/A
	1.07 (0.71-1.61)	0.46 (0.32-0.65)	1.23 (0.76-2.00)	N/A	Crush	0.91 (0.64-1.29)
	1.00 (0.55-1.81)	0.72 (0.43-1.19)	1.29 (0.57-2.94)	N/A	1.13 (0.49-2.59)	Provisional
Myocardial Infarction (16 trials)	Culotte	0.55 (0.34-0.89)	1.35 (0.80-2.30)	1.19 (0.28-5.06)	1.05 (0.61-1.81)	0.99 (0.68-1.44)
	1.82 (1.13-2.94)	DK Crush	2.46 (1.42-4.28)	2.17 (0.51-9.27)	1.92 (1.21-3.04)	1.81 (1.21-2.69)
	0.74 (0.43-1.26)	0.41 (0.23-0.70)	DBS	0.88 (0.21-3.75)	0.78 (0.43-1.42)	0.73 (0.50-1.08)
	0.39 (0.10-1.54)	0.18 (0.05-0.67)	0.54 (0.06-4.60)	T/TAP	0.57 (0.12-2.70)	0.36 (0.11-1.14)
	0.95 (0.55-1.63)	0.52 (0.33-0.83)	1.29 (0.71-2.35)	1.13 (0.26-4.93)	Crush	0.94 (0.59-1.50)
	1.01 (0.69-1.46)	0.55 (0.37-0.82)	1.36 (0.93-2.00)	1.20 (0.30-4.85)	1.06 (0.67-1.69)	Provisional
Stent Thrombosis (15 trials)	Culotte	0.47 (0.20-1.09)	1.36 (0.21-9.03)	2.54 (0.65-9.95)	1.45 (0.49-4.33)	0.91 (0.44-1.87)
	2.15 (0.92-5.03)	DK Crush	2.93 (0.44-19.43)	5.46 (1.50-19.86)	3.12 (1.08-8.98)	1.95 (1.11-3.46)
	0.73 (0.11-4.86)	0.34 (0.05-2.27)	DBS	1.86 (0.22-16.01)	1.07 (0.14-8.40)	0.67 (0.11-4.09)
	0.39 (0.10-1.54)	0.18 (0.05-0.67)	0.54 (0.06-4.60)	T/TAP	0.57 (0.12-2.70)	0.36 (0.11-1.14)
	0.69 (0.23-2.06)	0.32 (0.11-0.92)	0.94 (0.12-7.40)	1.75 (0.37-8.27)	Crush	0.63 (0.22-1.76)
	1.10 (0.53-2.26)	0.51 (0.29-0.90)	1.50 (0.24-9.17)	2.79 (0.88-8.90)	1.60 (0.57-4.49)	Provisional
Target Lesion Revascularization (15 trials)	Culotte	0.42 (0.24-0.73)	1.30 (0.60-2.81)	2.00 (0.40-10.03)	1.18 (0.60-2.31)	0.95 (0.62-1.45)
	2.37 (1.37-4.12)	DK Crush	3.08 (1.48-6.42)	4.76 (0.97-23.39)	2.79 (1.35-5.79)	2.25 (1.58-3.20)
	0.77 (0.36-1.67)	0.32 (0.16-0.68)	DBS	1.54 (0.29-8.30)	0.91 (0.37-2.24)	0.73 (0.38-1.39)

	0.50 (0.10-2.50)	0.21 (0.04-1.03)	0.65 (0.12-3.48)	T/TAP	0.59 (0.11-3.15)	0.47 (0.10-2.24)
	1.08 (0.64-1.84)	0.41 (0.26-0.65)	1.22 (0.55-2.66)	1.97 (0.38-10.01)	Crush	0.93 (0.58-1.49)
	1.06 (0.69-1.61)	0.44 (0.31-0.63)	1.37 (0.72-2.61)	2.11 (0.45-9.99)	1.24 (0.65-2.35)	Provisional
Target Vessel Revascularization (12 trials)	Culotte	0.76 (0.38-1.49)	2.00 (0.85-4.72)	1.64 (0.45-5.94)	1.41 (0.72-2.75)	1.41 (0.84-2.37)
	1.32 (0.67-2.60)	DK Crush	2.65 (1.19-5.92)	2.17 (0.62-7.60)	1.86 (0.88-3.93)	1.87 (1.21-2.87)
	0.50 (0.21-1.18)	0.38 (0.17-0.84)	DBS	0.82 (0.21-3.19)	0.70 (0.28-1.75)	0.70 (0.36-1.39)
	0.61 (0.17-2.22)	0.46 (0.13-1.62)	1.22 (0.31-4.77)	T/TAP	0.86 (0.23-3.24)	0.86 (0.26-2.80)
	0.95 (0.54-1.66)	0.43 (0.27-0.70)	1.25 (0.53-2.97)	1.13 (0.31-4.16)	Crush	0.97 (0.59-1.62)
	0.71 (0.42-1.20)	0.54 (0.35-0.82)	1.42 (0.72-2.80)	1.16 (0.36-3.77)	1.00 (0.54-1.84)	Provisional

Table S17. Sensitivity analysis excluding trials that did not include left main bifurcations.

All-cause Mortality (11 trials)	Culotte	0.60 (0.27-1.32)	0.36 (0.08-1.62)	0.53 (0.11-2.49)	1.01 (0.55-1.84)	0.69 (0.38-1.26)
	1.67 (0.76-3.68)	DK Crush	0.60 (0.13-2.63)	0.88 (0.16-4.78)	1.69 (0.78-3.65)	1.16 (0.69-1.95)
	2.80 (0.62-12.71)	1.68 (0.38-7.41)	DBS	1.47 (0.17-12.41)	2.83 (0.63-12.74)	1.94 (0.48-7.80)
	1.90 (0.40-9.01)	1.14 (0.21-6.21)	0.68 (0.08-5.72)	T/TAP	1.92 (0.37-9.90)	1.32 (0.26-6.62)
	0.99 (0.54-1.80)	0.59 (0.27-1.28)	0.35 (0.08-1.59)	0.52 (0.10-2.68)	Crush	0.69 (0.39-1.22)
	1.44 (0.80-2.61)	0.86 (0.51-1.45)	0.51 (0.13-2.07)	0.76 (0.15-3.81)	1.46 (0.82-2.58)	Provisional
Cardiac Death (14 trials)	Culotte	0.42 (0.18-0.98)	0.12 (0.01-1.20)	1.92 (0.32-11.47)	1.16 (0.53-2.56)	0.73 (0.32-1.69)
	2.39 (1.02-5.57)	DK Crush	0.30 (0.03-2.54)	4.59 (0.78-27.00)	2.78 (1.18-6.51)	1.75 (1.16-2.63)
	8.09 (0.84-78.34)	3.39 (0.39-29.14)	DBS	15.56 (1.0-240.1)	9.41 (0.98-90.55)	5.93 (0.72-49.04)
	0.52 (0.09-3.10)	0.22 (0.04-1.28)	0.60 (0.00-0.99)	T/TAP	0.60 (0.10-3.83)	0.38 (0.07-2.17)
	0.86 (0.39-1.89)	0.36 (0.15-0.84)	0.11 (0.01-1.02)	1.65 (0.26-10.47)	Crush	0.63 (0.28-1.42)
	1.36 (0.59-3.14)	0.57 (0.38-0.86)	0.17 (0.02-1.39)	2.62 (0.46-14.94)	1.59 (0.70-3.58)	Provisional
Major Adverse Cardiac Events (15 trials)	Culotte	0.46 (0.34-0.62)	0.79 (0.45-1.37)	2.09 (1.04-4.19)	1.14 (0.861.51)	0.84 (0.64-1.10)
	2.16 (1.61-2.91)	DK Crush	1.70 (1.00-2.91)	4.51 (2.13-9.55)	2.47 (1.82-3.35)	1.82 (1.44-2.30)
	1.27 (0.73-2.20)	0.59 (0.34-1.00)	DBS	2.65 (1.10-6.38)	1.45 (0.83-2.53)	1.07 (0.66-1.73)
	0.48 (0.24-0.96)	0.22 (0.10-0.47)	0.38 (0.16-0.91)	T/TAP	0.55 (0.26-1.15)	0.40 (0.19-0.84)
	0.87 (0.66-1.16)	0.40 (0.30-0.55)	0.69 (0.39-1.20)	1.83 (0.87-3.85)	Crush	0.73 (0.56-0.97)
	1.19 (0.91-1.56)	0.55 (0.43-0.70)	0.94 (0.58-1.52)	2.48 (1.19-5.18)	1.36 (1.03-1.80)	Provisional
Myocardial Infarction (16 trials)	Culotte	0.60 (0.35-1.03)	0.61 (0.16-2.35)	1.54 (0.28-8.51)	1.27 (0.74-2.19)	1.07 (0.67-1.70)
	1.67 (0.97-2.88)	DK Crush	1.02 (0.26-3.95)	2.57 (0.44-14.84)	2.13 (1.23-3.69)	1.78 (1.10-2.88)
	1.63 (0.42-6.27)	0.98 (0.25-3.77)	DBS	2.51 (0.30-20.99)	2.08 (0.52-8.24)	1.74 (0.49-6.14)
	0.65 (0.12-3.61)	0.39 (0.07-2.25)	0.40 (0.05-3.34)	T/TAP	0.83 (0.14-4.85)	0.69 (0.13-3.83)
	0.78 (0.46-1.35)	0.47 (0.27-0.82)	0.48 (0.12-1.91)	1.21 (0.21-7.04)	Crush	0.84 (0.48-1.45)
	0.94 (0.59-1.50)	0.56 (0.35-0.91)	0.58 (0.16-2.03)	1.44 (0.26-7.97)	1.20 (0.69-2.07)	Provisional
Stent Thrombosis (16 trials)	Culotte	0.45 (0.20-1.00)	1.51 (0.17-13.58)	2.26 (0.65-7.90)	0.76 (0.42-1.38)	0.95 (0.48-1.88)
	2.25 (1.00-5.04)	DK Crush	3.39 (0.39-29.60)	5.08 (1.48-17.46)	1.70 (0.73-3.97)	2.12 (1.19-3.80)
	0.66 (0.07-5.95)	0.29 (0.03-2.57)	DBS	1.50 (0.14-15.82)	0.50 (0.05-4.61)	0.63 (0.08-5.04)
	0.44 (0.13-1.54)	0.20 (0.06-0.68)	0.67 (0.06-7.05)	T/TAP	0.33 (0.09-1.23)	0.42 (0.14-1.25)
	1.32 (0.73-2.41)	0.59 (0.25-1.38)	2.00 (0.22-18.39)	2.99 (0.81-11.01)	Crush	1.25 (0.59-2.67)
	1.06 (0.53-2.10)	0.47 (0.26-0.84)	1.60 (0.20-12.88)	2.39 (0.80-7.17)	0.80 (0.38-1.71)	Provisional
Target Lesion Revascularization (16 trials)	Culotte	0.43 (0.29-0.64)	1.19 (0.61-2.33)	1.55 (0.89-2.68)	1.07 (0.72-1.60)	0.89 (0.64-1.24)
	2.34 (1.57-3.49)	DK Crush	2.78 (1.43-5.41)	3.62 (1.97-6.65)	2.51 (1.70-3.72)	2.08 (1.51-2.86)
	0.84 (0.43-1.65)	0.36 (0.18-0.70)	DBS	1.30 (0.59-2.89)	0.90 (0.46-1.79)	0.74 (0.42-1.34)

	0.65 (0.37-1.12)	0.28 (0.15-0.51)	0.77 (0.35-1.70)	T/TAP	0.69 (0.37-1.29)	0.57 (0.33-0.99)
	0.93 (0.63-1.39)	0.40 (0.27-0.59)	1.11 (0.56-2.19)	1.44 (0.77-2.68)	Crush	0.83 (0.58-1.18)
	1.13 (0.81-1.57)	0.48 (0.35-0.66)	1.34 (0.47-2.40)	1.74 (1.01-2.99)	1.21 (0.84-1.73)	Provisional
Target Vessel Revascularization (12 trials)	Culotte	0.49 (0.31-0.77)	1.27 (0.64-2.55)	1.32 (0.60-2.90)	1.19 (0.79-1.79)	0.93 (0.60-1.43)
	2.03 (1.30-3.19)	DK Crush	2.59 (1.31-5.13)	2.68 (1.23-5.86)	2.42 (1.58-3.72)	1.88 (1.23-2.87)
	0.78 (0.39-1.57)	0.39 (0.19-0.77)	DBS	1.04 (0.44-2.42)	0.94 (0.48-1.83)	0.73 (0.42-1.25)
	0.76 (0.34-1.67)	0.37 (0.17-0.82)	0.97 (0.41-2.26)	T/TAP	0.90 (0.42-1.96)	0.70 (0.36-1.36)
	0.84 (0.56-1.26)	0.41 (0.27-0.63)	1.07 (0.55-2.09)	1.11 (0.51-2.39)	Crush	0.78 (0.52-1.16)
	1.08 (0.70-1.67)	0.53 (.0.35-0.81)	1.37 (0.80-2.35)	1.42 (0.74-2.75)	1.29 (0.86-1.93)	Provisional

Table S18. Sensitivity analysis of trials that included only non-left main bifurcations.

All-cause Mortality (7 trials)	Culotte	N/A	1.86 (0.12-28.03)	2.33 (0.18-29.33)	3.06 (0.19-49.33)	1.88 (0.17-20.44)
	N/A	DK Crush	N/A	N/A	N/A	N/A
	0.54 (0.04-8.14)	N/A	DBS	1.26 (0.27-5.94)	1.65 (0.24-11.38)	1.02 (0.28-3.72)
	0.43 (0.03-5.41)	N/A	0.80 (0.17-3.77)	T/TAP	1.32 (0.25-6.95)	0.81 (0.34-1.90)
	0.33 (0.02-5.26)	N/A	0.61 (0.09-4.17)	0.76 (0.14-4.02)	Crush	0.61 (0.15-2.56)
	0.53 (0.05-5.76)	N/A	0.98 (0.27-3.61)	1.24 (0.53-2.91)	1.63 (0.39-6.78)	Provisional
Cardiac Death (8 trials)	Culotte	N/A	N/A	N/A	N/A	N/A
	N/A	DK Crush	1.68 (0.05-53.06)	1.55 (0.02-127.3)	1.72 (0.10-29.10)	1.69 (0.21-13.52)
	N/A	0.60 (0.02-18.89)	DBS	0.92 (0.01-108.7)	1.03 (0.04-29.57)	1.01 (0.06-15.93)
	N/A	0.65 (0.01-53.31)	1.08 (0.01-127.9)	T/TAP	1.12 (0.01-85.22)	1.09 (0.02-53.52)
	N/A	0.58 (0.03-9.80)	0.97 (0.03-27.97)	0.90 (0.01-68.53)	Crush	0.98 (0.14-6.66)
	N/A	0.59 (0.07-4.73)	0.99 (0.06-15.66)	0.91 (0.02-44.74)	1.02 (0.15-6.92)	Provisional
Major Adverse Cardiac Events (10 trials)	Culotte	0.20 (0.06-0.64)	1.04 (0.40-2.70)	0.75 (0.27-2.09)	0.90 (0.36-2.27)	0.75 (0.31-1.83)
	4.96 (1.55-15.80)	DK Crush	5.18 (2.28-11.74)	3.73 (1.51-9.21)	4.45 (2.01-9.83)	3.73 (1.77-7.87)
	0.96 (0.37-2.47)	0.19 (0.09-0.44)	DBS	0.72 (0.39-1.33)	0.86 (0.56-1.32)	0.72 (0.51-1.01)
	1.33 (0.48-3.69)	0.27 (0.11-0.66)	1.39 (0.75-2.55)	T/TAP	1.19 (0.67-2.12)	1.00 (0.60-1.66)
	1.11 (0.44-2.81)	0.22 (0.10-0.50)	1.16 (0.76-1.79)	0.84 (0.47-1.49)	Crush	0.84 (0.64-1.10)
	1.33 (0.55-3.22)	0.27 (0.13-0.56)	1.39 (0.99-1.94)	1.00 (0.60-1.66)	1.19 (0.91-1.56)	Provisional
Myocardial Infarction (12 trials)	Culotte	0.21 (0.02-2.15)	0.65 (0.18-2.37)	0.34 (0.07-1.67)	0.67 (0.20-2.29)	0.47 (0.15-1.47)
	4.85 (0.47-50.54)	DK Crush	3.16 (0.37-26.77)	1.67 (0.16-16.98)	3.27 (0.40-26.55)	2.28 (0.29-17.70)
	1.53 (0.42-5.58)	0.32 (0.04-2.67)	DBS	0.53 (0.15-1.84)	1.03 (0.49-2.19)	0.72 (0.39-1.32)
	2.91 (0.60-14.15)	0.60 (0.06-6.12)	1.90 (0.54-6.64)	T/TAP	1.96 (0.60-6.39)	1.37 (0.46-4.10)
	1.48 (0.44-5.04)	0.31 (0.04-2.49)	0.97 (0.46-2.05)	0.51 (0.16-1.66)	Crush	0.70 (0.45-1.09)
	2.12 (0.68-6.64)	0.44 (0.06-3.39)	1.39 (0.76-2.54)	0.73 (0.24-2.18)	1.43 (0.92-2.23)	Provisional
Stent Thrombosis (11 trials)	Culotte	0.15 (0.01-4.15)	0.51 (0.02-10.65)	0.70 (0.05-9.50)	0.78 (0.06-10.24)	0.31 (0.03-2.97)
	6.75 (0.24-189.0)	DK Crush	3.46 (0.14-84.60)	4.72 (0.29-77.39)	5.27 (0.33-83.63)	2.12 (0.18-24.84)
	1.94 (0.09-40.56)	0.29 (0.01-7.07)	DBS	1.36 (0.12-15.56)	1.52 (0.14-16.73)	0.61 (0.08-4.71)
	1.43 (0.11-19.47)	0.21 (0.01-3.48)	0.73 (0.06-8.37)	T/TAP	0.73 (0.06-8.37)	1.12 (0.18-6.97)
	1.28 (0.10-16.79)	0.19 (0.01-3.01)	0.66 (0.06-7.20)	0.89 (0.14-5.57)	Crush	0.40 (0.11-1.41)
	3.19 (0.34-30.11)	0.47 (0.04-5.53)	1.63 (0.21-12.55)	2.23 (0.59-8.41)	2.49 (0.71-8.74)	Provisional
Target Lesion Revascularization (9 trials)	Culotte	N/A	N/A	N/A	N/A	N/A
	N/A	DK Crush	5.54 (1.85-16.62)	4.32 (1.47-12.68)	3.78 (1.25-11.39)	4.05 (1.66-9.86)
	N/A	0.18 (0.06-0.54)	DBS	0.78 (0.32-1.89)	0.68 (0.27-1.71)	0.73 (0.38-1.39)

	N/A	0.23 (0.08-0.68)	1.28 (0.53-3.11)	T/TAP	0.87 (0.36-2.13)	0.94 (0.51-1.72)
	N/A	0.26 (0.09-0.80)	1.47 (0.59-3.67)	1.14 (0.47-2.79)	Crush	1.07 (0.56-2.06)
	N/A	0.25 (0.10-0.60)	1.37 (0.72-2.61)	1.07 (0.58-1.96)	0.93 (0.49-1.79)	Provisional
Target Vessel Revascularization (10 trials)	Culotte	0.71 (0.06-8.11)	4.01 (0.38-41.95)	2.20 (0.18-26.81)	3.03 (0.31-29.99)	2.83 (0.30-26.70)
	1.40 (0.12-15.97)	DK Crush	5.63 (1.78-17.86)	3.08 (0.73-13.05)	4.26 (1.51-12.00)	3.96 (1.56-10.07)
	0.25 (0.02-2.60)	0.18 (0.06-0.56)	DBS	0.55 (0.15-2.00)	0.76 (0.33-1.71)	0.70 (0.36-1.39)
	0.45 (0.04-5.55)	0.32 (0.08-1.37)	1.83 (0.50-6.66)	T/TAP	1.38 (0.42-4.53)	1.29 (0.43-3.86)
	0.33 (0.03-3.26)	0.23 (0.08-0.66)	1.32 (0.59-2.99)	0.72 (0.22-2.38)	Crush	0.93 (0.59-1.46)
	0.35 (0.04-3.35)	0.25 (0.10-0.64)	1.42 (0.72-2.80)	0.78 (0.26-2.34)	1.07 (0.68-1.69)	Provisional

Table S19. Sensitivity analysis excluding trials with multiple bifurcation techniques in either arm.

All-cause Mortality (14 trials)	Culotte	0.87 (0.25-3.06)	0.82 (0.23-2.94)	1.01 (0.32-3.22)	0.88 (0.43-1.79)	0.97 (0.33-2.85)
	1.15 (0.33-4.03)	DK Crush	0.94 (0.31-2.83)	1.16 (0.42-3.18)	1.01 (0.27-3.77)	1.12 (0.58-2.14)
	1.22 (0.34-4.34)	1.06 (0.35-3.20)	DBS	1.23 (0.39-3.90)	1.07 (0.28-4.14)	1.18 (0.48-2.89)
	0.99 (0.31-3.15)	0.86 (0.31-2.37)	0.81 (0.26-2.58)	T/TAP	0.87 (0.25-3.04)	0.96 (0.44-2.08)
	1.13 (0.56-2.31)	0.99 (0.27-3.69)	0.93 (0.24-3.60)	1.15 (0.33-4.00)	Crush	1.10 (0.35-3.47)
	1.03 (0.35-3.01)	0.90 (0.47-1.72)	0.85 (0.35-2.06)	1.04 (0.48-2.25)	0.91 (0.29-2.85)	Provisional
Cardiac Death (19 trials)	Culotte	0.44 (0.17-1.17)	0.34 (0.06-1.94)	1.79 (0.33-9.81)	1.11 (0.48-2.57)	0.84 (0.31-2.27)
	2.26 (0.86-5.94)	DK Crush	0.76 (0.15-3.75)	4.03 (0.79-20.67)	2.50 (0.87-7.20)	1.91 (1.22-2.98)
	2.97 (0.52-17.11)	1.32 (0.27-6.52)	DBS	5.31 (0.59-47.98)	3.29 (0.53-20.31)	2.51 (0.53-11.80)
	0.56 (0.10-3.08)	0.25 (0.05-1.27)	0.19 (0.02-1.70)	T/TAP	0.62 (0.10-3.75)	0.47 (0.10-2.34)
	0.90 (0.39-2.10)	0.40 (0.14-1.15)	0.30 (0.05-1.87)	1.61 (0.27-9.75)	Crush	0.76 (0.26-2.22)
	1.18 (0.44-3.18)	0.52 (0.34-0.82)	0.40 (0.08-1.87)	2.11 (0.43-10.44)	1.31 (0.45-3.81)	Provisional
Major Adverse Cardiac Events (20 trials)	Culotte	0.48 (0.34-0.66)	1.12 (0.74-1.71)	1.25 (0.78-1.99)	1.04 (0.78-1.40)	0.94 (0.68-1.29)
	2.10 (1.51-2.92)	DK Crush	2.36 (1.61-3.46)	2.62 (1.62-4.23)	2.19 (1.63-2.95)	1.97 (1.51-2.57)
	0.89 (0.59-1.35)	0.42 (0.29-0.62)	DBS	1.11 (0.67-1.84)	0.93 (0.64-1.35)	0.83 (0.63-1.10)
	0.80 (0.50-1.28)	0.38 (0.24-0.62)	0.90 (0.54-1.50)	T/TAP	0.84 (0.52-1.34)	0.75 (0.49-1.15)
	0.96 (0.72-1.28)	0.46 (0.34-0.61)	1.08 (0.74-1.57)	1.19 (0.75-1.91)	Crush	0.90 (0.70-1.16)
	1.07 (0.78-1.46)	0.51 (0.39-0.66)	1.20 (0.91-1.58)	1.33 (0.87-2.03)	1.11 (0.86-1.44)	Provisional
Myocardial Infarction (23 trials)	Culotte	0.53 (0.31-0.91)	1.06 (0.57-1.96)	0.77 (0.29-2.08)	0.94 (0.58-1.50)	0.84 (0.51-1.39)
	1.89 (1.10-3.24)	DK Crush	2.00 (1.11-3.61)	1.45 (0.54-3.93)	1.76 (0.54-2.74)	1.58 (0.99-2.52)
	0.94 (0.51-1.75)	0.50 (0.28-0.90)	DBS	0.73 (0.28-1.91)	0.88 (0.54-1.45)	0.79 (0.55-1.14)
	1.30 (0.48-3.49)	0.69 (0.25-1.86)	1.37 (0.52-3.60)	T/TAP	1.21 (0.47-3.12)	1.09 (0.44-2.65)
	1.07 (0.67-1.72)	0.57 (0.36-0.88)	1.13 (0.69-1.86)	0.82 (0.32-2.12)	Crush	0.90 (0.64-1.25)
	1.19 (0.72-1.98)	0.63 (0.40-1.01)	1.26 (0.88-1.82)	0.92 (0.38-2.25)	1.12 (0.80-1.56)	Provisional
Stent Thrombosis (22 trials)	Culotte	0.22 (0.08-0.59)	0.64 (0.13-3.15)	0.92 (0.27-3.12)	0.73 (0.39-1.37)	0.37 (0.14-0.95)
	4.56 (1.71-12.19)	DK Crush	2.92 (0.64-13.20)	4.20 (1.44-12.24)	3.35 (1.29-8.73)	1.69 (0.87-3.31)
	1.56 (0.32-7.71)	0.34 (0.08-1.55)	DBS	1.44 (0.29-7.18)	1.15 (0.23-5.70)	0.58 (0.15-2.29)
	1.08 (0.32-3.67)	0.24 (0.08-0.69)	0.69 (0.14-3.46)	T/TAP	0.80 (0.24-2.69)	0.40 (0.17-0.94)
	1.36 (0.73-2.54)	0.30 (0.11-0.78)	0.87 (0.18-4.32)	1.25 (0.37-4.24)	Crush	0.51 (0.20-1.28)
	2.69 (1.05-6.92)	0.59 (0.30-1.16)	1.72 (0.44-6.77)	2.48 (1.06-5.79)	1.98 (0.78-4.99)	Provisional
Target Lesion Revascularization (21 trials)	Culotte	0.42 (0.26-0.68)	1.23 (0.65-2.34)	1.31 (0.77-2.22)	0.95 (0.60-1.51)	0.94 (0.58-1.52)
	2.36 (1.47-3.79)	DK Crush	2.91 (1.67-5.08)	3.08 (1.84-5.15)	2.25 (1.47-3.43)	2.21 (1.54-3.16)
	0.81 (0.43-1.54)	0.34 (0.20-0.60)	DBS	1.06 (0.58-1.92)	0.77 (0.42-1.42)	0.76 (0.49-1.17)

	0.77 (0.45-1.30)	0.32 (0.19-0.54)	0.94 (0.52-1.71)	T/TAP	0.73 (0.42-1.27)	0.72 (0.47-1.09)
	1.05 (0.66-1.66)	0.45 (0.29-0.68)	1.29 (0.71-2.37)	1.37 (0.97-2.38)	Crush	0.98 (0.64-1.52)
	1.07 (0.66-1.73)	0.45 (0.32-0.65)	1.32 (0.86-2.02)	1.40 (0.92-2.11)	1.02 (0.66-1.57)	Provisional
Target Vessel Revascularization (19 trials)	Culotte	0.43 (0.38-0.66)	1.06 (0.57-1.97)	0.97 (0.47-2.00)	0.99 (0.67-1.46)	0.78 (0.47-1.28)
	2.33 (1.51-3.60)	DK Crush	2.47 (1.43-4.28)	2.26 (1.17-4.35)	2.30 (1.56-3.39)	1.81 (1.23-2.68)
	0.94 (0.51-1.76)	0.40 (0.23-0.70)	DBS	0.91 (0.47-1.76)	0.93 (0.52-1.65)	0.73 (0.50-1.09)
	1.03 (0.50-2.13)	0.44 (0.23-0.85)	1.09 (0.57-2.11)	T/TAP	1.02 (0.52-2.01)	0.80 (0.47-1.36)
	1.01 (0.68-1.50)	0.43 (0.30-0.64)	1.07 (0.60-1.91)	0.98 (0.50-1.93)	Crush	0.79 (0.51-1.21)
	1.29 (0.78-2.11)	0.55 (0.37-0.81)	1.36 (0.92-2.02)	1.24 (0.74-1.94)	1.27 (0.83-1.94)	Provisional

Figure S1. Forest plot of two-stent versus provisional stent on all-cause mortality.




Figure S2. Forest plot of two-stent versus provisional stent on cardiac death.




Figure S3. Forest plot of two-stent versus provisional stent on major adverse cardiac events.




Figure S4. Forest plot of two-stent versus provisional stent on myocardial infarction.




Figure S5. Forest plot of two-stent versus provisional stent on stent thrombosis.




Figure S6. Forest plot of two-stent versus provisional stent on target lesion revascularization.




Figure S7. Forest plot of two-stent versus provisional stent on target vessel revascularization.




Figure S8. Comparison of two-stent versus provisional stent stratified to lesion length of side branch on all-cause mortality.

Study or Subgroup	Experimental Events	Experimental Total	Control Events	Control Total	Risk Ratio Random, 95% CI
Group = SB <10mm					
EBC MAIN	10.0	237	7.0	230	1.39 [0.54; 3.58]
NBBS IV	5.0	228	5.0	218	0.96 [0.28; 3.26]
POLBOS II	1.0	102	3.0	100	0.33 [0.03; 3.09]
TRYTON	4.0	355	4.0	349	0.98 [0.25; 3.90]
POLBOS I	2.0	120	3.0	123	0.68 [0.12; 4.02]
NBS	21.0	202	12.0	202	1.75 [0.88; 3.46]
Colombo et al. (2004)	1.0	63	0.0	22	1.08 [0.05; 25.54]
Total (95% CI)		**1308**		**1245**	**1.27 [0.81; 1.97]**
Heterogeneity: $Tau^2 = 0$; $Chi^2 = 3.1$, df = 6 (P = 0.80); $I^2 = 0\%$					
Group = SB ≥10mm					
DEFINITION II	9.0	328	11.0	325	0.81 [0.34; 1.93]
DKCRUSH-V	16.0	240	18.0	242	0.90 [0.47; 1.72]
EBC TWO	1.0	97	2.0	103	0.53 [0.05; 5.76]
BBK I	10.0	101	8.0	101	1.25 [0.51; 3.04]
PERFECT	3.0	213	2.0	206	1.45 [0.24; 8.59]
Total (95% CI)		**979**		**977**	**0.96 [0.62; 1.47]**
Heterogeneity: $Tau^2 = 0$; $Chi^2 = 0.97$, df = 4 (P = 0.91); $I^2 = 0\%$					
Test for subgroup differences: $Chi^2 = 0.80$, df = 1 (P = 0.37)					




Figure S9. Comparison of two-stent versus provisional stent stratified to lesion length of side branch on cardiac death.




Figure S10. Comparison of two-stent versus provisional stent stratified to lesion length of side branch on major adverse cardiac events.

Study or Subgroup	Experimental Events	Experimental Total	Control Events	Control Total	Risk Ratio Random, 95% CI
Group = SB <10mm					
EBC MAIN	42.0	237	34.0	230	1.20 [0.79; 1.81]
NBBS IV	5.0	228	12.0	218	0.40 [0.14; 1.11]
POLBOS II	12.0	102	15.0	100	0.78 [0.39; 1.59]
SMART-STRATEGY	8.0	130	1.0	128	7.88 [1.00; 62.08]
TRYTON	65.0	355	45.0	349	1.42 [1.00; 2.02]
POLBOS I	16.0	120	15.0	123	1.09 [0.57; 2.11]
NBS	44.0	202	32.0	202	1.38 [0.91; 2.07]
CACTUS	28.0	177	26.0	173	1.05 [0.64; 1.72]
Total (95% CI)		**1551**		**1523**	**1.20 [1.00; 1.44]**
Heterogeneity: Tau^2 = < 0.0001; Chi^2 = 10.67, df = 7 (P = 0.15); I^2 = 34%					
Group = SB ≥10mm					
DEFINITION II	20.0	328	37.0	325	0.54 [0.32; 0.90]
DKCRUSH-V	20.0	240	41.0	242	0.49 [0.30; 0.81]
DKCRUSH-II	29.0	183	44.0	183	0.66 [0.43; 1.00]
EBC TWO	10.0	97	8.0	103	1.33 [0.55; 3.22]
Zhang et al. (2016)	4.0	52	2.0	52	2.00 [0.38; 10.45]
BBK I	23.0	101	23.0	101	1.00 [0.60; 1.66]
PERFECT	38.0	213	38.0	206	0.97 [0.64; 1.45]
Ye et al. (2012)	1.0	38	5.0	30	0.16 [0.02; 1.28]
Lin et al. (2010)	6.0	54	21.0	54	0.29 [0.13; 0.65]
Ye et al. (2010)	0.0	25	1.0	26	0.35 [0.01; 8.12]
Total (95% CI)		**1332**		**1323**	**0.68 [0.50; 0.93]**
Heterogeneity: Tau^2 = 0.0966; Chi^2 = 17.5, df = 9 (P = 0.04); I^2 = 49%					
Test for subgroup differences: Chi^2 = 9.64, df = 1 (P < 0.01)					




Figure S11. Comparison of two-stent versus provisional stent stratified to lesion length of side branch on myocardial infarction.




Figure S12. Comparison of two-stent versus provisional stent stratified to lesion length of side branch on stent thrombosis.




Figure S13. Comparison of two-stent versus provisional stent stratified to lesion length of side branch on target lesion revascularization.




Figure S14. Comparison of two-stent versus provisional stent stratified to lesion length of side branch on target vessel revascularization.





Figure S15. Bar graph showing SUCRA scores of each bifurcation technique for every outcome.



	All-cause Mortality	Cardiac Death	Major Adverse Cardiac Events	Myocardial Infarction	Stent Thrombosis	Target Lesion Revascularization	Target Vessel Revascularization
Provisional	0.5826	0.4923	0.6759	0.5278	0.6474	0.6271	0.5822
Culotte	0.3111	0.3676	0.5046	0.4266	0.3854	0.5055	0.573
DK Crush	0.7048	0.8292	0.9998	0.9324	0.975	0.9999	0.9987
DBS	0.6655	0.8639	0.3968	0.41019	0.3596	0.2486	0.1864
T/TAP	0.5119	0.1853	0.1721	0.5663	0.1264	0.1576	0.3228
Crush	0.2242	0.2627	0.2508	0.145	0.5061	0.4613	0.3369

Figure S16. Bayesian network meta-analysis.

