Supplementary Table 4. GRADE of Each Comparison

Certainty assessment							Summary of findings				
							Study event rates (%)			Anticipated absolute effects	
Participant (studies) follow-up	Risk of bias	Inconsistency	Indirectness	Imprecision	Publication bias	Overall certainty of evidence	With before the COVID-19 pandemic	With during the COVID-19 pandemic	Relative effect (95% CI)	Risk with before the COVID-19 pandemic	Risk difference with during the COVID-19 pandemic
OHCA at home											
2251 (2 observational studies)	Not serious	Not serious	Not serious	Not serious	None	⨁⨁◯◯ Low	704/1036 (68.0)	888/1215 (73.1)	OR 1.29 (1.08 to 1.55)	680 per 1,000	53 more per 1,000 (from 17 more to 87 more)
Bystander CPR											
3598 (5 observational studies)	Not serious	Serious	Not serious	Not serious	None	⨁◯◯◯ Very low	1152/1951 (59.0)	966/1647 (58.7)	OR 1.09 (0.81 to 1.47)	590 per 1,000	21 more per 1,000 (from 52 fewer to 89 more)
Unwitnessed arrest											
3598 (5 observational studies)	Not serious	Serious	Not serious	Not serious	None	⨁◯◯◯ Very low	934/1951 (47.9)	823/1647 (50.0)	OR 0.83 (0.59 to 1.16)	479 per 1,000	46 fewer per 1,000 (from 127 fewer to 37 more)
AED usage											
2580 (3 observational studies)	Not serious	Not serious	Not serious	Not serious	None	⨁⨁◯◯ Low	129/1194 (10.8)	115/1386 (8.3)	OR 0.74 (0.57 to 0.97)	108 per 1,000	26 fewer per 1,000 (from 43 fewer to 3 fewer)
Shockable cardiac rhythm											
3598 (5 observational studies)	Not serious	Serious	Not serious	Not serious	None	⨁◯◯◯ Very low	373/1951 (19.1)	250/1647 (15.2)	OR 1.17 (0.87 to 1.59)	191 per 1,000	25 more per 1,000 (from 21 fewer to 82 more)
Intubation											
2283 (2 observational studies)	Not serious	Serious	Not serious	Not serious	None	⨁◯◯◯ Very low	70/1049 (6.7)	29/1234 (2.4)	OR 0.36 (0.14 to 0.97)	67 per 1,000	42 fewer per 1,000 (from 57 fewer to 2 fewer)
Supraglottic airway											
2283 (2 observational studies)	Not serious	Very serious	Not serious	Not serious	None	⨁◯◯◯ Very low	725/1049 (69.1)	1011/1234 (81.9)	OR 1.22 (0.16 to 9.26)	691 per 1,000	41 more per 1,000 (from 427 fewer to 263 more)
ROSC											
3840 (5 observational studies)	Not serious	Serious	Not serious	Not serious	None	⨁◯◯◯ Very low	430/1815 (23.7)	447/2025 (22.1)	OR 0.90 (0.71 to 1.13)	237 per 1,000	19 fewer per 1,000 (from 56 fewer to 23 more)
Survival to discharge											
3368 (4 observational studies)	Not serious	Serious	Not serious	Not serious	None	⨁◯◯◯ Very low	231/1822 (12.7)	116/1546 (7.5)	OR 0.74 (0.43 to 1.26)	127 per 1,000	30 fewer per 1,000 (from 68 fewer to 28 more)
Ravorable neurological outcome											
3598 (5 observational studies)	Not serious	Serious	Not serious	Not serious	None	⨁◯◯◯ Very low	189/1951 (9.7)	91/1647 (5.5)	OR 0.77 (0.43 to 1.37)	97 per 1,000	21 fewer per 1,000 (from 53 fewer to 31 more)

CI, confidence interval; OR, odds ratio.