Table S1. Core-based group and control group in the included parallel group trials.

Author, Year	Core-based Group	Control Group	Duration
Schreiber, 2015[23] and Schreiber, 2016[24]	***Schroth exercises:*** with the corrective movements, the targeted curve type, the level of passive support involved. ***Individual exercises:*** 60 min/session during the first 2 weeks ***Group exercises:*** 60 min/session ***Frequency:*** 1 time/week ***Home exercises:*** 30-45min/day ***Frequency:*** 1 time/week	***Observation or bracing:*** in accordance with SRS recommendations	24 weeks
Gür, 2017[25]	***Core stabilization exercises:*** local and global muscle stability training, global muscle mobility training, strength training of the core muscles; ***Traditional treatment:*** breathing exercises, posture training, spinal flexibility exercises, general strengthening exercises	***Traditional treatment:*** breathing exercises; posture training; spinal flexibility exercises; general strengthening exercises ***Exercise:*** 2h of removal for hygienic activities and exercise ***Bracing:*** 22h/day	10 weeks
Ko, 2017[26]	***Warm-up exercises:*** 10 min stretching and breathing ***Core stabilization exercises:*** 40 min; 12 repetitions/3 sets pelvic tilt, cat-camel pose (back raises), basic trunk curl (crunch), bridge (with knee extension), double-leg abdominal press, supermans, arm/leg raises, quadruped arm/leg raises, and hand walkouts ***Cool-down exercises:*** 10 min stretching and breathing ***Frequency:*** 3 times /week	***Observation***	12 weeks
Yongnam, 2014[27]	***Warm-up exercises:*** 10 min walking and stretching of various body parts ***Core stabilization exercises:*** 40 min, 10 repetitions Stretching exercise used a swiss ball or other adequate equipment Strengthening trunk muscle on the convex side ***Cool-down exercises:*** 10 min walking and stretching of various body parts ***Frequency:*** 3 times/week	***Observation***	8 weeks
Cheon, 2013[28]	***Warm-up exercises:*** 5 min stationary bicycle	***Chiropractic:*** 10 min; cervical spine lateral correction (left and right);	12 weeks

	Stretching: 10 min Static/dynamic stretching (iliopsoas m/piriformis m /hamstring m/trunk twist) ***Lumbar exercise program:*** 20 min; 5 repetitions (10 s hold)/3 sets dumbbell side bending/thoracic lateral flexion/trunk extension exercise scapula/lumbar stabilizing exercise ***Cool-down exercises:*** 5 min stationary bicycle ***Frequency:*** 3 times/week ***Chiropractic:*** 10 min cervical spine lateral correction (left and right) thoracic spine rotation correction in sitting lumbar spine rotation correction in supine pelvic, prone, and posteroinferior correction ***Frequency:*** 3times/week	thoracic spine rotation correction in sitting; lumbar spine rotation correction in supine pelvic, prone, and posteroinferior correction ***Frequency:*** 3 times/week	
Kuru, 2016[29]	***Schroth exercises:*** spinal elongation; de-rotation/de-flexion; stretching; strengthening and rotational breathing exercises ***Supervised exercises:*** 90 min/day ***Frequency:*** 3 times/week ***Home exercises:*** 90 min/day ***Frequency:*** 3 times/week	***Observation***	6 weeks
Zheng, 2018[30]	**SEAS:** spinal stabilization, strengthening, muscular retraction, motor coordination ***Learning exercises:*** 90 min/session ***Frequency:*** 1 time/week ***Supervised exercises:*** 40 min/session ***Frequency:*** 1 time/week ***Home exercises:*** 10-15 min/session ***Frequency:*** 1 time/day	***Bracing:*** 23h/day rigid thoracolumbosacral orthosis	48 weeks
Gao C, 2019[31]	**SEAS:** active self-correction, spinal stabilization, breathing ***Learning exercises frequency:*** 1 time/2-3 months ***Supervised exercises:*** 40 min/session ***Frequency:*** 1 time/week	***Bracing:*** 23h/day rigid thoracolumbosacral orthosis ***Re-evaluation frequency:*** 1 time/3 months	24 weeks

	Home exercises: 10-15 min/session ***Frequency:*** 1 time/day ***Bracing:*** 23h/day rigid thoracolumbosacral orthosis ***Re-evaluation frequency:*** 1 time/3 months		
Alves, 2012[32]	***Pilates method:*** ***Warm-up:*** 8 min of walking followed by stretching for 5 min ***Specific exercises:*** Pilates technique with swiss balls ***Cool-down:*** 3 movements, with 3 repetitions per movement, and lasted 5 min ***Supervised exercises:*** 60 min/day ***Frequency:*** 2 times/week	***Observation***	12 weeks

SEAS: scientific exercises approach to scoliosis.

Table S2. The Cochrane tool of assessing risk of bias for methodological assessment (RoB 2.0 tool).

Article, Year	Randomization process	Deviations From Intended Interventions	Missing Outcome Data	Measurement of the Outcome	Selection of the reported	Overall
Schreiber, 2015[23], Schreiber, 2016[24]	Low	Low	Low	Low	Low	Low
Gür, 2017[25]	Low	Low	Low	Unclear	Low	Unclear
Yongnam, 2014[27]	Unclear	Low	Low	Unclear	Low	Unclear
Kuru, 2015[29]	Low	Low	Low	Unclear	Low	Unclear
Zheng, 2018[30]	Low	Low	Low	Unclear	Low	Unclear
Gao C, 2019[31]	Low	Low	Low	Low	Low	Low
Alves, 2012[32]	Low	Low	Low	High	Low	High