Detail searching strategy was as follows.

PubMed:

(‘‘extracorporeal membrane oxygenation’’ [Mesh] OR ECMO[Title/Abstract] OR extracorporeal cardiopulmonary resuscitation [Title/Abstract] OR ECPR[Title/Abstract] OR extracorporeal life support [Title/Abstract] OR ECSL[Title/Abstract]) AND (‘‘seizure’’[Mesh] OR “status epilepticus” [Mesh] OR epileptic state [Title/Abstract] OR neurologic outcome [Title/Abstract] OR neurologic prognosis [Title/Abstract] OR neurological complication [Title/Abstract] OR neurological injury [Title/Abstract])

Search results: 190 articles.

Embase:

('extracorporeal membrane oxygenation'/exp OR 'extracorporeal membrane oxygenation' OR 'extracorporeal cardiopulmonary resuscitation' 'ECPR' OR 'extracorporeal life support' OR 'ECSL') AND ('seizure'/exp OR 'seizure' OR 'epileptic state' OR 'neurologic outcome' OR 'neurologic prognosis' OR 'neurological complication' OR 'neurological injury')

Search results: 352 articles.

Web of Science:

(Ts=(extracorporeal membrane oxygenation) OR TI=(extracorporeal membrane oxygenation OR ECMO OR extracorporeal cardiopulmonary resuscitation OR ECPR OR extracorporeal life support OR ECSL)) AND ((Ts=seizure)

OR TI=(seizure OR epileptic state OR neurologic outcome OR neurologic prognosis OR neurological complication OR neurological injury))

Search results: 226 articles.




Figure S1 Subgroup analysis for the rate of seizure in neonatal and pediatric patients. The incidence of seizures was calculated with a random-effects model. Abbreviations: CI, confident interval; ES, estimate.