Author	year	total	male	Female	Age	follow-up	rec	Htransf	Ctransf	IDHmt	codel	surgery (T//ST//partial or biopsy	aduvant radiio-therapy	chemo-therapy	Astro	Olig	mixed	comment	institue and year
Ahmadi	**2009**	130	79	51	M36.9	M94mo(24-264)	128	43				91//?//39	36	14	60	34	36	total removal, >=90% removal	1985-2003 Heiderberg University
Aoki	**2021**	276	178	98	m39.7, M37	m5.87 M5.1			107	153+123	123	35/34/207	55	109	153	123	0	multi-center	1990-2008 Nagoya University
Bauman	**2009**	145	92	53	m39	M105mo	110	43			in part	71/43/31	71	0	74	51	20		1979-1995University of Western Ontario, Canada
Cappelle	**2013**	929	522	407	M44(18-76)	M6.3y,m7.4y	539		382			235, >=90%removal	258	209	NA				1985-2007French multicenter study group
Cesselli	**2020**	241	144	97	M39(19-75)	M66mo(13-239)	185		152	213	77	99/97/45	0	0	164	77		no initial adjuvant therapy, biopsy were excluded	2000-2016Hospital of Udine. Italy
Chaichana	**2010**	191	111	80	m34.9/15.3	m36mo SD	83		44			69//122	16	0	89	89	13	biopsy cases were excluded	1996-2006 Johns Hopkins School of Medicine
Esseonu	**2017**	109	63	46	M37(19-74)	M5.2y	28		20	assessed in 51 patien		59, >=90% removal	34	16	73	36	0	chemotherapy for patients with subtotally removed tumors	2006- 2015 Johns Hopkins School of Medicine
Dahlrot	**2013**	66	42	24	M48	M36mo(0.4-79)	38	13		30/52	9//16	16/17/26・None7	NA	NA		39	20		Odense Univ. Hosp Denmark
El-Hateer	**2009**	69	44	25	M39(20-78)	M6.1y(1.3-16.3)	47	9			38/68	19/41/9	15	6	0	69			1992-2006 McGill Univ Health Center
Etxaniz	**2017**	58	33	25	m45.9(12-77)	M7.6y	42	8		26/42	25/54	24//11/24	30	6	23	13	22	CHT tmz	1990-2012 Hospital Univ. Germans Trias I Pujol, Spain

Fukuya	**2019**	227				M6.7y (1.0-14.8)	81		38	64/81	28/81	Mvol90%	23	NA	NA	NA	NA	biopsy cases were excluded	2000-2013Tokyo Women Med Univ
Gousias	**2014**	148	83	65	M38(18 -74.1)	m69mo SD59(1-196)	81	59				62/75/11	8	2	76	54	18		1996-2011University Hospital of Bonn,
Heo	**2017**	49	31	18	m43.8S D10.9	m73.0mo SD14.3	41	15				26//	25	NA		49	0	gemistocytic astrocytomas were excluded	2005-2015Busan Park Hosp. Korea
Iwadate	**2015**	94	57	37	m41(22 -84)	M8.5y		20		78(im muno) /89	38/69	46/27/21	0	NA		38	47	9	oligo with residual tumor,PCV	1996-2013 Chiba Univ
Jaber	**2019**	74	42	32	m42.8S D13.4	M46.4mo CI(41.8-51.1)	41	17		46+12	12	24/50	24	9	62	12		total removal, >=90% removal	2010-2016 Univ. Hospital Mu¨nster,
Jakola A	**2012/ 2017**	66	41	25	m45 SD15	M7y(4.5-10.9)			37	48/64	23/64te	19//47	19	13	55	6	5	biopsy preffered	1998-2009St Olavs Univ. &Norweigian Univ
Jakola B	**2012/ 2017**	87	47	40	m48 SD56	M7.1y(4.2 -9.9)			31	56/81	20/81te	75//12	37	18	62	16	9	resection preffered	1998-2009St Olavs Univ. &Norweigian Univ
Jansen E	**2019**	110	57	53	M37	M126mo 95%CI(109 -143)	93	72		53+39	39	58/28/24	0	0	71	39		no iniitial adjuvant therapy	1985-2018Univ Cologne and Freiburg
Jansen NL	2014	59	32	27	m42.6S D11.3	M47mo	29	20	21			7//52	14	15	59	0	0	GII astro with 18F-FETPET	2005-2011 University of Munich,
Jung	2011	86	47	39	M41(SD 17.4)	m4.25y SD2.8	36	13				T40/22/24	27	7PCV		32	42	12	LGG including 10 adol	2000-09 Chonnam National University, Korea
Jungk	**2016**	46	21	25	M35 (17-54)	M69mo (17.5-164.6)	26		19	38(IDH 1)	0	21, ＞ 90％ removal	3	2	46	0		astrocytoma only	Portland Veterans Medical Center, Oregon

Jurati	**2012**	71			19-72	M9.6y (3.6-19)	60	45		54		24/37/10		22	0	55	0	16	astrocytoma only	93-2007Dresden Univ Hosp
Karschnia	2020	155	76	79	codel 43 ± 1.4 years	M35mo(3-136)	69	34		51+81	81	21//14//12		41	67	74	81		40 patients with wait and see after diagnosis	2005-2019 Ludwig Maximilians University School of Medicine
Kavouridis	**2020**	326	178	148	M36(IQR30-46)	M5.4y (IQR2.6-9.5)	176	80		astro154	140	157/156/0		83	43+63	154+ wild32	140			2000-2017 Brigham and Women's Hospital
Koo	**2018**	555	292	263	M40(30-77)	M83.4y (6-184.7)	301	114				200/224/131		269	20+65	210	85	260	>=90% removal, 353	00-2010Korea multi-institution
Leu	**2016**	210	126	84	m41.9	m6.87y		69		150/185	39/160	NA	NA		57/162	92	53	63		University Hospital Basel,
Majchrzak	**2012**	68	39	29	M36	M34mo (IQR21-49)	22	11				21/13/34		50	NA	46	5	17		2005-2011 Medical Univ.of Silesia, Sosnowiec, Poland
Obara	**2020**	339	188	151	M38(18-80)	m9.6y (M8.5)			158	139/175	64/182	48/147/42		40	106	86	199	54	delayed treatment, 56 cases, excluded pts with death unrelated to the	1982-2017 Nancy France University Hospital
Patel	**2018**	74	36	38	M43(20-71)	M44.4mo (0.6-187.2)		23	31	52	NA	NA(%resect)/biopsy 8		41	11+37	43	19	12		2001-2012 Univ. of Texas Southwestern Medical center Dallas
Ross	**2014**	89	51	38	m40.3SD13.8	m93.2mo	47		30	9//13	29/49te	NA	NA		NA	0	89		oligodendroglioma or	Portland Veterans Medical Center, Oregon
Sakarunch	**2013**	77	48	29	M43(18-74)	m40mo (3-300)	20	12	15			4/34/39		63	3	55	19	3		00-2009Prince of Songkla Univ. Thailand
Smith	**2008**	216	134	82	M38(19-72)	M52.8mo	95	32	44			101, >=90% removal		45	51	93	91	32	biopsy excluded	1985-2005 UCSF

Snyder	**2014**	93	58	35	NA	M75.4mo	31	20		39/65te	>90% removal,45%	NA	NA		0	93		oligodendroglioma or	1999-2010Barrow Brain Tumor Research Center,
Suzuki	**2018**	102	67	35	M40	M2.74y m2.49y	16	11	81	56	Mvol90%	24	NA		46	56		biopsy cases excluded	2008-2013Tokyo Women Med Univ
Tanaka	**2015**	55	38	17	M46(16-76)	M52mo		5	35/48(immuno)	10//20	21/18/16	36	22		28	21	6	>=95%, total removal	1983-2013 Kobe University hospital
Thon	**2012**	127	60	67	M37(18-75)	M173mo (36-306)	106		80	89/114(IDH1)	NA	T58 biopsy21	18	0	127	0	0	astrocytoma only	1991-1998 Ludwig Maximilians University Munich,
Tom	**2019**	486	262	224	M39(18-78)	M5.3y (0.02-28.4)		56	84	125+169	169	204/104/160	144	82+81		317	169	0	1980-2018 Cleveland Clinic
Turkoglu	**2013**	63	36	27	M38	M4.9y (1.1-11)		13				35/19/9	22	17	28	27	8		1999-2011 Diskapi Yildirim Beyazit Education and Research Hospital
Youland	**2013**	554	324	230	m39.5(18.6-76.0)	M5.2y	330	103			133	171/168/215	322	100	120	247	187		1992-2011Mayo clinic
Zhang	**2019**	68	31	37	35.3(SD 15.1, n=15) and 40.3(SD	m4.68y		15		33/41	25/30te	NA	26	27	32	36	0	for Htrans; at least 2 years follow-up	2000-2017 Tongii Hospital Wuhan China
Zeng	**2021**	75	33	42	M38 (21-53)	m7.2			27	44+31	31	59/13/3	22	22	44	31		for Ctrans: icluding 49 incidental cases	2008-2015 Tongii Hospital, Wuhan China
For forest plots only																			
Goze	**2014**	131	79	52	m38SD10.8M38	m70SD54.7M55		20	56	107	38/11	18//4//64	23	14	25	71	35	(excluding gemistcytic; overlapping in Capella et al.	92-2012 French glioma study group

Murphy	**2017**	599	327	272	M36	M7.36Y		76	48		147/4	209//17 2//203	91+47	88+47	240	255	104	overlapping in Tom et al.	1988-2014Cleveland Clinic
Pallude	**2013**	407	227	180	m39S D11	m86.5SD 59.4M73		109	209		67/20	38//64//278			76	246	85	overlapping in Capella et al.	92-2011 French glioma study group
Wellar	**2021**	142	68	74	M42(2 0-80)	m6.5M5.9		20		142	142	10/17/115		56				overlapping in Karschnia et al. oligodendroglioma only	03-2019 Univ Hosp Munich
Wenn	**2018**	168	74	94	M45(1 8-61)	NA	42		29	IDH1, 113	NA	115//	91	108TM	168	0		astrocytoma only; follow-up period was not availble	09-2017 Central Hospital of Yiyang, China

Htrans, histologically provrn malignant transformation

Ctrans, clinically defined malignant transformation

CE, contrast enhancement on MRI

T//ST//Partial bipopsy, number of total removal, sbtotal removal and partialremoval or biopsy

M, median; m, mean; SD, standarddeviation