Supplementary Table 1. Type of scope, sedation, and technical details of endoscopic drainage

Study	Scope used	Sedation	Technique
Sharma and Maharshi, 2008 [15]	Olympus IT-10 and TJF-100 endoscopes (Olympus, Tokyo, Japan)	Ketamine and diazepam	Needle knife sphincterotome for puncture, guidewire, 5 to 6 mm dilatation with a standard papillotome
Jazrawi et al., 2011 [16]	TJF160; GFUC140-P; GFUCT140 (Olympus America Corp., Melville, NY)	GA	Cyst wall puncture with 19G or 22G Echotip needle (Cook Medical, Winston-Salem, NC) under EUS guidance, 0.035-inch standard guidewire, Huibregtse triple lumen needle knife (Cook Medical, Winston-Salem, NC) for dilatation of tract, 6- to 10-mm balloon dilatation (Hurricane Rx; Microvasive Endoscopy, Boston Sci Corp., Natick, MA or Quantum TTC; Cook Medical, Winston-Salem, NC)
Makin et al., 2012 [17]	Pediatric 11-mm side-viewing endoscope (Olympus, JF 240, Olympus UK Ltd)	GA	Puncture using endoscopic cystotome CST10 (Cook Medical, Bloomington, IN), followed by a guidewire and stent placement, no dilatation of tract
Ramesh et al., 2013 [20]	Curvilinear array echoendoscope (GF-UCT 140; Olympus America, Melville, NY)	GA	19-gauge FNA needle (Echotip, Cook Medical Inc, Bloomington, IN), 0.035-in. guidewire, dilatation by 4.5F ERCP canula (Proforma HF, Conmed Technologies, Utica, NY) and 8- to 15-mm balloon dilator
Agarwal et al., 2013 [18]	Curvilinear GF-UCT140 echoendoscopes (Olympus, Tokyo, Japan)	Ketamine with midazolam f/b propofol	NR
Bai, 2013 [19]	NR	NR	NR
Bang and Varadarajulu, 2016 [21]	GF UCT180T (Olympus America Inc, Center Valley, PA)	GA	19G FNA needle (Expect needle, Boston Scientific Corporation, Natick, MA), 0.025-in. guide wire, dilatation using a 4.5Fr cannula and a 12- to 15-mm balloon (CRE balloon, Boston Scientific Corporation, Natick, MA), stent placement
Nabi et al., 2017 [22]	Adult linear EUS scope (Olympus 180, Japan)	Ketamine with midazolam f/b propofol	19G FNA needle, guidewire with flexible tip (0.025"; 450 cm), 6 Fr cystotome (Endoflex, Voerde, Germany), balloon dilatation (Hurricane, 4 mm, Boston Scientific, Natick, MA Boston Scientific), stent placement
Nabi et al., 2019 [23]	NR	Ketamine with midazolam f/b propofol	NR
Farr et al., 2020 [24]	NR	NR	NR
Lal et al., 2020 [25]	NR	NR	NR
Poddar et al., 2021 [27]	Duodenoscope (Olympus, Japan, TJF180V)	Moderate (midazolam/ ketamine)	Cyst wall puncture with needle knife (5.5 F, triple lumen, micro-knife, Boston Scientific Corp. USA), guide wire coiling (0.35 Inches, 480 cm, Tracer Metro, Wilson Cook, USA), dilatation with 8 mm biliary balloon (Quantum TTC Biliary Balloon Dilator [QBD-8X3], Wilson Cook, USA), stent placement
Seol et al., 2021 [28]	NR	NR	NR
Ghoneem et al., 2021 [26]	NR	NR	NR

NR: not reported, GA: general anesthesia, f/b: followed by, FNA: fine-needle aspiration.