Supplementary Table 2. Recurrence and re-intervention rates after endoscopic drainage of pancreatic fluid collection

Study	ERP or MRCP	PD anatomy	Stent removal	Stent migration
Sharma and Maharshi, 2008 [15]	2	DPD: 2	6 (1 m: 4, 2 m: 2)	3
Jazrawi et al., 2011 [16]	NR	NR	NR	NR
Makin et al., 2012 [17]	3 (PD stenting)	NR	8 w (6-40)	1
Ramesh et al., 2013 [20]	ERP 4, Unsuccessful 3	PD stent 2, DPD 2	NR	0
Agarwal et al., 2013 [18]	NR	NR	NR	NR
Bai, 2013 [19]	NR	NR	NR	NR
Bang and Varadarajulu, 2016 [21]	ERP all	DPD 1, normal 5	8 w	0
Nabi et al., 2017 [22]	Yes	Normal=12, DPD 9, leak 2, divisum 2, stricture 1	N=13; 8-12 w=8, 12-16 w=3, >16 w=2	4
Nabi et al., 2019 [23]	Yes	DPD 25, normal 3, stricture 1, CP 1	4 w	NR
Farr et al., 2020 [24]	NR	NR	NR	NR
Lal et al., 2020 [25]	NR	NR	NR	NR
Poddar et al., 2021 [27]	Not done	NR	12 w (7-20) n=26	3
Seol et al., 2021 [28]	NR	NR	NR	1
Ghoneem et al., 2021 [26]	NR	NR	NR	NR

ERP: endoscopic retrograde cholangiopancreatography, MRCP: magnetic resonance cholangiopancreatography, PD: pancreatic duct, NR: not reported, DPD: disconnected pancreatic duct, CP: chronic pancreatitis, m: month, w: week.