

**A**


anti-EGFR Control Hazard Ratio Hazard Ratio
Study or Subgroup log[Hazard Ratio] SE Total Total Weight IV, Random, 95% CI IV, Random, 95% CI
2.1.1 EGFR+CT
Lorenzen 2009 -0.2236 0.2821 32 30 3.9% 0.80 [0.46, 1.39]
Lu 2017 -0.4734 0.226 40 40 5.5% 0.62 [0.40, 0.97]
Subtotal (95% CI) 72 70 9.4% 0.69 [0.49, 0.97]
Heterogeneity: Tau² = 0.00; Chi² = 0.48, df = 1 (P = 0.49); I² = 0%
Test for overall effect: Z = 2.13 (P = 0.03)
2.1.2 EGFR+CRT
Castro 2018 -0.3275 0.1968 53 54 6.6% 0.72 [0.49, 1.06]
Chen 2016a -0.7186 0.199 68 68 6.5% 0.49 [0.33, 0.72]
Chen 2016b -1.4092 1.0701 32 32 0.3% 0.24 [0.03, 1.99]
Crosby 2017 0.2103 0.1389 129 129 9.8% 1.23 [0.94, 1.62]
Dutton 2014 -0.1036 0.0987 224 225 12.8% 0.90 [0.74, 1.09]
Luo 2015 -0.2521 0.3389 28 27 2.9% 0.78 [0.40, 1.51]
Sun 2014 -0.4586 0.2733 21 21 4.1% 0.63 [0.37, 1.08]
Suntharalingam 2017 -0.09 0.1217 159 169 11.0% 0.91 [0.72, 1.16]
Wu 2018 -0.239 0.122 176 176 11.0% 0.79 [0.62, 1.00]
Ye 2015 -0.4027 0.2494 37 36 4.8% 0.67 [0.41, 1.09]
Subtotal (95% CI) 927 937 69.8% 0.80 [0.67, 0.95]
Heterogeneity: Tau² = 0.04; Chi² = 20.33, df = 9 (P = 0.02); I² = 56%
Test for overall effect: Z = 2.60 (P = 0.009)
2.1.3 EGFR+IC+CRT+S
Ruhstaller 2018 -0.2989 0.1525 149 151 8.9% 0.74 [0.55, 1.00]
Subtotal (95% CI) 149 151 8.9% 0.74 [0.55, 1.00]
Heterogeneity: Not applicable
Test for overall effect: Z = 1.96 (P = 0.05)
2.1.5 EGFR+RT
Shi 2014 -0.2868 0.2176 32 32 5.8% 0.75 [0.49, 1.15]
Zhou 2015 -0.3845 0.2112 40 40 6.0% 0.68 [0.45, 1.03]
Subtotal (95% CI) 72 72 11.8% 0.71 [0.53, 0.96]
Heterogeneity: Tau² = 0.00; Chi² = 0.10, df = 1 (P = 0.75); I² = 0%
Test for overall effect: Z = 2.22 (P = 0.03)
Total (95% CI) 1220 1230 100.0% 0.78 [0.69, 0.88]
Heterogeneity: Tau² = 0.02; Chi² = 23.44, df = 14 (P = 0.05); I² = 40%
Test for overall effect: Z = 3.95 (P < 0.0001)
Test for subgroup differences: Chi² = 0.84, df = 3 (P = 0.84), I² = 0%
0.05 0.2 1 5 20
Favours [anti-EGFR] Favours [Control]

**B**


anti-EGFR Control Hazard Ratio Hazard Ratio
Study or Subgroup log[Hazard Ratio] SE Total Total Weight IV, Random, 95% CI IV, Random, 95% CI
2.2.1 mAb
Castro 2018 -0.3275 0.1968 53 54 6.6% 0.72 [0.49, 1.06]
Chen 2016b -1.4092 1.0701 32 32 0.3% 0.24 [0.03, 1.99]
Crosby 2017 0.2103 0.1389 129 129 9.8% 1.23 [0.94, 1.62]
Lorenzen 2009 -0.2236 0.2821 32 30 3.9% 0.80 [0.46, 1.39]
Lu 2017 -0.4734 0.226 40 40 5.5% 0.62 [0.40, 0.97]
Luo 2015 -0.2521 0.3389 28 27 2.9% 0.78 [0.40, 1.51]
Ruhstaller 2018 -0.2989 0.1525 149 151 8.9% 0.74 [0.55, 1.00]
Shi 2014 -0.2868 0.2176 32 32 5.8% 0.75 [0.49, 1.15]
Sun 2014 -0.4586 0.2733 21 21 4.1% 0.63 [0.37, 1.08]
Suntharalingam 2017 -0.09 0.1217 159 169 11.0% 0.91 [0.72, 1.16]
Subtotal (95% CI) 675 685 58.9% 0.81 [0.69, 0.96]
Heterogeneity: Tau² = 0.02; Chi² = 13.92, df = 9 (P = 0.13); I² = 35%
Test for overall effect: Z = 2.47 (P = 0.01)
2.2.2 TKI
Chen 2016a -0.7186 0.199 68 68 6.5% 0.49 [0.33, 0.72]
Dutton 2014 -0.1036 0.0987 224 225 12.8% 0.90 [0.74, 1.09]
Wu 2018 -0.239 0.122 176 176 11.0% 0.79 [0.62, 1.00]
Ye 2015 -0.4027 0.2494 37 36 4.8% 0.67 [0.41, 1.09]
Zhou 2015 -0.3845 0.2112 40 40 6.0% 0.68 [0.45, 1.03]
Subtotal (95% CI) 545 545 41.1% 0.73 [0.59, 0.89]
Heterogeneity: Tau² = 0.03; Chi² = 8.52, df = 4 (P = 0.07); I² = 53%
Test for overall effect: Z = 3.04 (P = 0.002)
Total (95% CI) 1220 1230 100.0% 0.78 [0.69, 0.88]
Heterogeneity: Tau² = 0.02; Chi² = 23.44, df = 14 (P = 0.05); I² = 40%
Test for overall effect: Z = 3.95 (P < 0.0001)
Test for subgroup differences: Chi² = 0.75, df = 1 (P = 0.39), I² = 0%
0.05 0.2 1 5 20
Favours [anti-EGFR] Favours [Control]

**C**





**Figure S1** Subgroup analysis of overall survival (OS) according to different treatment regimens: (A) different treatment modalities; (B) different anti-EGFR groups; (C) different anti-EGFR agents