

A


anti-EGFR
Control
Risk Ratio
Risk Ratio
Study or Subgroup Events Total Events Total Weight M-H, Fixed, 95% CI
M-H, Fixed, 95% CI
7.1.1 CRT
Castro 2018 0 53 0 54 Not estimable
Crosby 2013 12 129 8 129 32.7% 1.50 [0.63, 3.55]
Dutton 2014 13 224 2 225 8.1% 6.53 [1.49, 28.60]
Ramos-Suzarte 2012 6 33 2 30 8.6% 2.73 [0.60, 12.49]
Suntharalingam 2017 11 153 4 166 15.7% 2.98 [0.97, 9.17]
Wu 2018 8 158 8 156 32.9% 0.99 [0.38, 2.56]
Subtotal (95% CI) 750 760 97.9% 2.09 [1.30, 3.36]
Total events 50 24
Heterogeneity: Chi² = 5.73, df = 4 (P = 0.22); I² = 30%
Test for overall effect: Z = 3.05 (P = 0.002)
7.1.2 CT
Lorenzen 2009 5 32 0 30 2.1% 10.33 [0.60, 179.22]
Subtotal (95% CI) 32 30 2.1% 10.33 [0.60, 179.22]
Total events 5 0
Heterogeneity: Not applicable
Test for overall effect: Z = 1.60 (P = 0.11)
Total (95% CI) 782 790 100.0% 2.26 [1.42, 3.60]
Total events 55 24
Heterogeneity: Chi² = 7.13, df = 5 (P = 0.21); I² = 30%
Test for overall effect: Z = 3.46 (P = 0.0005)
Test for subgroup differences: Chi² = 1.17, df = 1 (P = 0.28), I² = 14.7%
0.005 0.1 1 10 200
Favours [anti-EGFR] Favours [Control]

B


anti-EGFR
Control
Risk Ratio
Risk Ratio
Study or Subgroup Events Total Events Total Weight M-H, Random, 95% CI
M-H, Random, 95% CI
7.2.1 mAb
Castro 2018 0 53 0 54 Not estimable
Crosby 2013 12 129 8 129 26.8% 1.50 [0.63, 3.55]
Lorenzen 2009 5 32 0 30 4.2% 10.33 [0.60, 179.22]
Ramos-Suzarte 2012 6 33 2 30 12.5% 2.73 [0.60, 12.49]
Suntharalingam 2017 11 153 4 166 19.5% 2.98 [0.97, 9.17]
Subtotal (95% CI) 400 409 63.0% 2.21 [1.20, 4.06]
Total events 34 14
Heterogeneity: Tau² = 0.00; Chi² = 2.31, df = 3 (P = 0.51); I² = 0%
Test for overall effect: Z = 2.55 (P = 0.01)
7.2.2 TKI
Dutton 2014 13 224 2 225 13.1% 6.53 [1.49, 28.60]
Wu 2018 8 158 8 156 23.8% 0.99 [0.38, 2.56]
Subtotal (95% CI) 382 381 37.0% 2.34 [0.35, 15.45]
Total events 21 10
Heterogeneity: Tau² = 1.47; Chi² = 4.65, df = 1 (P = 0.03); I² = 79%
Test for overall effect: Z = 0.88 (P = 0.38)
Total (95% CI) 782 790 100.0% 2.20 [1.20, 4.05]
Total events 55 24
Heterogeneity: Tau² = 0.17; Chi² = 7.13, df = 5 (P = 0.21); I² = 30%
Test for overall effect: Z = 2.54 (P = 0.01)
Test for subgroup differences: Chi² = 0.00, df = 1 (P = 0.95), I² = 0%
0.005 0.1 1 10 200
Favours [anti-EGFR] Favours [Control]

C




Figure S9 Subgroup analysis of grade 3–5 diarrhoea according to different treatment regimens: (A) different treatment modalities; (B) different anti-EGFR groups; (C) different anti-EGFR agents