

A


anti-EGFR
Control
Risk Ratio
Risk Ratio
Study or Subgroup Events Total Events Total Weight M-H, Fixed, 95% CI
M-H, Fixed, 95% CI
8.1.1 CRT
Castro 2018 1 53 0 54 12.4% 3.06 [0.13, 73.37]
Crosby 2013 9 129 0 129 12.5% 19.00 [1.12, 323.06]
Dutton 2014 5 224 1 225 25.0% 5.02 [0.59, 42.65]
Suntharalingam 2017 9 153 0 166 12.0% 20.60 [1.21, 351.02]
Wu 2018 11 158 1 156 25.2% 10.86 [1.42, 83.12]
Subtotal (95% CI) 717 730 87.1% 10.59 [3.56, 31.49]
Total events 35 2
Heterogeneity: Chi² = 1.43, df = 4 (P = 0.84); I² = 0%
Test for overall effect: Z = 4.24 (P < 0.0001)
8.1.2 CT
Lorenzen 2009 2 32 0 30 12.9% 4.70 [0.23, 94.01]
Subtotal (95% CI) 32 30 12.9% 4.70 [0.23, 94.01]
Total events 2 0
Heterogeneity: Not applicable
Test for overall effect: Z = 1.01 (P = 0.31)
Total (95% CI) 749 760 100.0% 9.83 [3.54, 27.28]
Total events 37 2
Heterogeneity: Chi² = 1.61, df = 5 (P = 0.90); I² = 0%
Test for overall effect: Z = 4.39 (P < 0.0001)
Test for subgroup differences: Chi² = 0.25, df = 1 (P = 0.62), I² = 0%
0.002 0.1 1 10 500
Favours [anti-EGFR] Favours [Control]

B


anti-EGFR
Control
Risk Ratio
Risk Ratio
Study or Subgroup Events Total Events Total Weight M-H, Fixed, 95% CI
M-H, Fixed, 95% CI
8.2.1 mAb
Castro 2018 1 53 0 54 12.4% 3.06 [0.13, 73.37]
Crosby 2013 9 129 0 129 12.5% 19.00 [1.12, 323.06]
Lorenzen 2009 2 32 0 30 12.9% 4.70 [0.23, 94.01]
Suntharalingam 2017 9 153 0 166 12.0% 20.60 [1.21, 351.02]
Subtotal (95% CI) 367 379 49.8% 11.71 [2.80, 48.99]
Total events 21 0
Heterogeneity: Chi² = 1.31, df = 3 (P = 0.73); I² = 0%
Test for overall effect: Z = 3.37 (P = 0.0007)
8.2.2 TKI
Dutton 2014 5 224 1 225 25.0% 5.02 [0.59, 42.65]
Wu 2018 11 158 1 156 25.2% 10.86 [1.42, 83.12]
Subtotal (95% CI) 382 381 50.2% 7.95 [1.84, 34.37]
Total events 16 2
Heterogeneity: Chi² = 0.27, df = 1 (P = 0.61); I² = 0%
Test for overall effect: Z = 2.78 (P = 0.005)
Total (95% CI) 749 760 100.0% 9.83 [3.54, 27.28]
Total events 37 2
Heterogeneity: Chi² = 1.61, df = 5 (P = 0.90); I² = 0%
Test for overall effect: Z = 4.39 (P < 0.0001)
Test for subgroup differences: Chi² = 0.14, df = 1 (P = 0.71), I² = 0%
0.005 0.1 1 10 200
Favours [anti-EGFR] Favours [Control]

C




Figure S10 Subgroup analysis of grade 3–5 acne-like rashes according to different treatment regimens: (A) different treatment modalities; (B) different anti-EGFR groups; (C) different anti-EGFR agents