Supplementary Material

Methods

Search strategy and screening

Table S1. PubMed Search Strategy

Search	Query
#1	(("lung diseases"[MeSH Terms] OR "chronic lung disease"[Text Word]) OR "chronic respiratory disease"[Text Word]) OR "cystic fibrosis"[Text Word]) OR "pulmonary arterial hypertension"[Text Word]) OR "pulmonary hypertension"[Text Word]) OR "ILD"[Text Word]) OR "pulmonary fibrosis"[Text Word]) OR "asthma"[Text Word]) OR "chronic obstructive pulmonary disease"[Text Word]) OR "COPD"[Text Word]) OR "lung transplantation"[MeSH Terms]) OR "bronch*"[Text Word]) OR "pulmonary disease"[Text Word]))
#2	(("interval"[Text Word] OR "intermittent"[Text Word]) OR "high intensity"[Text Word])
#3	(("exercise"[MeSH Terms] OR "exercise therapy"[MeSH Terms]) OR "rehabilitation"[MeSH Terms]) OR "rehabilit*"[Text Word]) OR "exercis*"[Text Word]) OR "train*"[Text Word]) OR "continuous"[Text Word]) OR "endurance"[Text Word]) OR "aerobic"[Text Word]) OR "moderate intensity"[Text Word]))
#4	(("exercise tolerance"[MeSH Terms] OR "exercise tolerance"[Text Word]) OR "exercise capacity"[Text Word]) OR "quality of life"[MeSH Terms]) OR "quality of life"[Text Word]) OR "oxygen consumption"[MeSH Terms]) OR "treatment outcome"[MeSH Terms]) OR "dyspnea"[MeSH Terms]) OR "dyspnea"[Text Word]) OR "physical fitness"[MeSH Terms]) OR "forced expiratory volume"[MeSH Terms]) OR "leg fatigue"[Text Word]) OR "leg discomfort"[Text Word]) OR "breathlessness"[Text Word])
#5	1 AND 2 AND 3 AND 4

Results



Figure S4. Comparison of the total volume of training between IET and CET for the duration of each programme as per study design.

Analysis S1. Meta-analysis and subgroup analysis

Figure S1.1. Comparison of the effect of IET versus CET on peak oxygen uptake ($VO2_{peak}$) in L/min.



Figure S1.2. Comparison of the effect of IET versus CET on peak heart rate (HR_{peak}) in beats/min. Subgroup analysis by BMI (<30Kg/m^2).



Figure S1.3. Comparison of the effect of IET versus CET on peak minute ventilation (VE_{peak}) in L/min. Subgroup analysis by BMI (<30Kg/m^2).



Figure S1.4. Comparison of the effect of IET versus CET on lactate threshold (LAT) in L/min. Subgroup analysis by BMI (<30Kg/m2).



Figure S1.5. Comparison of the effect of IET versus CET on leg discomfort (Borg's scale CR 0-10). Subgroup analysis by BMI (<30Kg/m2).



Analysis S2. Sensitivity Analysis

Figure S2.1. Forest plot of comparison between good to excellent quality studies and poor to fair quality studies rated on PEDro scale on peak work rate (WR_{peak}).



Figure S2.2. Forest plot of comparison between good to excellent quality studies and poor to fair quality studies rated on PEDro scale on peak oxygen uptake (VO2peak).



Figure S2.3. Forest plot of comparison between good to excellent quality studies and poor to fair quality studies rated on PEDro scale on peak heart rate (HR_{peak}).



Figure S2.4. Forest plot of comparison between good to excellent quality studies and poor to fair quality studies rated on PEDro scale on peak minute ventilation (VE_{peak}).



Figure S2.5. Forest plot of comparison between good to excellent quality studies and poor to fair quality studies rated on PEDro scale on lactate threshold (LAT).



Figure S2.6. Forest plot of comparison between good to excellent quality studies and poor to fair quality studies rated on PEDro scale on dyspnoea (Borg's scale CR 0-10).



Figure S2.7. Forest plot of comparison between good to excellent quality studies and poor to fair quality studies rated on PEDro scale on leg discomfort (Borg's scale CR 0-10).



Analysis S3. Publication bias

Figure S3.1. Funnel plot illustration of the impact of publication bias on WR_{peak}.