SUPPORTING DOCUMENTS

Table S1: Search strategy used for Databases

Database	Search terms
Pubmed/MEDLINE	(Africa OR African OR "West* Africa" OR "East* Africa" OR "South* Africa" OR "North* Africa" OR "Central Africa" OR Algeria OR Angola OR Benin OR Botswana OR "Burkina Faso" OR Burundi OR Cabo Verde OR Cameroon OR Central African Republic OR "CAR" OR Chad OR Comoros OR "DRC" OR Congo OR "Republic of Congo" OR "Ivory Coast" OR Cote d'Ivoire OR Djibouti OR Egypt OR "Equatorial Guinea" OR Eritrea OR Eswatini OR Swaziland OR Ethiopia OR Gabon OR Gambia OR Ghana OR Guinea OR "Guinea-Bissau" OR Kenya OR Lesotho OR Liberia OR Libya OR Madagascar [PL] OR Malawi OR Mali OR Mauritania OR Mauritanie OR Mauritius OR Morocco OR Mozambique OR Namibia OR Niger OR Nigeria OR Rwanda OR "Sao Tome and Principe" OR Senegal OR Seychelles OR "Sierra Leone" OR Somalia OR "South Africa" OR "South Sudan" OR Sudan OR Tanzania OR Togo OR Tunisia OR Uganda OR Zambia OR Zimbabwe) AND (TNBC OR "triple-negative*" OR "triple negative*" OR "triple negative") AND (rate* OR prevalence OR incidence OR epidemiology)
Web of Science	TS = (Africa or African or "West* Africa" or "East* Africa" or "South* Africa" or "North* Africa" or "Central Africa" or Algeria or Angola OR Benin OR Botswana OR "Burkina Faso" OR Burundi OR Cabo Verde OR Cameroon OR Central African Republic OR "CAR" OR Chad OR Comoros OR "DRC" OR Congo OR "Republic of Congo" OR "Ivory Coast" OR Cote d'Ivoire OR Djibouti OR Egypt OR "Equatorial Guinea" OR Eritrea OR Eswatini OR Swaziland OR Ethiopia OR Gabon OR Gambia OR Ghana OR Guinea OR "Guinea-Bissau" OR Kenya OR Lesotho OR Liberia OR Libya OR Madagascar [PL] OR Malawi OR Mali OR Mauritania OR Mauritanie OR Mauritius OR Morocco OR Mozambique OR Namibia OR Niger OR Nigeria OR Rwanda OR "Sao Tome and Principe" OR Senegal OR Seychelles OR "Sierra Leone" OR Somalia OR "South Africa" OR "South Sudan" OR Sudan OR Tanzania OR Togo OR Tunisia OR Uganda OR Zambia OR Zimbabwe) AND TS= (TNBC OR "triple-negative*" OR "triple negative*" OR "triple negative") AND TS= (rate* OR prevalence OR incidence OR epidemiology)
Embase	(Africa OR African OR "West* Africa" OR "East* Africa" OR "South* Africa" OR "North* Africa" OR "Central Africa" OR Algeria OR Angola OR Benin OR Botswana OR "Burkina Faso" OR Burundi OR Cabo Verde OR Cameroon OR Central African Republic OR "CAR" OR Chad OR Comoros OR "DRC" OR Congo OR "Republic of Congo" OR "Ivory Coast" OR Cote d'Ivoire OR Djibouti OR Egypt OR "Equatorial Guinea" OR Eritrea OR Eswatini OR Swaziland OR Ethiopia OR Gabon OR Gambia OR Ghana OR Guinea OR "Guinea-Bissau" OR Kenya OR Lesotho OR Liberia OR Libya OR Madagascar OR Malawi OR Mali OR Mauritania OR Mauritanie OR Mauritius OR Morocco OR Mozambique OR Namibia OR Niger OR Nigeria OR Rwanda OR "Sao Tome and Principe" OR Senegal OR Seychelles OR "Sierra Leone" OR Somalia OR "South Africa" OR "South Sudan" OR Sudan OR Tanzania OR Togo OR Tunisia OR Uganda OR Zambia OR Zimbabwe) AND (TNBC OR "triple-negative*" OR "triple negative*" OR "triple negative") AND (rate* OR prevalence OR incidence OR epidemiology)
African Journals Online (AJOL)	(TNBC OR "triple-negative*" OR "triple negative*" OR "triple negative") AND (rate* OR prevalence OR incidence OR epidemiology)

Table S2: Summary of data from included studies

Author	Population	Region	Mean age ± SD	Median age (min-max)	Grade 3 tumors	Positive Nodes	Number of TNBC cases	Participants studied with known receptor status (all participants)	Frequency of TNBC	Risk of Bias Assessment
Adani-Ifè A, 2020 [1]	Togo	West Africa	52.05 ±12.38	(30-85)	42.70%	41.90%	44	117 (312)	37.60%	Moderate
Adisa AO, 2011 [2]	Nigeria	West Africa	45.90	(20-81)	/	/	30	64 (385)	46.88%	High
Agboola A, 2014 [3]	Nigeria	West Africa	/	/	35.70%	99.20%	99	155 (199)	54.36%	High
Agboola A, 2014 [4]	Nigeria	West Africa	/	/	/	94.20%	55	100 (308)	55.00%	Moderate
Agboola A, 2013 [5]	Nigeria	West Africa	/	/	/	92.70%	81	149 (308)	63.87%	High
Agboola A, 2016 [6]	Nigeria	West Africa	/	/	34.12%	91.76%	114	187 (255)	60.96%	Moderate
Agboola A, 2014 [7]	Nigeria	West Africa	/	/	38.50%	92.20%	115	231 (231)	49.78%	High
Akinyi MO, 2019 [8]	Kenya	East Africa	48.00 ±14.5	/	/	71%	22	79 (79)	27.85%	High
Alshenawy HA, 2012 [9]	Egypt	North Africa	58.30 ±10.1	/	20.00%	/	48	150 (150)	26.67%	Moderate
Ayadi EZ, 2018 [10]	Tunisia	North Africa	35.00 ±4.2	(17-40)	46.85%	64.10%	47	203 (238)	23.15%	Moderate
Babalanda JNP, 2014 [11]	Uganda	East Africa	47	(13-87)	53.51%	77.19%	23	114 (114)	20.18%	Moderate
Bahnassy A, 2015 [12]	Egypt	North Africa	TNBC: 51.91± 12.34 non-TNBC: 52.77 ± 12.13 Controls: 35 ± 13.94	TNBC: (30-78) non-TNBC: (27-81) Controls: (22-64)	15.63%	72.91%	43	96 (96)	44.79%	Moderate
Bakkach J, 2017 [13]	Morocco	North Africa	/	36 (/)	40.20%	58.50%	19	82 (331)	23.00%	High
Bekkouche Z, 2013 [14]	Algeria	North Africa	49.72	(30-84)	73.33%	35.83%	30	120 (120)	25.00%	High
Belhadj A, 2021 [15]	Algeria	North Africa	/	50 (21-88)	38.86%	64.70%	286	1066 (1140)	27.00%	Moderate
Ben Ayed-Guerfali D, 2021 [16]	Tunisia	North Africa	/	42.78 (27-65)	26.78%	59.70%	26	110 (134)	23.60%	High
Ben Gacem R, 2011 [17]	Tunisia	North Africa	48.70	47 (31-75)	43.62%	55.26%	28	94 (94)	29.79%	Moderate
Bhatia RK, 2019 [18]	Botswana	Southern Africa	/	54 (27-96)	25.76%	/	45	211 (384)	21.30%	Low

Bogan D, 2017 [19]	Egypt	North Africa	52.38 ±12.17	(27-81)	13.94%	66.67%	43	96 (96)	44.79%	Moderate
Brandão M, 2020 [20]	Mozambique	East Africa	/	48 (/)	31.25%	61.43%	52	210 (262)	25.00%	Moderate
Chahbouni S, 2017 [21]	Morocco	North Africa	48.00	(22-99)	/	/	53	371 (390)	13.59%	High
Cherbal F, 2015 [22]	Algeria	North Africa	/	48.5 (22-84)	31.30%	/	627	3014 (3014)	20.80%	Moderate
Corsini C, 2017 [23]	Multiple (France, Algeria, Egypt, Morocco, Tunisia)	North Africa	/	/	44.74%	42%	35	174 (456)	20.10%	Moderate
Cubasch H, 2018 [24]	South Africa	Southern Africa	54.40 ±14.2	/	39.53%	/	126	581 (602)	21.70%	Low
Debouki-Joudi S, 2016 [25]	Tunisia	North Africa	49.30	(24-80)	26.22%	55.20%	53	80 (91)	66.25%	High
Dickens C, 2014 [26]	Multiple (South Africa & Namibia)	Southern Africa	/	/	33.87%	/	1482	7246 (12361)	20.45%	Low
Ekpe E, 2019 [27]	Kenya	East Africa	47.00 ±12	(21-79)	50.00%	/	24	120 (125)	20.00%	Moderate
El-Amir MI, 2020 [28]	Egypt	North Africa	54.20 ±10.6	(32-77)	18.00%	72%	3	50 (50)	6.00%	High
Elidrissi Errahhali M, 2017 [29]	Morocco	North Africa	48.70	(19-105)	18.00%	64.80%	320	2260 (2406)	14.20%	Moderate
Fitzpatrick MB, 2019 [30]	Senegal	West Africa	47.00 ±12.7	/	76.17%	/	92	197 (522)	46.70%	Moderate
Fouhi ME, 2020 [31]	Morocco	North Africa	51.60	(23-89)	32.60%	35.70%	87	568 (668)	15.30%	High
Gaceb H, 2017 [32]	Algeria	North Africa	48.44	(22-84)	28.50%	/	175	877 (877)	19.95%	Low
Groenewald C, 2019 [33]	South Africa	Southern Africa	53.60 ±13.4	/	/	68.10%	71	504 (505)	14.10%	Moderate
Guedouar Y, 2014 [34]	Algeria	North Africa	47.00	(23-78)	24.58%	38.33%	60	240 (240)	25.00%	High
Hachana M, 2012 [35]	Tunisia	North Africa	/	/	40.18%	21.40%	32	123 (123)	26.02%	Moderate
Hadgu E, 2020 [36]	Ethiopia	East Africa	43.00 ±14	50 (22-75)	50.00%	/	25	110 (112)	23.00%	Moderate
Huo D, 2009 [37]	Multiple (Nigeria & Senegal)	West Africa	44.80	/	39.15%	72%	103	271 (378)	27.00%	Low
Hussein O, 2013 [38]	Egypt	North Africa	53.00 ±11.6	52	17.39%	/	50	263 (263)	19.00%	High

Kakudji BK, 2021 [39]	South Africa	Southern Africa	56.35 ±14.09	/	62.07%	64.66%	26	116 (116)	22.50%	High
Kallel-Bayoudh I, 2011 [40]	Tunisia	North Africa	50.00 ±11	(27-75)	35.44%	65%	11	84 (84)	13.00%	High
Khalil AI, 2016 [41]	Morocco	North Africa	50.20 ±11.34	(17-93)	35.48%	49.40%	146	889 (1277)	16.40%	Moderate
Luyey Mvila G, 2015 [42]	Democratic Republic of Congo	Central Africa	50.00	49 (29-82)	47.13%	/	13	87 (87)	14.94%	Moderate
Ly M, 2012 [43]	Mali	West Africa	46.00	(25-82)	78.18%	91%	53	114 (114)	46.00%	Moderate
Manai M, 2019 [44]	Tunisia	North Africa	/	42 (24-62)	42.20%	95%	49	194 (210)	25.00%	Moderate
McCormack VA, 2013 [45]	South Africa	Southern Africa	55.30	/	42.34%	/	209	1027 (1247)	20.40%	Low
Mehemmai C, 2019 [46]	Algeria	North Africa	/	48.5 (21-84)	26.99%	/	243	1144 (1162)	21.24%	Moderate
Mejri N, 2020 [47]	Tunisia	North Africa	/	48.5 IBCa 49 non-IBCa (/)	46.20% IBCa 36.10% non-IBCa	50%	121	740 (740)	16.40%	High
Mouh FZ, 2020 [48]	Morocco	North Africa	/	47 (/)	34.92%	60.10%	85	500 (500)	17.00%	High
Newman LA, 2019 [49]	Multiple (Ghana, Ethiopia, United States)	West and East Africa	/	56 (14-95)	/	/	Ghanaian: 142 Ethiopian: 24	Ghana: 292 (618) Ethiopia: 144 (252)	Ghana: 49.00% Ethiopia: 17.00%	High
Ohene-Yeboah M, 2012 [50]	Ghana	West Africa	49.10 ±15.3	(20-96)	53.70%	/	23	54 (330)	42.70%	Moderate
Ramdas Y, 2020 [51]	South Africa	Southern Africa	60.80 ±9.3	60 (43-86)	9.57%	12.10%	6	107 (107)	5.60%	High
Rayne S, 2019 [52]	South Africa	Southern Africa	/	56	51.10%	/	31	223 (365)	14.00%	Moderate
Roy I, 2011 [53]	Uganda	East Africa	/	/	40.00%	/	16	44 (64)	36.00%	Moderate
Samaka RM, 2016 [54]	Egypt	North Africa	/	(30-81)	44.44%	77.80%	19	81 (81)	23.50%	Moderate
Sawe RT, 2017 [55]	Kenya	East Africa	51.90	48.5 (16-84)	42.22%	/	18	49 (49)	36.70%	Moderate
Schwartz T, 2012 [56]	Ghana	West Africa	49.00 ±13.4	/	52.52%	/	58	103 (104)	56.30%	Moderate
Sengal AT, 2017 [57]	Sudan	North Africa	48.80 ±13.5	(20-90)	41.79%	52.70%	143	415 (560)	34.50%	Moderate
Seshie B, 2015 [58]	Ghana	West Africa	52.50 ±12.1	(24-77)	31.58%	66.00%	77	156 (165)	49.40%	Moderate
Shaikh AJ, 2018 [59]	Kenya	East Africa	54.00	(27-83)	33.33%	49%	21	116 (123)	17.00%	Moderate
Stark A, 2010 [60]	Ghana	West Africa	48.00 ±13.7	/	76.00%	/	37	45 (75)	82.20%	Moderate

Sung H, 2019 [61]	Multiple (USA, Caribbean, West Africa, East Africa)	West and East Africa	/	/	West Africa: 56.92% East Africa: 43.95%	/	West Africa: 173 East Africa: 51	West Africa: 717 (811) East Africa: 441 (484)	West Africa: 24.10% East Africa: 11.60%	Moderate
Tanimowo MO, 2019 [62]	Nigeria	West Africa	43.07 ±11.19	(23-70)	29.51%	/	38	61 (61)	62.30%	High
Traoré B, 2019 [63]	Guinea	West Africa	48.20 ±10.9	(31-80)	36.21%	62.10%	15	58 (58)	25.90%	Moderate
Ukah CO, 2017 [64]	Nigeria	West Africa	46.30 ±12.3	(24-85)	26.83%	/	50	123 (397)	40.70%	Moderate
Uyisenga JP, 2020 [65]	Rwanda	East Africa	49.70 ±12.98	(17-86)	/	70.06%	52	138 (138)	37.70%	Low
Youngblood VM, 2020 [66]	Malawi	East Africa	/	49 (21-80)	37.00%	77%	25	100 (100)	25.00%	Low
Znati K, 2012 [67]	Morocco	North Africa	30.62	(18-35)	45.07%	51%	13	74 (74)	18.00%	High

/ data not stated

IBCa- inflammatory breast cancer

Table S3: Questions for Risk of Bias Assessment, modified from Hoy *et al*., 2012

1. Was the study's target population a close representation of the national population in relation to relevant variables, e.g. age, sex, occupation?
2. Was the sampling frame a true or close representation of the target population?
3. Was some form of random selection used to select the sample, or was a census undertaken?
4. Was the likelihood of non-response bias minimal?
5. Were data collected directly from the subjects (as opposed to a proxy)?
6. Was an acceptable case definition used in the study? Was TNBC defined as ER, PR, HER2 negative?
7. Was the study instrument that measured the parameter of interest (e.g. appropriate guidelines for cut-offs ASCO-CAP) shown to have reliability and validity?
8. Was the same mode of data collection used for all subjects? Was the information from each patient collected in the same way (e.g clinical referral appointment)?
9. Was there a shortest prevalence or incidence period identified?
10.Were the numerator(s) and denominator(s) for the parameter of interest appropriate?

Table S4: Raw Values for Risk of Bias Assessment for Included Studies

Author	Q1	Q2	Q3	Q4	Q5	Q6	Q7	Q8	Q9	Q10	TOTAL	Overall ROB
Adani-Ifè A, 2020 [1]	0	0	0	0	1	1	1	1	1	1	6	**Moderate**
Adisa AO, 2011 [2]	0	0	1	0	1	0	0	0	1	0	3	**High**
Agboola A, 2014 [3]	0	0	0	0	1	1	1	1	1	0	5	**High**
Agboola A, 2014 [4]	0	0	1	0	1	1	1	1	1	1	7	**Moderate**
Agboola A, 2013 [5]	0	0	0	0	1	1	1	1	0	1	5	**High**
Agboola A, 2016 [6]	0	0	0	1	1	0	1	1	1	1	6	**Moderate**
Agboola A, 2014 [7]	0	0	0	0	1	0	1	1	1	1	5	**High**
Akinyi MO, 2019 [8]	0	0	0	1	1	1	0	1	0	0	4	**High**
Alshenawy HA, 2012 [9]	0	0	1	1	1	1	0	1	1	1	7	**Moderate**
Ayadi EZ, 2018 [10]	0	0	0	1	1	1	1	1	1	1	7	**Moderate**
Babalanda JNP, 2014 [11]	0	0	1	1	1	1	1	1	1	1	8	**Moderate**
Bahnassy A, 2015 [12]	0	0	1	1	1	1	0	1	1	1	7	**Moderate**
Bakkach J, 2017 [13]	0	0	0	1	1	0	0	1	1	0	4	**High**
Bekkouche Z, 2013 [14]	0	0	1	0	1	1	0	1	1	0	5	**High**

Belhadj A, 2021 [15]	1	0	1	1	1	1	0	1	1	1	8	**Moderate**
Ben Ayed-Guerfali D, 2021 [16]	0	0	0	0	1	0	0	1	1	0	3	**High**
Ben Gacem R, 2011 [17]	0	0	1	1	1	1	0	1	0	1	6	**Moderate**
Bhatia RK, 2019 [18]	1	1	1	1	1	1	1	1	1	1	10	**Low**
Bogan D, 2017 [19]	0	0	0	1	1	1	1	1	1	1	7	**Moderate**
Brandão M, 2020 [20]	0	0	1	1	1	1	1	0	1	1	7	**Moderate**
Chahbouni S, 2017 [21]	0	0	0	0	1	1	0	1	1	0	4	**High**
Cherbal F, 2015 [22]	1	1	1	0	1	1	0	1	1	1	8	**Moderate**
Corsini C, 2017 [23]	0	0	1	1	1	0	0	1	1	1	6	**Moderate**
Cubasch H, 2018 [24]	1	1	1	1	1	1	0	1	1	1	9	**Low**
Debouki-Joudi S, 2016 [25]	0	0	0	0	1	1	0	1	0	1	4	**High**
Dickens C, 2014 [26]	1	1	1	1	1	1	1	1	1	1	10	**Low**
Ekpe E, 2019 [27]	0	0	0	0	1	1	1	1	1	1	6	**Moderate**
El-Amir MI, 2020 [28]	0	0	0	0	1	0	0	1	1	1	4	**High**
Elidrissi Errahhali M, 2017 [29]	0	1	1	0	1	1	0	1	1	1	7	**Moderate**
Fitzpatrick MB, 2019 [30]	0	0	0	0	1	1	1	1	1	1	6	**Moderate**
Fouhi ME, 2020 [31]	0	1	0	0	1	0	0	1	1	0	4	**High**
Gaceb H, 2017 [32]	1	1	1	0	1	1	1	1	1	1	9	**Low**
Groenewald C, 2019 [33]	0	0	0	1	1	1	0	1	1	1	6	**Moderate**
Guedouar Y, 2014 [34]	0	1	0	0	1	1	0	1	1	0	5	**High**
Hachana M, 2012 [35]	0	0	1	1	1	1	0	1	1	1	7	**Moderate**
Hadgu E, 2020 [36]	0	0	0	1	1	1	1	1	1	1	7	**Moderate**
Huo D, 2009 [37]	0	1	1	1	1	1	1	1	1	1	9	**Low**
Hussein O, 2013 [38]	0	0	1	0	1	0	0	1	1	1	5	**High**
Kakudji BK, 2021 [39]	0	0	0	1	1	1	0	0	1	1	5	**High**
Kallel-Bayoudh I, 2011 [40]	0	0	0	0	1	1	0	1	1	1	5	**High**
Khalil AI, 2016 [41]	0	1	1	0	1	1	0	1	1	1	7	**Moderate**
Luyey Mvila G, 2015 [42]	0	0	1	0	1	1	1	1	1	1	7	**Moderate**
Ly M, 2012 [43]	0	0	1	1	1	1	1	1	1	1	8	**Moderate**
Manai M, 2019 [44]	0	1	1	0	1	1	0	1	1	1	7	**Moderate**
McCormack VA, 2013 [45]	1	1	1	1	1	1	0	1	1	1	9	**Low**
Mehemmai C, 2019 [46]	0	0	1	1	1	1	1	1	1	1	8	**Moderate**
Mejri N, 2020 [47]	0	0	0	0	1	0	0	1	1	0	3	**High**
Mouh FZ, 2020 [48]	0	0	0	0	1	1	0	1	1	1	5	**High**
Newman LA, 2019 [49]	0	0	0	0	1	0	1	1	1	1	5	**High**
Ohene-Yeboah M, 2012 [50]	0	0	1	0	1	1	0	1	1	1	6	**Moderate**

Ramdas Y, 2020 [51]	0	0	0	1	1	0	0	1	1	1	5	**High**
Rayne S, 2019 [52]	0	0	1	0	1	1	1	0	1	1	6	**Moderate**
Roy I, 2011 [53]	0	0	0	0	1	1	1	1	1	1	6	**Moderate**
Samaka RM, 2016 [54]	0	0	0	1	1	1	1	1	1	1	7	**Moderate**
Sawe RT, 2017 [55]	0	1	1	0	1	1	0	1	1	0	6	**Moderate**
Schwartz T, 2012 [56]	0	0	0	1	1	1	0	1	1	1	6	**Moderate**
Sengal AT, 2017 [57]	0	0	1	1	1	1	1	1	1	1	8	**Moderate**
Seshie B, 2015 [58]	0	1	1	1	1	1	0	1	1	0	7	**Moderate**
Shaikh AJ, 2018 [59]	0	0	1	1	1	1	1	1	1	1	8	**Moderate**
Stark A, 2010 [60]	0	0	1	0	1	1	1	1	1	0	6	**Moderate**
Sung H, 2019 [61]	0	1	1	1	1	1	0	1	1	1	8	**Moderate**
Tanimowo MO, 2019 [62]	0	0	0	1	1	0	1	0	1	1	5	**High**
Traoré B, 2019 [63]	0	1	1	0	1	1	0	1	1	0	6	**Moderate**
Ukah CO, 2017 [64]	0	1	0	1	1	1	0	1	1	1	7	**Moderate**
Uyisenga JP, 2020 [65]	0	1	1	1	1	1	1	1	1	1	9	**Low**
Youngblood VM, 2020 [66]	0	1	1	1	1	1	1	1	1	1	9	**Low**
Znati K, 2012 [67]	0	0	0	0	1	0	0	1	1	0	3	**High**

Table S5: Pooled TNBC Frequcny in Africa by Country

Country (no. of studies)	Total participants	Pooled TNBC Frequency, % [95% CI]	Heterogeneity (I^2), %	Weight, %
Algeria (*n*= 6)	6461	22.7 [20.3- 25.2]	77	12.0
Botswana (*n*= 1)	211	21.3 [16.3- 27.4]	not applicable	1.9
Democratic Republic of Congo (*n*= 1)	87	14.9 [8.9- 24.1]	not applicable	1.6
Egypt (*n*= 5)	640	24.5 [15.3- 36.7]	88	8.3
Ethiopia (*n*= 1)	110	22.7 [15.8- 31.5]	not applicable	1.8
Ghana (*n*= 4)	358	57.2 [43.6- 69.8]	82	6.8
Guinea (*n*= 1)	58	25.9 [16.2- 38.6]	not applicable	1.6
Kenya (*n*= 4)	364	24.5 [17.8- 32.8]	63	6.9
Malawi (*n*= 1)	100	25.0 [17.5- 34.4]	not applicable	1.8
Mali (*n*= 1)	114	46.5 [37.6- 55.7]	not applicable	1.9
Morocco (*n*= 7)	4744	15.6 [14.3- 17.1]	31	13.4
Mozambique (*n*= 1)	210	24.8 [19.4- 31.0]	not applicable	1.9
Nigeria (*n* = 4)	479	49.2 [41.4- 57.0]	61	7.3
Rwanda (*n*= 1)	138	37.7 [30.0- 46.0]	not applicable	1.9
Senegal (*n*= 1)	197	46.7 [39.8- 53.7]	not applicable	2.0
South Africa (*n*= 4)	950	14.0 [9.6- 19.8]	75	7.0
Sudan (*n*= 1)	415	34.5 [30.0- 39.2]	not applicable	2.0
Togo (*n*= 1)	117	37.6 [29.3- 46.7]	not applicable	1.9
Tunisia (*n*= 8)	1628	26.4 [18.5- 36.2]	92	14.6
Uganda (*n*= 2)	158	27.0 [14.3- 45.1]	77	3.4
Total	17539	27.0 [24.0- 30.2]	94	100

Figure S1: Risk of bias assessment of all articles by (A) region, (B) country and (C) overall risk of bias assessment for all articles included. Southern Africa had the highest percentage of low risk of bias. Multiple denotes more than one country and/or region or was investigated within that study.




Figure S2: Risk of bias (ROB) assessment per question for all articles included in systematic review. High risk of bias was noted specifically for questions on data acquisition




Figure S3: Pooled TNBC frequency in Africa by risk of bias assessment. Cases are defined as participants in a study that were identified as triple negative, and total is the number of breast cancer participants with known receptor status in the study.




Figure S4: Pooled TNBC frequency in Africa by use of validated tool for receptor status. Cases are defined as participants in a study that were identified as triple negative, and total is the number of breast cancer participants with known receptor status in the study.




Figure S5: Meta-regression shows association with year of publication but not with age or use of validated tool for receptor status. The association of use of validated tool to assess receptor status (A), year of publication (B), mean age at diagnosis (C) and median age at diagnosis (D) on heterogeneity of treatment effects between studies was investigated by a meta-regression. P <0.05 indicates significant association.




Figure S6: Influence analysis highlights studies that overly contribute to heterogeneity. Influence analysis via Baujat plot shows studies that two studies [7] and [29] contribute largely to overall heterogeneity.




Figure S7: Egger's test shows high publication bias. Publication bias between studies for TNBC frequency investigated using Egger's test and a significant association was found, inferring significant publication bias. P <0.05 indicates significant association.





Figure S8: Funnel plot shows high publication bias. Publication bias between studies was investigated using funnel plot to visualize asymmetry and multiple studies do not follow the funnel shape inferring significant publication bias. P <0.05 indicates significant association.



References for studies included in review:

1. Adani-Ife, A., et al., *Breast cancer in togolese women: immunohistochemistry subtypes.* Bmc Womens Health, 2020. **20**(1).
2. Adisa, A.O., et al., *Metastatic breast cancer in a Nigerian tertiary hospital.* Afr Health Sci, 2011. **11**(2): p. 279-84.
3. Agboola, A.O.J., et al., *Clinicopathological and molecular significance of Sumolyation marker (ubiquitin conjugating enzyme 9 (UBC9)) expression in breast cancer of black women.* Pathology - Research and Practice, 2014. **210**(1): p. 10-17.
4. Agboola, A., et al., *Molecular profiling of breast cancer in Nigerian women identifies an altered p53 pathway as a major mechanism underlying its poor prognosis compared with British counterpart.* The Malaysian journal of pathology, 2014. **36 1**: p. 3-17.
5. Agboola, A.O.J., et al., *Cell Proliferation (KI-67) Expression Is Associated with Poorer Prognosis in Nigerian Compared to British Breast Cancer Women.* ISRN Oncology, 2013. **2013**: p. 675051.
6. Agboola, A., et al., *Tumour cell membrane laminin expression is associated with basal-like phenotype and poor survival in Nigerian breast cancer.* The Malaysian journal of pathology, 2016. **38**: p. 83-92.
7. Agboola, A., et al., *PIASγ expression in relation to clinicopathological, tumour factors and survival in indigenous black breast cancer women.* Journal of Clinical Pathology, 2014. **67**(4): p. 301.
8. Akinyi, M.O., et al., *Correlation between receptor status and presence of axillary lymph node metastasis in breast cancer in Kenya.* Annals of African Surgery, 2019. **16**(2).
9. Alshenawy, H.A., *Prevalence of androgen receptors in invasive breast carcinoma and its relation with estrogen receptor, progesterone receptor and Her2/neu expression.* Journal of the Egyptian National Cancer Institute, 2012. **24**(2): p. 77-83.
10. Ayadi, E.Z., et al., *Prognostic Value of BCL2 in Women Patients with Invasive Breast Cancer.* Asian Pac J Cancer Prev, 2018. **19**(12): p. 3557-3564.
11. Babalanda, J., et al., *Receptor Status and Associated Clinico- Histopathological Characteristics among Women with Breast Cancer in a Ugandan Tertiary Hospital.* The Annals of African Surgery, 2014. **11**.
12. Bahhnassy, A., et al., *Transforming growth factor-β, insulin-like growth factor I/insulin-like growth factor I receptor and vascular endothelial growth factor-A: Prognostic and predictive markers in triple-negative and non-triple-negative breast cancer.* Mol Med Rep, 2015. **12**(1): p. 851-864.
13. Bakkach, J., et al., *Clinicopathologic and prognostic features of breast cancer in young women: a series from North of Morocco.* BMC Women's Health, 2017. **17**(1): p. 106.
14. Bekkouche, Z., et al., *Caractéristiques des carcinomes mammaires triple-négatifs dans l'Ouest-algérien.* Journal Africain du Cancer / African Journal of Cancer, 2013. **5**(3): p. 155-161.
15. Belhadj, A., et al., *Prevalence and prognosis of molecular phenotypes in breast cancer patients by age: a population-based retrospective cohort study in western Algeria.* Pan Afr Med J, 2021. **38**: p. 88.
16. Ben Ayed-Guerfali, D., et al., *Novel and recurrent BRCA1/BRCA2 germline mutations in patients with breast/ovarian cancer: a series from the south of Tunisia.* J Transl Med, 2021. **19**(1): p. 108.
17. Ben Gacem, R., et al., *Clinicopathologic significance of DNA methyltransferase 1, 3a, and 3b overexpression in Tunisian breast cancers.* Human Pathology, 2012. **43**(10): p. 1731-1738.

18. Bhatia, R.K., et al., *Report of clinico-pathological features of breast cancer in HIV-infected and uninfected women in Botswana.* Infect Agent Cancer, 2019. **14**: p. 28.
19. Bogan, D., et al., *The role of BRCA1-IRIS in the development and progression of triple negative breast cancers in Egypt: possible link to disease early lesion.* BMC Cancer, 2017. **17**(1): p. 329.
20. Brandão, M., et al., *Breast cancer subtypes: implications for the treatment and survival of patients in Africa-a prospective cohort study from Mozambique.* ESMO Open, 2020. **5**(5): p. e000829.
21. Chahbouni, S., et al. *CLINICOPATHOLOGIC FEATURES AND MOLECULAR SUBTYPES OF BREAST CANCER IN FEZ-MEKNES REGION (MOROCCO): A STUDY OF 390 PATIENTS.* 2017.
22. Cherbal, F., et al., *Distribution of molecular breast cancer subtypes among Algerian women and correlation with clinical and tumor characteristics: A population-based study.* Breast disease, 2015. **35**: p. 95-102.
23. Corsini, C., et al., *Early onset breast cancer: differences in risk factors, tumor phenotype, and genotype between North African and South European women.* Breast Cancer Research and Treatment, 2017. **166**(2): p. 631-639.
24. Cubasch, H., et al., *Breast cancer survival in Soweto, Johannesburg, South Africa: A receptor-defined cohort of women diagnosed from 2009 to 11.* Cancer Epidemiology, 2018. **52**: p. 120-127.
25. Debouki-Joudi, S., et al., *CpG methylation of APC promoter 1A in sporadic and familial breast cancer patients.* Cancer Biomarkers, 2016. **16**: p. 1-9.
26. Dickens, C., et al., *Racial Comparison of Receptor-Defined Breast Cancer in Southern African Women: Subtype Prevalence and Age–Incidence Analysis of Nationwide Cancer Registry Data.* Cancer Epidemiology Biomarkers & Prevention, 2014. **23**(11): p. 2311.
27. Ekpe, E., et al., *Metastatic Breast Cancer in Kenya: Presentation, Pathologic Characteristics, and Patterns-Findings From a Tertiary Cancer Center.* Journal of Global Oncology, 2019. **5**.
28. El-Amir, M.I., et al., *Role of ATG16LI (rs2241880) and Interleukin 10 (rs1800872) Polymorphisms in Breast Cancer Among Egyptian Patients.* Egypt J Immunol, 2020. **27**(1): p. 65-76.
29. Elidrissi Errahhali, M., et al., *First report on molecular breast cancer subtypes and their clinico-pathological characteristics in Eastern Morocco: series of 2260 cases.* BMC Women's Health, 2017. **17**(1): p. 3.
30. Fitzpatrick, M.B., et al., *Pathology of Senegalese breast cancers.* Pan Afr Med J, 2019. **34**: p. 67.
31. Fouhi, M.E., et al., *[Epidemiological and anatomopathological profile of breast cancer at the Ibn Rochd University Hospital, Casablanca].* Pan Afr Med J, 2020. **37**: p. 41.
32. Gaceb, H., et al., *Clinicopathological and Molecular Study of Triple-Negative Breast Cancer in Algerian Patients.* Pathology & Oncology Research, 2017. **24**(2): p. 297-308.
33. Groenewald, C., et al., *Axillary lymph node dissection for patients with invasive breast cancer at Charlotte Maxeke and Chris Hani Baragwanath Academic Hospitals.* S Afr J Surg, 2019. **57**(4): p. 18-24.
34. Guedouar, Y., et al., *Évaluation phénotypique des sous-types moléculaires en carcinologie mammaire dans une population de l'Ouest algérien.* Journal Africain du Cancer / African Journal of Cancer, 2014. **6**(3): p. 150-158.
35. Hachana, M., et al., *Investigation of human JC and BK polyomaviruses in breast carcinomas.* Breast Cancer Research and Treatment, 2012. **133**(3): p. 969-977.

36. Hadgu, E., et al., *Distribution and characteristics of androgen receptor (AR) in breast cancer among women in Addis Ababa, Ethiopia: A cross sectional study.* Plos One, 2020. **15**(5).
37. Huo, D., et al., *Population Differences in Breast Cancer: Survey in Indigenous African Women Reveals Over-Representation of Triple-Negative Breast Cancer.* Journal of Clinical Oncology, 2009. **27**(27): p. 4515-4521.
38. Hussein, O., et al., *Hormone Receptors and Age Distribution in Breast Cancer Patients at a University Hospital in Northern Egypt.* Breast Cancer: Basic and Clinical Research, 2013. **7**: p. BCBCR.S12214.
39. Kakudji, B.K., et al., *Breast cancer molecular subtypes and receptor status among women at Potchefstroom Hospital: a cross-sectional study.* Pan Afr Med J, 2021. **38**: p. 85.
40. Kallel-Bayoudh, I., et al., *Bcl-2 expression and triple negative profile in breast carcinoma.* Medical Oncology, 2011. **28**(1): p. 55-61.
41. Khalil, A., et al., *Cancer du sein au Maroc: profil phénotypique des tumeurs.* The Pan African Medical Journal, 2016. **25**.
42. Luyeye Mvila, G., et al., *Prognostic features of breast cancer differ between women in the Democratic Republic of Congo and Belgium.* The Breast, 2015. **24**(5): p. 642-648.
43. Ly, M., et al., *High Incidence of Triple-Negative Tumors in Sub-Saharan Africa: A Prospective Study of Breast Cancer Characteristics and Risk Factors in Malian Women Seen in a Bamako University Hospital.* Oncology, 2012. **83**(5): p. 257-263.
44. Manai, M., et al., *Inflammatory breast cancer in 210 patients: A retrospective study on epidemiological, anatomo-clinical features and therapeutic results.* Mol Clin Oncol, 2019. **10**(2): p. 223-230.
45. McCormack, V.A., et al., *Breast cancer receptor status and stage at diagnosis in over 1,200 consecutive public hospital patients in Soweto, South Africa: a case series.* Breast cancer research : BCR, 2013. **15**(5): p. R84-R84.
46. Mehemmai, C., et al., *BRCA1 and BRCA2 Germline Mutation Analysis in Hereditary Breast/Ovarian Cancer Families from the Aures Region (Eastern Algeria): First Report.* Pathology & Oncology Research, 2019. **26**(2): p. 715-726.
47. Mejri, N., et al., *Reproductive Risk Factors of Inflammatory Breast Cancer according to Luminal, HER2-Overexpressing, and Triple-Negative Subtypes: A Case Comparison Study.* Oncol Res Treat, 2020. **43**(5): p. 204-210.
48. Mouh, F.Z., et al., *Clinicopathological, Treatment and Event-Free Survival Characteristics in a Moroccan Population of Triple-Negative Breast Cancer.* Breast Cancer (Auckl), 2020. **14**: p. 1178223420906428.
49. Newman, L.A., et al., *Hereditary Susceptibility for Triple Negative Breast Cancer Associated With Western Sub-Saharan African Ancestry: Results From an International Surgical Breast Cancer Collaborative.* Ann Surg, 2019. **270**(3): p. 484-492.
50. Ohene-Yeboah, M. and E. Adjei, *Breast cancer in Kumasi, Ghana.* Ghana Med J, 2012. **46**(1): p. 8-13.
51. Ramdas, Y., C.A. Benn, and M. van Heerden, *First Intraoperative Radiation Therapy Center in Africa: First 2 Years in Operation, Including COVID-19 Experiences.* JCO Glob Oncol, 2020. **6**: p. 1696-1703.
52. Rayne, S., et al., *Unraveling the South African Breast Cancer Story: The Relationship of Patients, Delay to Diagnosis, and Tumor Biology With Stage at Presentation in an Urban Setting.* Journal of Surgical Research, 2019. **235**: p. 181-189.

53. Roy, I. and E. Othieno, *Breast carcinoma in Uganda: microscopic study and receptor profile of 45 cases.* Arch Pathol Lab Med, 2011. **135**(2): p. 194-9.
54. Samaka, R.M. and S.F. Younes, *Androgen Receptor Expression in Breast Carcinoma of Egyptian Patients.* J Clin Diagn Res, 2016. **10**(11): p. Ec17-ec21.
55. Sawe, R.T., et al., *Tumor infiltrating leukocyte density is independent of tumor grade and molecular subtype in aggressive breast cancer of Western Kenya.* Tropical Medicine and Health, 2017. **45**(1): p. 19.
56. Schwartz, T., et al., *Expression of aldehyde dehydrogenase 1 as a marker of mammary stem cells in benign and malignant breast lesions of Ghanaian women.* Cancer, 2013. **119**(3): p. 488-494.
57. Sengal, A.T., et al., *Comparison of Receptor-Defined Breast Cancer Subtypes Between German and Sudanese Women: A Facility-Based Cohort Study.* Journal of Global Oncology, 2017(4): p. 1-12.
58. Seshie, B., et al., *A retrospective analysis of breast cancer subtype based on ER/PR and HER2 status in Ghanaian patients at the Korle Bu Teaching Hospital, Ghana.* BMC Clinical Pathology, 2015. **15**(1): p. 14.
59. Shaikh, A.J., et al., *Mammographic Breast Density and Breast Cancer Molecular Subtypes: The Kenyan-African Aspect.* BioMed Research International, 2018. **2018**: p. 6026315.
60. Stark, A., et al., *African ancestry and higher prevalence of triple-negative breast cancer.* Cancer, 2010. **116**(21): p. 4926-4932.
61. Sung, H., et al., *Breast cancer subtypes among Eastern-African–born black women and other black women in the United States.* Cancer, 2019. **125**(19): p. 3401-3411.
62. Tanimowo, M.O., et al., *Histopathological and immunohistochemical characteristics of breast carcinomas in Uyo, subtropical region of Africa.* Medical Journal of Zambia, 2019. **46**(2): p. 100-108.
63. Traore, B., et al., *Molecular profile of breast cancers in Guinean oncological settings.* Pan African Medical Journal, 2019. **33**.
64. CO., U., et al., *The Immunohistochemical Profile of Breast Cancer in Indigenous Women of Southeast Nigeria* Annals of Medical and Health Sciences Research, 2017. **7**(2): p. 83-87.
65. Uyisenga, J.P., et al., *Prevalence of Histological Characteristics of Breast Cancer in Rwanda in Relation to Age and Tumor Stages.* Horm Cancer, 2020. **11**(5-6): p. 240-249.
66. Youngblood, V.M., et al., *Outcomes and prognostic factors for women with breast cancer in Malawi.* Cancer Causes Control, 2020. **31**(4): p. 393-402.
67. Znati, K., et al., *Cancer du sein chez la femme jeune dans le Nord-Est du Maroc.* Gynécologie Obstétrique & Fertilité, 2012. **42**(3): p. 149-154.