Supplementary Materials for

"Efficacy and Safety of Huangqi Injection in the treatment of Hypertensive Nephropathy : A Systematic Review and Meta-Analysis"

ZhongChi Xu[1,†], LiChao Qian[2,†], RuGe Niu[1], Ying Yang[1], ChunLing Liu[1*], Xin Lin[1*]

1 *Jiangsu Provincial Hospital of Chinese Medicine, Affiliated Hospital of Nanjing University of Chinese Medicine, Nanjing, Jiangsu 210029, China*

2 *Nanjing Hospital of Chinese Medicine, Affiliated Hospital of Nanjing University of Chinese Medicine, Nanjing, Jiangsu 210001, China*

†Both authors contributed equally to this paper.

*Corresponding author:Xin Lin. *E-mail address*: linxin@njucm.edu.cn

Studies included: **8**
Participants included: **908**

Meta-analysis pooling of **Weighted Mean Differences**
using the **random-effects inverse-variance** model
with **DerSimonian-Laird** estimate of tau²

var1 (var2)	Effect	[95% Conf.	Interval]	% Weight
chen (2015)	**-0.440**	**-0.633**	**-0.247**	**11.51**
guo (2017)	**-0.150**	**-0.292**	**-0.008**	**13.69**
huang (2017)	**-0.300**	**-0.475**	**-0.125**	**12.29**
tang (2006)	**-0.143**	**-0.211**	**-0.074**	**16.58**
wu (2007)	**-0.250**	**-0.504**	**0.004**	**9.16**
yang (2015)	**-0.410**	**-0.724**	**-0.096**	**7.30**
zhao (2015)	**-0.450**	**-0.545**	**-0.355**	**15.67**
zhao (2017)	**-0.280**	**-0.420**	**-0.140**	**13.80**
Overall, DL	**-0.294**	**-0.405**	**-0.183**	**100.00**

Test of overall effect = **0**: **z** = **-5.193** p = **0.000**

Heterogeneity measures, calculated from the data
with Conf. Intervals based on **Gamma (random-effects)** distribution for Q

Measure	Value	df	p-value
Cochran's Q	**33.66**	**7**	**0.000**
		−[95% Conf.	Interval]−
H	**2.193**	**1.000**	**3.546**
I² (%)	**79.2%**	**0.0%**	**92.0%**

H = relative excess in Cochran's Q over its degrees-of-freedom
I² = proportion of total variation in effect estimate due to between-study heterogeneity (based on Q)

Heterogeneity variance estimates

Method	tau²
DL	**0.0181**

Supplementary Fig.1 Meta-analysis result of 24h UTP



Supplementary Fig.2 Funnel plot for publication bias of 24 h UTP

```
Egger's test for small-study effects:
Regress standard normal deviate of intervention
  effect estimate against its standard error

.
Number of studies =  8                                  Root MSE      =    2.204

------------------------------------------------------------------------------
     Std_Eff |      Coef.   Std. Err.      t    P>|t|     [95% Conf. Interval]
-------------+----------------------------------------------------------------
       slope |  -.1610176   .1123587    -1.43   0.202    -.4359494    .1139142
        bias |  -1.700827   1.762445    -0.97   0.372    -6.013375    2.611721
------------------------------------------------------------------------------

Test of H0: no small-study effects          P = 0.372
```

Supplementary Fig. 3 Egger's test for 24h UTP



Supplementary Fig. 4 Sensitivity analysis of 24h UTP

Meta-analysis regression

No of studies = **8**
tau^2 method **reml**
tau^2 estimate = **0**

Successive values of tau^2 differ by less than 10^-4 :convergence achieved

	Coef.	Std. Err.	z	P>|z|	[95% Conf.	Interval]
var9	**-.1732052**	**.1664156**	**-1.04**	**0.298**	**-.4993738**	**.1529635**
var10	**-.0100001**	**.1097002**	**-0.09**	**0.927**	**-.2250084**	**.2050083**
var11	**.2501514**	**.1972454**	**1.27**	**0.205**	**-.1364425**	**.6367453**
var12	**(dropped)**					
var13	**.2132052**	**.0654306**	**3.26**	**0.001**	**.0849634**	**.3414469**
_cons	**-.7201513**	**.3013447**	**-2.39**	**0.017**	**-1.310776**	**-.1295265**

Supplementary Fig. 5 Regression-analysis result of 24h UTP

Studies included: **8**
Participants included: **908**

Meta-analysis pooling of **Weighted Mean Differences**
using the **random-effects inverse-variance** model
with **DerSimonian-Laird** estimate of tau²

Subgroup and var1 (var2)	Effect	[95% Conf.	Interval]	% Weight
1				
chen (2015)	-0.440	-0.633	-0.247	11.51
zhao (2015)	-0.450	-0.545	-0.355	15.67
Subgroup, DL	-0.448	-0.533	-0.363	27.18
2				
guo (2017)	-0.150	-0.292	-0.008	13.69
huang (2017)	-0.300	-0.475	-0.125	12.29
tang (2006)	-0.143	-0.211	-0.074	16.58
wu (2007)	-0.250	-0.504	0.004	9.16
yang (2015)	-0.410	-0.724	-0.096	7.30
zhao (2017)	-0.280	-0.420	-0.140	13.80
Subgroup, DL	-0.212	-0.287	-0.138	72.82
Overall, DL	-0.294	-0.405	-0.183	100.00

Tests of subgroup effect size = **0**:
1 **z = -10.339 p = 0.000**
2 **z = -5.604 p = 0.000**
Overall **z = -5.193 p = 0.000**

Cochran's Q statistics for heterogeneity
(other heterogeneity measures are stored in matrices **r(ovstats)** and **r(bystats)**)

Measure	Value	df	p-value	I²
1	0.01	1	0.927	0.0%
2	7.36	5	0.195	32.1%
Overall	33.66	7	0.000	79.2%
Between	16.78	1	0.000	

Note: between-subgroup heterogeneity calculated using DL subgroup weights

.

Supplementary Fig. 6 Meta-analysis result of subgroups analysis of 24h UTP

Studies included: **4**
Participants included: **442**

Meta-analysis pooling of **Weighted Mean Differences**
using the **random-effects inverse-variance** model
with **DerSimonian-Laird** estimate of tau²

var1 (var2)	Effect	[95% Conf.	Interval]	% Weight
huang (2011)	**-12.340**	**-17.204**	**-7.476**	**31.51**
huang (2017)	**-14.100**	**-18.173**	**-10.027**	**33.51**
ji (2006)	**-27.730**	**-54.923**	**-0.537**	**4.46**
zhao (2017)	**-23.560**	**-28.800**	**-18.320**	**30.52**
Overall, DL	**-17.041**	**-23.141**	**-10.942**	**100.00**

Test of overall effect = **0**: **z** = **-5.476** p = **0.000**

Heterogeneity measures, calculated from the data
with Conf. Intervals based on **Gamma (random-effects)** distribution for Q

Measure	Value	df	p-value
Cochran's Q	**11.71**	**3**	**0.008**
		−[95% Conf.	Interval]−
H	**1.976**	**1.000**	**3.658**
I² (%)	**74.4%**	**0.0%**	**92.5%**

H = relative excess in Cochran's Q over its degrees-of-freedom
I² = proportion of total variation in effect estimate due to between-study heterogeneity (based on Q)

Heterogeneity variance estimates

Method	tau²
DL	**24.5828**

Supplementary Fig. 7 Meta-analysis result of mALB

Egger's test for small-study effects:
Regress standard normal deviate of intervention
effect estimate against its standard error

.
Number of studies = **4** Root MSE = **2.247**

Std_Eff	Coef.	Std. Err.	t	P>|t|	[95% Conf.	Interval]
slope	**-12.4539**	**7.227721**	**-1.72**	**0.227**	**-43.55227**	**18.64448**
bias	**-1.500431**	**2.653386**	**-0.57**	**0.629**	**-12.91703**	**9.916167**

Test of H0: no small-study effects P = **0.629**

.

Supplementary Fig. 8 Egger's test for mALB



Supplementary Fig. 9 Sensitivity analysis of mALB

Studies included: **4**
Participants included: **442**

Meta-analysis pooling of **Weighted Mean Differences**
using the **random-effects inverse-variance** model
with **DerSimonian-Laird** estimate of tau²

Subgroup and var1 (var2)	Effect	[95% Conf.	Interval]	% Weight
1				
huang (2011)	-12.340	-17.204	-7.476	31.51
huang (2017)	-14.100	-18.173	-10.027	33.51
Subgroup, DL	-13.375	-16.497	-10.252	65.02
2				
ji (2006)	-27.730	-54.923	-0.537	4.46
zhao (2017)	-23.560	-28.800	-18.320	30.52
Subgroup, DL	-23.709	-28.854	-18.564	34.98
Overall, DL	-17.041	-23.141	-10.942	100.00

Tests of subgroup effect size = **0**:
1 **z** = **-8.395** p = **0.000**
2 **z** = **-9.032** p = **0.000**
Overall **z** = **-5.476** p = **0.000**

Cochran's Q statistics for heterogeneity
(other heterogeneity measures are stored in matrices **r(ovstats)** and **r(bystats)**)

Measure	Value	df	p-value	I²
1	0.30	1	0.587	0.0%
2	0.09	1	0.768	0.0%
Overall	11.71	3	0.008	74.4%
Between	11.33	1	0.001	

Note: between-subgroup heterogeneity calculated using DL subgroup weights

.

Supplementary Fig. 10 Meta-analysis result of subgroups analysis of mALB

Studies included: **5**
Participants included: **531**

Meta-analysis pooling of **Weighted Mean Differences**
using the **random-effects inverse-variance** model
with **DerSimonian-Laird** estimate of tau²

var1 (var2)	Effect	[95% Conf.	Interval]	% Weight
guo (2017)	**-61.400**	**-69.848**	**-52.952**	**22.23**
zhao (2015)	**-61.300**	**-69.095**	**-53.505**	**22.29**
chen (2015)	**-8.790**	**-14.299**	**-3.281**	**22.46**
dong (2002)	**-29.100**	**-56.646**	**-1.554**	**19.01**
he (2004)	**-39.770**	**-89.240**	**9.700**	**14.01**
Overall, DL	**-40.392**	**-70.391**	**-10.393**	**100.00**

Test of overall effect = **0**: **z** = **-2.639** p = **0.008**

Heterogeneity measures, calculated from the data
with Conf. Intervals based on **Gamma (random-effects)** distribution for Q

Measure	Value	df	p-value
Cochran's Q	**168.14**	**4**	**0.000**
		−[95% Conf.	Interval]−
H	**6.484**	**1.164**	**12.243**
I² (%)	**97.6%**	**26.1%**	**99.3%**

H = relative excess in Cochran's Q over its degrees-of-freedom
I² = proportion of total variation in effect estimate due to between-study heterogeneity (based on Q)

Heterogeneity variance estimates

Method	tau²
DL	**1.0e+03**

Supplementary Fig. 11 Meta-analysis result of Scr

Egger's test for small-study effects:
Regress standard normal deviate of intervention
effect estimate against its standard error

.
Number of studies = **5** Root MSE = **7.206**

Std_Eff	Coef.	Std. Err.	t	P>|t|	[95% Conf.	Interval]
slope	**-22.66058**	**27.28169**	**-0.83**	**0.467**	**-109.4831**	**64.16195**
bias	**-2.980866**	**6.103437**	**-0.49**	**0.659**	**-22.40473**	**16.44299**

Test of H0: no small-study effects P = **0.659**

Supplementary Fig. 12 Egger’s test for Scr



Supplementary Fig. 13 Sensitivity analysis of Scr

Studies included: **4**
Participants included: **442**

Meta-analysis pooling of **Weighted Mean Differences**
using the **random-effects inverse-variance** model
with **DerSimonian-Laird** estimate of tau²

Subgroup and var1 (var2)	Effect	[95% Conf.	Interval]	% Weight
1				
huang (2011)	-12.340	-17.204	-7.476	31.51
huang (2017)	-14.100	-18.173	-10.027	33.51
Subgroup, DL	-13.375	-16.497	-10.252	65.02
2				
ji (2006)	-27.730	-54.923	-0.537	4.46
zhao (2017)	-23.560	-28.800	-18.320	30.52
Subgroup, DL	-23.709	-28.854	-18.564	34.98
Overall, DL	-17.041	-23.141	-10.942	100.00

Tests of subgroup effect size = **0**:
1 **z = -8.395 p = 0.000**
2 **z = -9.032 p = 0.000**
Overall **z = -5.476 p = 0.000**

Cochran's Q statistics for heterogeneity
(other heterogeneity measures are stored in matrices **r(ovstats)** and **r(bystats)**)

Measure	Value	df	p-value	I²
1	0.30	1	0.587	0.0%
2	0.09	1	0.768	0.0%
Overall	11.71	3	0.008	74.4%
Between	11.33	1	0.001	

Note: between-subgroup heterogeneity calculated using DL subgroup weights

.

Supplementary Fig. 14 Meta-analysis result of subgroups analysis of Scr

Studies included: **7**
Participants included: **810**

Meta-analysis pooling of **Weighted Mean Differences**
using the **random-effects inverse-variance** model
with **DerSimonian-Laird** estimate of tau²

var1 (var2)	Effect	[95% Conf.	Interval]	% Weight
chen (2015)	**-11.900**	**-17.343**	**-6.457**	**13.33**
guo (2017)	**-18.890**	**-20.046**	**-17.734**	**15.55**
han (2011)	**-4.500**	**-10.878**	**1.878**	**12.62**
song (2019)	**-9.580**	**-10.948**	**-8.212**	**15.50**
yang (2015)	**-4.950**	**-9.244**	**-0.656**	**14.12**
zhao (2017)	**-13.600**	**-17.134**	**-10.066**	**14.59**
zhaoYJ (2015)	**-1.720**	**-5.739**	**2.299**	**14.30**
Overall, DL	**-9.504**	**-14.642**	**-4.366**	**100.00**

Test of overall effect = **0**: **z** = **-3.625** p = **0.000**

Heterogeneity measures, calculated from the data
with Conf. Intervals based on **Gamma (random-effects)** distribution for Q

Measure	Value	df	p-value
Cochran's Q	**170.72**	**6**	**0.000**
		[95% Conf.	Interval]
H	**5.334**	**1.060**	**9.919**
I² (%)	**96.5%**	**11.0%**	**99.0%**

H = relative excess in Cochran's Q over its degrees-of-freedom
I² = proportion of total variation in effect estimate due to between-study heterogeneity (based on Q)

Heterogeneity variance estimates

Method	tau²
DL	**43.8563**

Supplementary Fig. 15 Meta-analysis result of SBP



Supplementary Fig. 16 Sensitivity analysis of SBP

```
Egger's test for small-study effects:
Regress standard normal deviate of intervention
  effect estimate against its standard error

.
Number of studies =  7                                  Root MSE      =   4.826
```

Std_Eff	Coef.	Std. Err.	t	P>|t|	[95% Conf.	Interval]
slope	-18.20857	3.500145	-5.20	0.003	-27.20598	-9.211162
bias	4.906297	3.215138	1.53	0.188	-3.358477	13.17107

```
Test of H0: no small-study effects          P = 0.188
```

Supplementary Fig. 17 Egger's test for SBP

Studies included: **7**
Participants included: **810**

Meta-analysis pooling of **Weighted Mean Differences**
using the **random-effects inverse-variance** model
with **DerSimonian-Laird** estimate of tau²

Subgroup and var1 (var2)	Effect	[95% Conf.	Interval]	% Weight
1				
chen (2015)	-11.900	-17.343	-6.457	13.33
guo (2017)	-18.890	-20.046	-17.734	15.55
song (2019)	-9.580	-10.948	-8.212	15.50
zhao (2017)	-13.600	-17.134	-10.066	14.59
Subgroup, DL	-13.570	-19.506	-7.633	58.96
2				
han (2011)	-4.500	-10.878	1.878	12.62
yang (2015)	-4.950	-9.244	-0.656	14.12
zhaoYJ (2015)	-1.720	-5.739	2.299	14.30
Subgroup, DL	-3.451	-6.116	-0.785	41.04
Overall, DL	-9.504	-14.642	-4.366	100.00

Tests of subgroup effect size = **0**:
1 **z** = **-4.480** p = **0.000**
2 **z** = **-2.537** p = **0.011**
Overall **z** = **-3.625** p = **0.000**

Cochran's Q statistics for heterogeneity
(other heterogeneity measures are stored in matrices **r(ovstats)** and **r(bystats)**)

Measure	Value	df	p-value	I²
1	105.57	3	0.000	97.2%
2	1.28	2	0.526	0.0%
Overall	170.72	6	0.000	96.5%
Between	9.29	1	0.002	

Note: between-subgroup heterogeneity calculated using DL subgroup weights

.

Supplementary Fig. 18 Meta-analysis result of subgroups analysis of SBP

Studies included: **7**
Participants included: **810**

Meta-analysis pooling of **Weighted Mean Differences**
using the **random-effects inverse-variance** model
with **DerSimonian-Laird** estimate of tau²

var1 (var2)	Effect	[95% Conf.	Interval]	% Weight
chen (2015)	**-4.830**	**-8.131**	**-1.529**	**12.01**
guo (2017)	**-7.310**	**-9.707**	**-4.913**	**17.03**
han (2011)	**-3.750**	**-7.532**	**0.032**	**10.05**
song (2019)	**-5.830**	**-8.060**	**-3.600**	**18.17**
yang (2015)	**-2.770**	**-4.573**	**-0.967**	**21.36**
zhao (2017)	**-4.040**	**-6.563**	**-1.517**	**16.21**
zhaoYJ (2015)	**-1.560**	**-7.388**	**4.268**	**5.17**
Overall, DL	**-4.588**	**-6.036**	**-3.140**	**100.00**

Test of overall effect = **0**: **z** = **-6.210** p = **0.000**

Heterogeneity measures, calculated from the data
with Conf. Intervals based on **Gamma (random-effects)** distribution for Q

Measure	Value	df	p-value
Cochran's Q	**11.47**	**6**	**0.075**
		[95% Conf.	Interval]
H	**1.383**	**1.000**	**2.171**
I² (%)	**47.7%**	**0.0%**	**78.8%**

H = relative excess in Cochran's Q over its degrees-of-freedom
I² = proportion of total variation in effect estimate due to between-study heterogeneity (based on Q)

Heterogeneity variance estimates

Method	tau²
DL	**1.7097**

Supplementary Fig. 19 Meta-analysis result of DBP

Egger's test for small-study effects:
Regress standard normal deviate of intervention
effect estimate against its standard error

.
Number of studies = **7** Root MSE = **1.512**

Std_Eff	Coef.	Std. Err.	t	P>|t|	[95% Conf.	Interval]
slope	**-4.852767**	**2.464015**	**-1.97**	**0.106**	**-11.18672**	**1.481186**
bias	**.2456837**	**1.853388**	**0.13**	**0.900**	**-4.518601**	**5.009968**

Test of H0: no small-study effects P = **0.900**

Supplementary Fig. 20 Egger's test for DBP



Supplementary Fig. 21 Sensitivity analysis of DBP

Studies included: **7**
Participants included: **810**

Meta-analysis pooling of **Weighted Mean Differences**
using the **random-effects inverse-variance** model
with **DerSimonian-Laird** estimate of tau²

Subgroup and var1 (var2)	Effect	[95% Conf.	Interval]	% Weight
1				
chen (2015)	-4.830	-8.131	-1.529	12.01
guo (2017)	-7.310	-9.707	-4.913	17.03
song (2019)	-5.830	-8.060	-3.600	18.17
zhao (2017)	-4.040	-6.563	-1.517	16.21
Subgroup, DL	-5.623	-7.033	-4.213	63.42
2				
han (2011)	-3.750	-7.532	0.032	10.05
yang (2015)	-2.770	-4.573	-0.967	21.36
zhaoYJ (2015)	-1.560	-7.388	4.268	5.17
Subgroup, DL	-2.851	-4.418	-1.283	36.58
Overall, DL	-4.588	-6.036	-3.140	100.00

Tests of subgroup effect size = **0**:
1 **z** = **-7.815** p = **0.000**
2 **z** = **-3.565** p = **0.000**
Overall **z** = **-6.210** p = **0.000**

Cochran's Q statistics for heterogeneity
(other heterogeneity measures are stored in matrices **r(ovstats)** and **r(bystats)**)

Measure	Value	df	p-value	I²
1	3.67	3	0.300	18.2%
2	0.41	2	0.813	0.0%
Overall	11.47	6	0.075	47.7%
Between	6.64	1	0.010	

Note: between-subgroup heterogeneity calculated using DL subgroup weights

.

Supplementary Fig. 22 Meta-analysis result of subgroups analysis of DBP

Studies included: **4**
Participants included: **422**

Meta-analysis pooling of **Weighted Mean Differences**
using the **random-effects inverse-variance** model
with **DerSimonian-Laird** estimate of tau²

var1 (var2)	Effect	[95% Conf.	Interval]	% Weight
guo (2017)	**-0.860**	**-1.076**	**-0.644**	**25.61**
huang (2011)	**-0.450**	**-0.626**	**-0.274**	**27.71**
huang (2017)	**-0.900**	**-1.216**	**-0.584**	**20.28**
zhao (2015)	**-0.830**	**-1.031**	**-0.629**	**26.40**
Overall, DL	**-0.747**	**-0.975**	**-0.518**	**100.00**

Test of overall effect = **0**: **z** = **-6.406** p = **0.000**

Heterogeneity measures, calculated from the data
with Conf. Intervals based on **Gamma (random-effects)** distribution for Q

Measure	Value	df	p-value
Cochran's Q	**12.98**	**3**	**0.005**
		[95% Conf.	Interval]
H	**2.080**	**1.000**	**3.720**
I² (%)	**76.9%**	**0.0%**	**92.8%**

H = relative excess in Cochran's Q over its degrees-of-freedom
I² = proportion of total variation in effect estimate due to between-study heterogeneity (based on Q)

Heterogeneity variance estimates

Method	tau²
DL	**0.0409**

.

Supplementary Fig. 23 Meta-analysis result of Cys-c

Egger's test for small-study effects:
Regress standard normal deviate of intervention
effect estimate against its standard error

.
Number of studies = **4** Root MSE = **1.904**

Std_Eff	Coef.	Std. Err.	t	P>|t|	[95% Conf.	Interval]
slope	**-.0542323**	**.5293197**	**-0.10**	**0.928**	**-2.331711**	**2.223246**
bias	**-6.133252**	**4.87931**	**-1.26**	**0.336**	**-27.12723**	**14.86072**

Test of H0: no small-study effects P = **0.336**

Supplementary Fig.24 Egger's test for Cys-c



Supplementary Fig. 25 Sensitivity analysis of Cys-c

Studies included: **3**
Participants included: **332**

Meta-analysis pooling of **Weighted Mean Differences**
using the **random-effects inverse-variance** model
with **DerSimonian-Laird** estimate of tau²

var1 (var2)	Effect	[95% Conf.	Interval]	% Weight
guo (2017)	**-0.860**	**-1.076**	**-0.644**	**38.21**
huang (2017)	**-0.900**	**-1.216**	**-0.584**	**17.78**
zhao (2015)	**-0.830**	**-1.031**	**-0.629**	**44.01**
Overall, DL	**-0.854**	**-0.987**	**-0.721**	**100.00**

Test of overall effect = **0**: **z** = **-12.553** p = **0.000**

Heterogeneity measures, calculated from the data
with Conf. Intervals based on **Gamma (random-effects)** distribution for Q

Measure	Value	df	p-value
Cochran's Q	**0.14**	**2**	**0.933**
		−[95% Conf.	Interval]−
H	**0.264**	**1.000**	**0.881**
I² (%)	**0.0%**	**0.0%**	**0.0%**

H = relative excess in Cochran's Q over its degrees-of-freedom
I² = proportion of total variation in effect estimate due to between-study heterogeneity (based on Q)

Heterogeneity variance estimates

Method	tau²
DL	**0.0000**

.

Supplementary Fig. 26 Meta-analysis result of Cys-c after exclude

Studies included: **5**
Participants included: **483**

Meta-analysis pooling of **Weighted Mean Differences**
using the **random-effects inverse-variance** model
with **DerSimonian-Laird** estimate of tau²

var1 (var2)	Effect	[95% Conf.	Interval]	% Weight
dong (2002)	**-2.140**	**-3.588**	**-0.692**	**21.56**
guo (2017)	**-5.400**	**-6.909**	**-3.891**	**21.35**
he (2004)	**-8.600**	**-12.639**	**-4.561**	**12.19**
zhao (2015)	**-5.100**	**-6.447**	**-3.753**	**21.89**
zhao (2017)	**-1.630**	**-2.600**	**-0.660**	**23.00**
Overall, DL	**-4.155**	**-6.152**	**-2.157**	**100.00**

Test of overall effect = **0**: **z** = **-4.077** p = **0.000**

Heterogeneity measures, calculated from the data
with Conf. Intervals based on **Gamma (random-effects)** distribution for Q

Measure	Value	df	p-value
Cochran's Q	**34.75**	**4**	**0.000**
		−[95% Conf.	Interval]−
H	**2.948**	**1.000**	**5.142**
I² (%)	**88.5%**	**0.0%**	**96.2%**

H = relative excess in Cochran's Q over its degrees-of-freedom
I² = proportion of total variation in effect estimate due to between-study heterogeneity (based on Q)

Heterogeneity variance estimates

Method	tau²
DL	**4.2712**

Supplementary Fig. 27 Meta-analysis result of BUN

Egger's test for small-study effects:
Regress standard normal deviate of intervention
effect estimate against its standard error

.
Number of studies = **5** Root MSE = **2.44**

Std_Eff	Coef.	Std. Err.	t	P>|t|	[95% Conf.	Interval]
slope	**.2412832**	**2.220401**	**0.11**	**0.920**	**-6.825022**	**7.307589**
bias	**-5.27867**	**3.132881**	**-1.68**	**0.191**	**-15.2489**	**4.691555**

Test of H0: no small-study effects P = **0.191**

Supplementary Fig. 28 Egger's test for BUN



Supplementary Fig. 29 Sensitivity analysis of BUN