Supplementary Online Content

This supplement contains the following items:

eFigure 1 Inconsistency assessment of network meta-analysis
eTable 1 Search strategy
eTable 2 Extracted data of included studies for the network meta-analysis
eTable 3 Characteristics of included studies
eTable 4 Risk of bias assessment of included studies
eTable 5 Mixed comparisons
eTable 6 GRADE summary of findings

eTable 1. Search strategies

The search strategy used in PubMed:

#1 "chronic prostatitis"[Title/Abstract]
#2 "chronic pelvic pain syndrome"[Title/Abstract]
#3 "nonbacterial prostatitis"[Title/Abstract]
#4 "chronic prostatitis/chronic pelvic pain syndrome"[Title/Abstract]
#5 "chronic prostatitis-chronic pelvic pain syndrome"[Title/Abstract]
#6 "CP/CPPS"[Title/Abstract]
#7 "CP-CPPS"[Title/Abstract]
#8 #1 OR #2 OR #3 OR #4 OR #5 OR #6 OR #7
#9 "Randomized Controlled Trial" [Publication Type]
#10 "Randomized Controlled Trials as Topic"[Mesh]
#11 "Pragmatic Clinical Trial" [Publication Type]
#12 "Pragmatic Clinical Trials as Topic"[Mesh]
#13 "Intention to Treat Analysis"[Mesh]
#14 "random allocation"[MeSH Terms]
#15 random*[Title/Abstract]
#16 #9 OR #10 OR #11 OR #12 OR #13 OR #14 OR #15
#17 #8 AND #16

The strategy used in EMBASE:

#1 'chronic prostatitis'/exp
#2 'chronic prostatitis':ti,ab
#3 'chronic pelvic pain syndrome':ti,ab
#4 'nonbacterial prostatitis':ti,ab
#5 'chronic prostatitis/chronic pelvic pain syndrome':ti,ab
#6 'chronic prostatitis-chronic pelvic pain syndrome':ti,ab
#7 'CP/CPPS':ti,ab
#8 'CP-CPPS':ti,ab
#9 OR/1-8
#10 'randomized controlled trial'/exp
#11 'randomized controlled trial (topic)'/exp
#12 random*:ti,ab
#13 OR/10-12
#14 #9 AND #13

The strategy used in Cochrane Library:

#1 [mh prostatitis]
#2 prostatit*:ti,ab
#3 prostatodyn*:ti,ab
#4 #1 or #2 or #3
#5 [mh "pelvic pain"]

#6 pelvic near/3 pain:ti,ab
#7 CPP*:ti,ab
#8 #5 or #6 or #7
#9 #4 and #8
#10 [mh "asymptomatic diseases"]
#11 asymptom*:ti,ab
#12 nonsymptom*:ti,ab
#13 non near/3 symptom*:ti,ab
#14 abacterial*:ti,ab
#15 nonbacterial*:ti,ab
#16 non near/3 bacterial*:ti,ab
#17 aseptic*:ti,ab
#18 nonseptic*:ti,ab
#19 non near/3 septic*:ti,ab
#20 #10 or #11 or #12 or #13 or #14 or #15 or #16 or #17 or #18 or #19
#21 #4 and #20
#22 prostatit* and (pelvic near/3 pain or CP CPP*)
#23 #9 or #21 or #22

The strategy used in Web of Science:

1. TS=(prostatit* NEAR/3 (chronic* or pelvic* or pain*))
2. TS=(chronic pelvic pain*)
3. TS=(CPP*)
4. TS=(nonbacterial prostatit* or abacterial prostatit*)
5. #1 OR #2 OR #3 OR #4
6. TS=(clinical trial* or controlled trial* or random* or research design or comparative stud* or evaluation stud* or prospective stud* or placebo*)
7. #5 AND #6
8. All languages

eTable 2. Extracted data of included studies for the network meta-analysis

Study ID	Treatment	N	NIH-CPSI Total scores		NIH-CPSI Pain scores		NIH-CPSI Urinary scores		NIH-CPSI Quality of life scores		Adverse events
			Mean	SD	Mean	SD	Mean	SD	Mean	SD	N*
Ye 2008	Tamsuloxin	42	NA	NA	-4.8	1.7	-2.5	2.0	-3.2	1.3	0
	Tamsuloxin + Levofloxacin	42	NA	NA	-7.1	1.8	-5.6	1.7	-5.3	1.3	0
	Levofloxacin	21	NA	NA	-3.4	1.7	-1.8	1.9	-1.6	1.3	0
Nickel 2005	Pentosan polysulfate sodium	51	-5.9	7.4	-2.7	3.9	-1.2	2.5	-2.0	2.3	11
	Placebo	49	-3.2	7.5	-1.7	3.9	-0.5	2.5	-1.0	2.3	4
Alexander 2004	Ciprofloxacin	49	-6.2	7.3	-3.0	4.6	-1.3	2.1	-1.9	2.4	19
	Tamsulosin	49	-4.4	6.3	-1.8	3.7	-1.1	1.8	-1.3	2.0	20
	Ciprofloxacin + Tamsulosin	49	-4.1	6.1	-1.6	2.9	-0.5	2.0	-1.3	1.9	17
	Placebo	49	-3.4	5.0	-2.3	3.7	-0.9	2.0	-1.2	1.5	20
Cai 2017	Pollen extract	36	-13.4	2.0	-4.8	0.3	NA	NA	NA	NA	1
	Bromelain	34	-3.1	2.2	-2.1	0.7	NA	NA	NA	NA	2
Shoskes 1999	Quercetin	15	-8.0	4.3	-4.1	2.3	-1.2	1.8	-3.1	2.0	2
	Placebo	15	-1.4	4.7	-0.4	2.1	-0.2	1.8	-0.9	1.9	1
Chen 2011	Tamsulosin	50	-7.5	1.9	-4.4	1.3	-2.0	1.1	-1.3	1.0	13
	Placebo	50	-4	2.3	-1.7	1.3	-1.1	0.7	-1.2	0.8	2
Wagenlehner	Pollen extract	70	-7.7	5.8	-4.5	3.5	-1.0	1.6	-2.2	2.2	9

2009	Placebo	69	-5.2	5.8	-2.9	3.5	-0.9	1.6	-1.4	2.2	10
Zhao 2009	Celecoxib	32	-8.0	3.0	-5.5	2.6	-0.7	1.7	-2.2	1.8	2
	Placebo	32	-4.8	2.9	-2.9	2.6	-0.8	1.7	-1.0	1.8	1
Nickel 2011a	Dutasteride	341	-5.2	6.6	-3.9	3.6	-0.4	2.4	-0.9	2.5	NA
	Placebo	343	-2.7	7.2	-2.9	4.3	0.3	2.6	-0.2	2.5	NA
Pontari 2010	Pregabalin	218	-6.5	5.2	-3.2	2.8	-1.2	1.7	-2.0	1.8	NA
	Placebo	106	-4.3	5.4	-2.3	2.9	-0.7	1.7	-1.5	1.9	NA
Nickel 2008	Alfuzosin	138	-7.1	9.0	-3.3	4.5	-1.2	2.6	-1.2	1.5	39
	Placebo	134	-6.5	8.5	-3.0	4.4	-1.0	2.6	-1.2	1.5	38
Kim 2011a	Ciprofloxacin + solifenacin	49	-7.7	4.5	-4.0	2.2	-1.0	1.3	-2.7	2.0	NA
	Ciprofloxacin	47	-11.6	3.6	-5.5	2.0	-2.8	1.6	-3.3	1.7	NA
Zhao 2019	Tamsulosin + dapoxetine	83	-14.3	4.7	NA	NA	-3.5	1.9	-3.5	1.2	3
	Tamsulosin	168	-16.9	3.6	NA	NA	-3.9	1.4	-5.3	1.2	14
Tuğcu 2007	Doxazosin	30	-12.4	1.1	-5.2	1.0	-3.0	0.9	-4.2	0.7	12
	Doxazosin + ibuprofen + thiocolchicoside	30	-12.7	0.9	-5.4	0.9	-3.5	0.9	-3.8	0.6	17
	Placebo	23	-1.0	0.8	-0.6	0.5	-0.2	0.9	-0.2	0.7	7

Kaplan 2004	Finasteride	32	-6.3	4.1	-2.2	0.6	NA	NA	NA	NA	2
	Saw palmetto	32	0.6	4.5	-0.4	0.6	NA	NA	NA	NA	3
Nickel 2004	Finasteride	33	-2.9	1.0	-2.0	0.5	-0.6	0.3	-1.1	0.4	5
	Placebo	31	-0.8	1.0	-0.1	0.6	0	0.3	-0.8	0.4	7
Macchione 2019	Pollen extract	40	-17.3	3.1	NA	NA	NA	NA	NA	NA	0
	Serenoa repens	43	-13.6	4.8	NA	NA	NA	NA	NA	NA	0
Cheah 2003	Terazosin	50	-14.3	5.4	-6.5	3.6	-3.1	2.2	-4.7	2.0	18
	Placebo	50	-10.2	7.5	-4.8	4.3	-1.8	2.3	-3.4	2.5	9
Kim 2011b	Tamsulosin	40	-15	3.3	-7.4	2.3	-2.6	1.2	-5.1	1.4	
	Tamsulosin + diclofenac	32	-16.2	3.5	-8.2	2.1	-2.5	1.3	-5.1	1.6	NA
	Tamsulosin + ciprofloxacin	28	-15.0	3.3	-7.2	2.4	-2.0	0.9	-5.7	1.3	NA
Nickel 2003	Levofloxacin	45	-5.4	5.7	-1.7	3.1	-1.2	1.9	-1.0	1.1	9
	Placebo	35	-2.9	5.5	-1.8	2.8	-0.2	1.8	-0.7	1.	6
Wang 2016	Levofloxacin	38	-10.9	3.9	NA	NA	NA	NA	NA	NA	NA
	Terazosin	38	-5.0	3.9	NA	NA	NA	NA	NA	NA	
	Levofloxacin + terazosin	39	-11.4	4.6	NA	NA	NA	NA	NA	NA	NA
Maurizi 2019	Pollen extract	27	-13.5	1.1	NA	NA	NA	NA	NA	NA	0
	Quercetin	27	-11.1	1.1	NA	NA	NA	NA	NA	NA	0

Nickel 2011b	Silodosin 4 mg	52	-12.1	9.3	-5.9	4.5	-2.2	2.7	-4.1	3.1	3
	Silodosin 8 mg	45	-10.2	8.8	-4.9	5.0	-1.8	2.4	-3.5	3.0	6
	Placebo	54	-8.5	7.2	-4.5	3.6	-1.3	3.0	-2.7	2.5	1
Morgia 2010	Profluss	51	-6.8	2.5	NA	NA	NA	NA	NA	NA	5
	Serenoa repens	51	-11.4	5.3	NA	NA	NA	NA	NA	NA	4
Cai 2014	Pollen extract	41	-12.1	1.4	-4.4	0.5	NA	NA	NA	NA	1
	Ibuprofen	46	-6.0	1.8	-2.2	0.5	NA	NA	NA	NA	7

ID: identification; N: number of patients; QoL: quality of life; SD: standard deviation; NA: not available; N: number of adverse events*

eTable 3. Study characteristics of the included trials

Study ID	No. arm (included/total)	Fund	No. randomized	Mean age (years)	Dose (mg daily)	Baseline (Pain/Total/Urinary/QoL)	Aftertreatment (Pain/Total/Urinary/QoL)	Length of trial	Recruitment place	Multi/ single center	No. adverse event
Ye 2008	3/5	Yes	105	NR	tamsulosin 0.2mg, levofloxacin 200mg	IIIA tamsulosin group: 10.57±3.54, NR, 4.62±2.67, 9.33±1.91; IIIA tamsulosin plus levofloxacin group: 16.19±2.79, NR, 8.71±1.65, 11.38±1.16; IIIA levofloxacin group: 11.48±2.84, NR, 4.57±3.03, 9.38±2.06; IIIB tamsulosin group: 11.0±2.70, NR, 4.43±3.79, 9.43±1.75; IIIB tamsulosin plus levofloxacin group: 11.24±2.96, NR, 6.14±2.10, 9.57±1.47	IIIA tamsulosin group: 5.86±2.99, NR, 2.24±2.10, 6.10±2.47; IIIA tamsulosin plus levofloxacin group: 4.90±1.58, NR, 1.81±1.54, 5.67±1.71; IIIA levofloxacin group: 8.05±2.16, NR, 2.76±2.05, 7.81±2.06; IIIB tamsulosin group: 6.19±1.54, NR, 1.86±2.20, 6.38±1.86; IIIB tamsulosin plus levofloxacin group: 4.10±2.72, NR, 1.81±1.86, 4.71±2.24	13 W	Asia	single	No
Nickel 2005	2/2	NR	100	pentosan polysulfate sodium (PPS): 40.8 (21-59); placebo group: 37.5 (25-55)	PPS 900 mg; placebo 900mg	PPS group: 12.4±0.56, 27.1±1.10, 6.0±0.44, 8.9±0.36; placebo group: 12.3±0.58, 25.8±1.13, 5.1±0.46, 8.5±0.38	PPS group: 9.7±NR, 21.2±NR, 4.8±NR, 6.9±NR; placebo group: 10.6±NR, 22.6±NR, 4.6±NR, 7.5±NR	16 W	North America	multi	PPS: 11; placebo: 4[A]

Study ID	No. arms (include	Fund	No. rando	Mean age (years)	Dose (mg daily)	Baseline (Pain/Total/Urinary/QoL)	Aftertreatment (Pain/Total/Urinary/QoL)	Length of trial	Recruitment place	Multi/ single	No. adverse event

	d/total)		mized							center	
Alexander 2004	4/4	Yes	196	Placebo group: 42.6±12.0; ciprofloxacin group: 45.9±11.7; tamsulosin group: 45.3±9.7; combination group: 44.5±11.4	ciprofloxacin 1000mg; tamsulosin 0.4 mg; combination of both drugs	placebo group: 12.2±3.0, 25.0±5.1, 4.5±2.5, 8.3±1.9; ciprofloxacin group: 11.7±3.1, 24.2±6.2, 4.8±2.8, 7.7±2.2; tamsulosin group: 11.3±3.6, 24.6±6.2, 5.1±2.9, 8.2±2.2; combination group: 12.0±3.4, 25.3±6.1, 5.0±2.4, 8.2±2.3	NR	6 W	North America	multi	placebo: 20 ciprofloxacin: 19 tamsulosin: 20 combination: 17[B]
Cai 2017	2/2	NR	70	Deprox 500® group: 32.4±4.3; Bromelain group: 32.8±4.9	Deprox 500® 2 tablets; bromelain 80mg	Deprox 500® group: 11.3±2.1, 25.1±2.1, NR, NR; bromelain group: 10.7±2.5, 25.6±2.9, NR, NR	Deprox 500® group: 6.7±1.9, 11.7±3.2, NR, NR; bromelain group: 8.1±2.3, 22.5±3.7, NR, NR	12 W	Europe	single	Deprox 500®: 1; Bromelain: 2 (mild effects: nausea)
Shoskes 1999	2/2	NR	30	quercetin group: 46.2±4.0; placebo group: 43.5±3.7	quercetin capsules 1000 mg; placebo 1000mg	quercetin group: 10.3±0.86, 21.0±1.8, 2.7±0.74, 8.0±0.84; placebo group: 9.4±0.63, 20.2±1.1, 3.2±0.76, 7.7±0.74	quercetin group: 6.2±1.0, 13.0±1.7, 1.5±0.5, 4.9±0.69; placebo group: 9.0±0.88, 18.8±1.9, 3.0±0.75, 6.8±0.8	4 W	North America	single	quercetin: 2; placebo: 1[C]

Study ID	No. arms (included/total)	Fund	No. randomized	Mean age (years)	Dose (mg daily)	Baseline (Pain/Total/Urinary/QoL)	Aftertreatment (Pain/Total/Urinary/QoL)	Length of trial	Recruitment place	Multi/ single center	No. adverse event

Study ID	No. arms (included/total)	Fund	No. randomized	Mean age (years)	Dose (mg daily)	Baseline (Pain/Total/Urinary/QoL)	Aftertreatment (Pain/Total/Urinary/QoL)	Length of trial	Recruitment place	Multi/ single center	No. adverse event
Chen 2011	2/2	NR	100	tamsulosin group: 35.3±6.8; placebo group: 33.3±7.2	tamsulosin 0.2mg; placebo 0.2mg	tamsulosin group: 12.4±2.2, 23.3±6.2, 5.4±2.0, 5.0±1.8; placebo group: 11.8±3.0, 22.5±5.6, 4.8±2.5, 5.2±2.1	tamsulosin group: NR, 15.7±4.0, NR, NR; placebo group: NR, 18.3±3.9, NR, NR	24 W	Asia	multi	tamsulosin: 13; placebo: 2[D]
Wagenlehner 2009	2/2	Yes	139	pollen extract group: 39.7±7.2 placebo group: 39.3±9.1	pollen extract 6 capsules; placebo 6 capsules	pollen extract group: 10.0±2.4, 19.3±5.1, 2.8±2.3, 6.5±2.5 placebo group: 10.2±2.6, 20.3±5.2, 3.5±2.5, 6.7±2.2		12 W	Europe	multi	pollen extract: 9; placebo: 10[E]
Zhao 2009	2/2	NR	64	NR	celecoxib 200mg; placebo 200mg	celecoxib group: NR, 23.91±5.27, NR, NR; placebo group: NR, 24.25±5.09, NR, NR	celecoxib group: NR, 15.88±2.51, NR, NR; placebo group: NR, 19.50±2.50, NR, NR	6 W	Asia	single	celecoxib: 2; placebo: 1[F]
Nickel 2011a	2/2	Yes	5379	NR	dutasteride 0.5mg; placebo 0.5mg	dutasteride group: 7.02±2.18, 15.03±5.39, 3.60±2.43, 4.47±2.40; placebo group: 6.98±2.09, 15.34±5.07, 3.72±2.41, 4.63±2.24	NR	192 W	North America	multi	dutasteride: 102; placebo: 150[G]

Pontari 2010	2/2	Yes	324	pregabalin group: 48.0±13.0; placebo group: 45.2±12.2	pregabalinand placebo (150mg/d for 2 W, 300mg/d for 2 W, 600mg/d for 2 W)	pregabalin group: 12.3±3.0, 26.2±5.6, 4.9±2.7, 8.9±2.0; placebo group: 12.4±3.1, 25.9±6.1, 4.7±2.7, 8.9±2.0	pregabalin group: 9.1±4.6, 19.7±8.5, 3.7±2.6, 6.9±2.9; placebo group: 10.1±4.7, 21.6±8.9, 4.0±2.7, 7.4±3.1	6 W	North America	multi	NR
Nickel 2008	2/2	Yes	272	alfuzosin group: 40.1±11.4 placebo group: 40.1±12.3	alfuzosin 10mg; placebo 10mg	alfuzosin group: 11.1±3.3, 23.8±6.3, 4.5±2.8, 4.5±1.1; placebo group: 11.5±3.4, 25.1±5.9, 4.9±2.9, 4.7±0.9	NR	12 W	North America and Asia	multi	alfuzosin: 39; placebo: 38[H]
Kim 2011a	2/2	NR	96	NR	ciprofloxacin 1000mg and solifenacin 5mg; ciprofloxacin 1000mg	ciprofloxacin group: 9.3±3.2, 19.8±6.4, 2.9±2.2, 7.8±3.0; ciprofloxacin and solifenacin group: 9.4±3.3, 22.4±6.0, 4.9±2.6, 8.4±2.7	ciprofloxacin group: 5.3±3.6, 12.1±7.4, 1.9±1.6, 5.1±3.3; ciprofloxacin and solifenacin group: 3.9±2.9, 10.8±5.4, 2.1±1.8, 5.1±1.8	8 W	Asia	multi	NR

Study ID	**No. arms (included/total)**	**Fund**	**No. randomized**	**Mean age (years)**	**Dose (mg daily)**	**Baseline (Pain/Total/Urinary/QoL)**	**Aftertreatment (Pain/Total/Urinary/QoL)**	**Length of trial**	**Recruitment place**	**Multi/ single center**	**No. adverse event**

Zhao 2019	2/2	Yes	251	tamsulosin group: 31.2±5.96; tamsulosin and dapoxetine group: 33.9±6.56	tamsulosin 0.2mg; dapoxetine 30mg before sexual activity	tamsulosin group: 10.59±4.66, 26.15±7.76, 6.37±3.15, 9.19±1.78; tamsulosin and dapoxetine group: 9.86±3.44, 24.29±5.92, 5.76±2.18, 8.64±1.86	tamsulosin group: NR, 11.81±6.10, 2.89±2.42, 5.74±1.99; tamsulosin and dapoxetine group: NR, 7.44±4.75, 1.82±1.40, 3.38±1.79	12 W	Asia	multi	tamsulosin: 3; tamsulosin and dapoxetine: 14[I]
Tuğcu 2007	3/3	NR	83	29.1±5.2	doxazosin 4mg; combination (doxazosin 4mg ibuprofen 400mg, thiocolchicoside 12mg);	doxazosin group: 9.9±1.6, 23.1±1.8, 5.2±1.0, 8.0±0.9; combination group: 8.8±1.4, 21.9±1.5, 6.9±1.4, 6.2±0.9 placebo group: 9.2±0.9, 22.9±1.2, 6.6±0.8, 7.1±1.1	doxazosin group: 4.7±1.2, 10.7±1.3, 2.2±0.8, 3.8±1.1; combination group: 3.4±0.7, 9.2±1.0, 3.4±0.8, 2.4±0.9; placebo group: 8.6±0.8, 21.9±1.2, 6.4±1.0, 6.9±1.1	24 W	Asia	multi	doxazosin: 12; combination: 17; placebo: 7[J]
Kaplan 2004	2/2	NR	64	finasteride group: 43.5±10.1; saw plametto group: 42.9±11.2;	finasteride 5mg; saw palmetto 325mg	finasteride group: 11.7±3.1, 23.9±5.7, 4.1±2.2, 6.9±2.2; saw plametto group: 10.9±3.2, 24.7±5.3, 4.2±2.1, 7.5±2.1	NR	48 W	North America	single	finasteride: 2; saw plametto: 3[K]

Study ID	**No. arms (included/total)**	**Fund**	**No. randomized**	**Mean age (years)**	**Dose (mg daily)**	**Baseline (Pain/Total/Urinary/QoL)**	**Aftertreatment (Pain/Total/Urinary/QoL)**	**Length of trial**	**Recruitment place**	**Multi/ single center**	**No. adverse event**

Nickel 2004	2/2	Yes	64	finasteride group: 46.9±1.7; placebo group: 41.7±2.1	finasteride 5mg qd; placebo 5mg	finasteride group: NR, 20.1±1.4, NR, NR; placebo group: NR, 22.5±1.6, NR, NR	finasteride group: NR, 19.1±1.5, NR, NR; placebo group: NR, 21.7±1.6, NR, NR	24 W	North America	multi	finasteride: 5; placebo: 7[L]
Macchione 2019	2/2	No	83	Deprox 500® group: 42.7±11.6; serenoa repens group: 39.9±11.6	Deprox 500® bid; Serenoa repens 320 mg	Deprox 500® group: NR, 20.4±2.1, NR, NR; serenoa repens group: NR, 19.6±2.4, NR, NR	Deprox 500® group: NR, 3.1±3.8, NR, NR; serenoa repens group: NR, 6.0±5.3, NR, NR	6 W	Europe	single	No
Cheah 2003	2/2	NR	100	terazosin group: 36 (24-49); placebo group: 35 (20-50)	terazosin dosage 1mg for 4 days, 2mg for 10 days and 5mg for 12 W; placebo had a similar appearance	Terazosin group: 11.7±3.4, 25.1±7.1, 5.1±3.7, 8.3±2.6 ; placebo group: 12.6±3.7, 27.2±7.7, 5.4±3.7, 8.9±2.3	Terazosin group: 6.4±5.2, 13.4±7.6, 2.5±2.8, 4.3±3.3; placebo group: 7.9±5.8, 18.1±10.2, 4.1±3.6, 5.6±3.5	14 W	North America and Asia	single	terazosin: 18; placebo: 9[M]

Study ID	**No. arms (included/total)**	**Fund**	**No. randomized**	**Mean age (years)**	**Dose (mg daily)**	**Baseline (Pain/Total/Urinary/QoL)**	**Aftertreatment (Pain/Total/Urinary/QoL)**	**Length of trial**	**Recruitment place**	**Multi/ single center**	**No. adverse event**
Kim 2011b	3/3	NR	100	tamsulosin group: 45.7; tamsulosin	tamsulosin 0.2 mg , diclofenac	tamsulosin group: 11.4±3.9, 23.6±5.5, 4.2±2.0, 8.1±2.3;	tamsulosin group: 4.0±3.2, 8.6±4.1, 1.6±1.5, 3.0±1.98; tamsulosin plus	12 W	Asia	multi	NR

				plus diclofenac group: 46.5; tamsulosin plus ciprofloxacin group: 46.1	100mg, ciprofloxacin 1000mg	tamsulosin plus diclofenac group: 11.6±3.5, 23.9±5.9, 4.0±2.1, 7.9±2.6; tamsulosin plus ciprofloxacin group: 10.9±4.0, 23.17±5.5, 3.7±1.5, 8.5±2.1	diclofenac group: 3.4±2.7, 7.7±4.9, 1.5±1.2, 2.81±1.75; tamsulosin plus ciprofloxacin group: 3.7±2.8, 8.1±4.2, 1.7±1.0, 2.8±1.5				
Nickel 2003	2/2	Yes	80	levofloxacin group: 56.0; placebo group: 56.2	levofloxacin 500mg; placebo 500mg	levofloxacin group: 10.6±4.7, 24.4±8.2, 5.4±3.0, 4.7±1.6; placebo group: 9.4±3.7, 21.3±7.2, 4.3±2.9, 4.5±1.2	levofloxacin group: 8.9±5.0, 19.0±9.5, 4.2±3.0, 3.7±1.8; placebo group: 7.6±4.7, 18.4±9.1, 4.1±2.8, 3.8±1.7	6 W	North America	multi	levofloxacin: 9; placebo: 6^{N}
Wang 2016	3/3	No	115	levofloxacin group: 36.6±8.1; terazosin group: 38.5±8.3; levofloxacin plus terazosin group: 37.8±7.9	levofloxacin 400mg; terazosin 2mg	levofloxacin group: NR, 24.2±7.1, NR, NR; terazosin group: NR, 22.3±6.6, NR, NR; levofloxacin plus terazosin group: NR, 22.6±6.8, NR, NR	NR	6 W	Asia	single	NR
Study ID	**No. arms (included/total)**	**Fund**	**No. randomized**	**Mean age (years)**	**Dose (mg daily)**	**Baseline (Pain/Total/Urinary/QoL)**	**Aftertreatment (Pain/Total/Urinary/QoL)**	**Length of trial**	**Recruitment place**	**Multi/ single center**	**No. adverse event**

Maurizi 2019	2/2	NR	54	Deprox 500® group: 34±5.9; quercitina group: 33.7±4.62	DEPROX 500® 2 capsules; quercetin 1000mg	Deprox 500® group: NR, 25.67±1.62, NR, NR; quercitina group: NR, 25.96±1.63, NR, NR	Deprox 500® group: NR, 12.22±1.84, NR, NR; quercitina group: NR, 14.85±1.85, NR, NR	4 W	Europe	single	No
Nickel 2011b	2/3	NR	151	silodosin 4mg group: 49.2±13.3; silodosin 8mg group: 46.7±15.6; placebo group: 49.0±11.6	silodosin 4mg; silodosin 8mg; placebo	silodosin 4mg group: 12.3±2.7, 26.0±6.3, 5.3±3.0, 8.3±2.1; silodosin 8mg group: 12.4±3.2, 26.8±5.9, 5.7±2.5, 8.7±2.2; placebo group: 12.8±3.1, 27.9±6.2, 6.5±2.8, 8.6± 2.4	NR	12 W	North America	multi	silodosin 4mg: 3; silodosin 8mg: 6; placebo: 1

Study ID	**No. arms (included/total)**	**Fund**	**No. randomized**	**Mean age (years)**	**Dose (mg daily)**	**Baseline (Pain/Total/Urinary/QoL)**	**Aftertreatment (Pain/Total/Urinary/QoL)**	**Length of trial**	**Recruitment place**	**Multi/ single center**	**No. adverse event**

Morgia 2010	2/2	NR	102	profluss group: 40.96±8.33; serenoa group: 35.9±6.76	Profluss group: 320mg of S. repens oil extract (85%), 5mg oil extracted lycopene (6%) and 0.05mg seleniated sodium, qd; serenoa group: 320mg of S. repens	profluss group: NR, 27.45±4.96, NR, NR; serenoa group: NR, 27.76±3.27, NR, NR	profluss group: NR, 16.02±8.36, NR, NR; serenoa group: NR, 21.00±4.24, NR, NR	8 W	Europe	multi	profluss: 5; serenoa: 4[O]

Study ID	**No. arms (included/total)**	**Fund**	**No. randomized**	**Mean age (years)**	**Dose (mg daily)**	**Baseline (Pain/Total/Urinary/QoL)**	**Aftertreatment (Pain/Total/Urinary/QoL)**	**Length of trial**	**Recruitment place**	**Multi/ single center**	**No. adverse event**

Cai 2014	2/2	NR	87	Deprox 500® group: 33.8±6.78; ibuprofen group: 33.7±5.44	Deprox 500® 2 capsules; ibuprofen 1800mg	Deprox 500® group: NR, 24.9±2.1, NR, NR; ibuprofen group: NR, 25.5±3.0, NR, NR	Deprox 500® group: NR,12.8±2.2, NR, NR; ibuprofen group: NR, 19.5±2.1, NR, NR	4 W	Europe	single	Deprox 500®: 1; ibuprofen: 7[P]

SD: standard difference; QoL: quality of life; NR: not reported; W: weeks;

A: 11 in the PPS group and 4 in the placebo group discontinued due to adverse events or intercurrent illnesses;

B: 41 reported grade 1 (mild), 33 reported grade 2 (moderate), 2 reported grade 3 (severe) adverse events, most adverse events involved nonpelvic pain and gastrointestinal disturbances, and the grade 3 events were progression of chronic pelvic pain in 1 patient receiving placebo and 1 patient receiving ciprofloxacin alone;

C: 1 patient taking placebo developed a rash that resolved when he stopped taking the capsules. One patient taking quercetin developed a headache after the first few doses, which resolved, and 1 patient taking quercetin noted mild tingling of the extremities after each dose;

D: Fifteen of the 100 participants reported at least one adverse event, two in placebo group and 13 in tamsulosin group. Most events, e.g. arrhythmia, heartburn, nausea, dizziness, were classified as mild or moderate;

E: mild gastrointestinal disorders that caused a short treatment interruption and moderate pain (not otherwise specified) that caused discontinuation of treatment. Serious adverse events were reported in 3 patients in the pollen extract group and in 2 patients in the placebo group;

F: 2 in the celecoxib group had mild diarrhea and a skin rash and 1 in the placebo group had nausea. However, the adverse experience did not prevent the patient from completing the study;

G: the adverse event of prostatitis was reported by 150 of 4126 men randomized to placebo vs 102 of 4105 randomized to dutasteride;

H: alfuzosin group (mild: 13; moderate: 19; severe: 7), placebo group (mild: 16; moderate: 19; severe: 3);

I: tamsulosin group (nausea 1, dizziness 2), tamsulosin and dapoxetine group (nausea 5, dizziness 7, dry 1, headache 1);

J: doxazosin group and combination group versus placebo group: dizziness (3 and 4 versus 2 patients), postural hypotension (3 and 4 versus 1), gastrointestinal complaints (2 and 6 versus 2), palpitation (1 and 2 versus 1), flulike syndrome (2 and 1 versus 0), and headache (1 and 0 versus 1);

K: headache (3 cases) in the saw palmetto group and decreased libido (2 cases) in the finasteride group;

L: finasteride group (decreased libido in 2, mood change, fatigue, gastrointestinal discomfort), placebo group (decreased libido, decreased ejaculatory volume, rash, dry mouth, throat constriction, increase in acne, weight gain);

M: Side effects in the terazosin and placebo groups included dizziness (7 versus 2 patients), asthenia (7 versus 3), postural hypotension (1 versus 0), palpitation (1 versus 2), flu syndrome (1 versus 0), drowsiness (2 versus 1), headache (2 versus 2) and rashes (0 versus 1);

N: levofloxacin group (3 with joint/muscular pain, 2 with upper respiratory complaints, 2 with depression, and 1 with a complaint of tendonitis and pruritus), placebo group (2 with sleep disorders, 2 with upper respiratory complaints, and 1 with a complaint of nausea and fatigue);

O: 7 patients reported gastrointestinal symptoms, including constipation, nausea, and meteorism 4 in the Profluss group and 3 in the Serenoa group, and 2 reported cephalea: 1 patient in each group;

P: in group A, 1/41 patients had mild adverse effects that did not require additional treatment (nausea), while in group B, 7/46 patients reported nausea and epigastric pain.

eTable 4. Risk of bias of included studies

(a) Summary of assessment results

Study ID	Bias arising from the randomization process	Bias due to deviations from intended intervention	Bias due to missing outcome data	Bias in measurement of the outcome	Bias in selection of the reported result	Overall
Ye 2008	Some concerns	Some concerns	Low	High	Some concerns	Some concerns
Nickel 2005	Some concerns	Some concerns	High	Low	Some concerns	Some concerns
Alexander 2004	Some concerns	Some concerns	Some concerns	Some concerns	Some concerns	Some concerns
Cai 2017	Low	Some concerns	High	High	Some concerns	Some concerns
Shoskes 1999	Some concerns	Some concerns	High	Low	Some concerns	Some concerns
Chen 2011	Low	Some concerns	Low	Low	Some concerns	Low
Wagenlehner 2009	Low	Low	High	Low	High	Some concerns
Zhao 2009	Low	Some concerns	Low	Low	High	Some concerns
Nickel 2011a	Low	Low	Low	Low	Low	Low
Pontari 2010	Low	Low	Low	Low	Low	Low

Nickel 2008	Low	Low	High	Low	Low	Low
Kim 2011a	Some concerns	Some concerns	High	High	Some concerns	Some concerns
Zhao 2019	Some concerns	Some concerns	High	High	Some concerns	Some concerns
Tuğcu 2007	Some concerns	Some concerns	Some concerns	Some concerns	Some concerns	Some concerns
Kaplan 2004	Some concerns	Some concerns	Low	Some concerns	Some concerns	Some concerns
Nickel 2004	Some concerns	Some concerns	High	Low	Some concerns	Some concerns
Macchione 2019	Low	Low	High	Low	Some concerns	Some concerns
Cheah 2003	High	Some concerns	Low	High	Some concerns	Some concerns
Kim 2011b	Some concerns	Some concerns	High	High	Some concerns	Some concerns
Nickel 2003	Low	Low	Low	Low	Some concerns	Low
Wang 2016	Low	Some concerns	Low	High	Low	Some concerns
Maurizi 2019	Some concerns	Some concerns	Low	Low	Some concerns	Some concerns

Nickel 2011b	Some concerns	Low	Some concerns	Low	Low	Low
Morgia 2010	Low	Low	Low	Some concerns	High	Low
Cai 2014	Low	Some concerns	Low	High	Some concerns	Some concerns

(b). Specific judgment

Ye 2008	
Bias \ Authors' judgment	Support for judgment
Random sequence generation (selection bias)	No mention was in the paper and author was not contacted.
Allocation concealment (selection bias)	No mention was in the paper and author was not contacted.
Blinding of participants and personnel (performance bias)	"Patients and healthcare workers were not blinded to the treatment regimens."
Blinding of outcome assessment (detection bias)	"Patients and healthcare workers were not blinded to the treatment regimens."
Incomplete outcome data (attrition)	"All patients completed the treatments and were scored according to prostatitis symptoms in follow-up checks"
Selective reporting (reporting bias)	Based on checking the outcome measures in Analysis and Method sections, there might be no selective report.
Other bias	Large baseline differences in NIH-CPSI scores. No other baseline characteristics were reported.
Nickel 2005	
Random sequence generation (selection bias)	"Randomization was stratified according to symptom severity at screening, as measured by the SSI", but no further information was provided.
Allocation concealment (selection bias)	"Randomization was stratified according to symptom severity at screening, as measured by the SSI", but no further information was provided.
Blinding of participants and personnel (performance bias)	"Double-blind", but no specific blind method was described.
Blinding of outcome assessment (detection bias)	"Double-blind" and identical-looking placebo was used.
Incomplete outcome data (attrition)	"Of the enrolled patients 73% completed the 16-week study. Of 27 patients who did not complete the study 11 in the PPS group and 4 in the placebo group discontinued due to adverse events or intercurrent illnesses"
Selective reporting (reporting bias)	Based on checking the outcome measures in Analysis and Method sections, there might be no selective report.
Other bias	No other sources of bias were identified.
Alexander 2004	
Random sequence generation (selection bias)	"Each patient was randomly assigned by computer"

Allocation concealment (selection bias)	"The research pharmacist at each site provided the blinded study drugs in 2 tamper-evident bottles."
Blinding of participants and personnel (performance bias)	"All clinical investigators, research nurses, and patients were blinded to treatment assignments until all patients had completed follow-up."
Blinding of outcome assessment (detection bias)	"All clinical investigators, research nurses, and patients were blinded to treatment assignments until all patients had completed follow-up."
Incomplete outcome data (attrition)	"174 men (89%) were available for evaluation at 6 weeks" and "Another 3 participants withdrew from study follow-up between 6 and 12 week".
Selective reporting (reporting bias)	All prespecified outcomes in the published protocol were reported.
Other bias	No other sources of bias were identified.
Cai 2017	
Random sequence generation (selection bias)	"All patients who met the inclusion criteria were randomized to either the treatment or control group by using a computer-generated allocation sequence"
Allocation concealment (selection bias)	No mention was in the paper and author was not contacted.
Blinding of participants and personnel (performance bias)	"Neither the physicians nor the patients were blinded to the treatment type."
Blinding of outcome assessment (detection bias)	"Neither the physicians nor the patients were blinded to the treatment type."
Incomplete outcome data (attrition)	Unbalanced attrition: outcome data were available for 32/36 participants in the pollen extract group and 33/34 in the bromelain group.
Selective reporting (reporting bias)	Protocol not available.
Other bias	No other sources of bias were identified.
Shoskes 1999	
Random sequence generation (selection bias)	Unclear based on "Patients were randomized 1:1 in a double-blind fashion", author was not contacted.
Allocation concealment (selection bias)	Unclear based on "Patients were randomized 1:1 in a double-blind fashion", author was not contacted.
Blinding of participants and personnel (performance bias)	"Patients were randomized 1:1 in a double-blind fashion", it is not clear if personnel were blinded.
Blinding of outcome assessment (detection bias)	"Patients were randomized 1:1 in a double-blind fashion" and "Both quercetin and placebo capsules were identical in appearance."
Incomplete outcome data (attrition)	2/15 participants in the placebo group left due to worsening symptoms (no outcome data).

Selective reporting (reporting bias)	Protocol not available.
Other bias	No other sources of bias were identified.
Chen 2011	
Random sequence generation (selection bias)	"Patients were randomly allocated using computer-generated random numbers into two equal groups"
Allocation concealment (selection bias)	No mention was in the paper and author was not contacted.
Blinding of participants and personnel (performance bias)	"Study investigators and subjects were unaware of the treatment assignments."
Blinding of outcome assessment (detection bias)	"Study investigators and subjects were unaware of the treatment assignments."
Incomplete outcome data (attrition)	"Study investigators and subjects were unaware of the treatment assignments."
Selective reporting (reporting bias)	Protocol not available and author was not contacted.
Other bias	No other sources of bias were identified.
Wagenlehner 2009	
Random sequence generation (selection bias)	"Randomisation was carried out in blocks (n = 4) within the centre using a random number generator."
Allocation concealment (selection bias)	"The investigators were instructed to use the study drug in ascendant order of random numbers available in the respective trial centre."
Blinding of participants and personnel (performance bias)	Double-blind study and "identical capsulation and weight" capsules.
Blinding of outcome assessment (detection bias)	Double-blind study and "identical capsulation and weight" capsules.
Incomplete outcome data (attrition)	12 participants due to outcome data not being available and Post-randomisation exclusion of 9 patients due "to wrong allocation".
Selective reporting (reporting bias)	Outcome was presented only graphically.
Other bias	"These sponsors contributed to the design and conduct of the study as well as data collection" and there were post-randomisation exclusions for "wrong allocation".
Zhao 2009	
Random sequence generation (selection bias)	Computer-generated (Excel spreadsheet).
Allocation concealment (selection bias)	No mention was in the paper and author was not contacted.
Blinding of participants and personnel (performance bias)	Double-blind study, participants were blinded, no information about blinding of study personnel and author was not contacted.

Blinding of outcome assessment (detection bias)	Participants were blinded using placebo.
Incomplete outcome data (attrition)	All outcomes: outcome data were available for all participants.
Selective reporting (reporting bias)	NIH-CPSI scores were presented graphically and author was not contacted.
Other bias	Baseline characteristics not available and author was not contacted.
Nickel 2011a	
Random sequence generation (selection bias)	According to previous study design, "randomization stratified by center."
Allocation concealment (selection bias)	"Randomization stratified by center".
Blinding of participants and personnel (performance bias)	Double-blind and "a steering committee and independent data monitoring committee will provide continuous oversight for the REDUCE study".
Blinding of outcome assessment (detection bias)	Double-blind and "a steering committee and independent data monitoring committee will provide continuous oversight for the REDUCE study".
Incomplete outcome data (attrition)	According to concerned previous publication, "8122 (98.7%) were in efficacy population".
Selective reporting (reporting bias)	"Predefined protocol".
Other bias	No other sources of bias were identified.
Pontari 2010	
Random sequence generation (selection bias)	"Randomly assigned 2:1 in each clinical site via a centrally controlled Web-based data management system to receive treatment with either pregabalin or matching placebo using a permuted block randomization".
Allocation concealment (selection bias)	"Randomly assigned 2:1 in each clinical site via a centrally controlled Web-based data management system to receive treatment with either pregabalin or matching placebo using a permuted block randomization".
Blinding of participants and personnel (performance bias)	"Study investigators and participants were unaware of treatment assignment".
Blinding of outcome assessment (detection bias)	"Study investigators and participants were unaware of treatment assignment".
Incomplete outcome data (attrition)	Outcome data were available for all but 8/218 and 3/106 participants in each group.
Selective reporting (reporting bias)	All outcomes matched the clinical registry.

Other bias	No other sources of bias were identified.
Nickel 2008	
Random sequence generation (selection bias)	"With the use of a centrally controlled, web based data-management system. A permuted-block randomization procedure with randomly assigned block sizes of 4, 6, and 8 was used".
Allocation concealment (selection bias)	"With the use of a centrally controlled, web based data-management system. A permuted-block randomization procedure with randomly assigned block sizes of 4, 6, and 8 was used".
Blinding of participants and personnel (performance bias)	Participants received identical-looking matched placebo and study investigators were unaware of the treatment assignments.
Blinding of outcome assessment (detection bias)	Participants were blinding using placebo.
Incomplete outcome data (attrition)	Loss of follow-up CPSI Experimental 22, placebo 17/134; IIEF 28/138, placebo 25/134; SF12 Experimental 23/138, placebo 21/134; Depression Experimental 23/138, placebo 17/134.
Selective reporting (reporting bias)	Outcomes matched those described in the clinical trial registry.
Other bias	No other sources of bias were identified.
Kim 2011a	
Random sequence generation (selection bias)	No information available and author was not contacted.
Allocation concealment (selection bias)	No information available and author was not contacted.
Blinding of participants and personnel (performance bias)	Single-blind study.
Blinding of outcome assessment (detection bias)	Single-blind study and it is not clear whether included participants.
Incomplete outcome data (attrition)	The author have not specified the reasons for missing outcome data and was not contacted.
Selective reporting (reporting bias)	No protocol available and author was not contacted.
Other bias	Baseline characteristics were not available. Small differences in baseline NIH-CPSI score between groups.
Zhao 2019	
Random sequence generation (selection bias)	"Patients were randomly divided into two treatment groups in a ratio of 1:2" which is insufficient information.
Allocation concealment (selection bias)	"Patients were randomly divided into two treatment groups in a ratio of 1:2" which is insufficient information.
Blinding of participants and personnel (performance bias)	Blinding was impossible cause the interventions were visibly different.

Blinding of outcome assessment (detection bias)	Blinding was impossible cause the interventions were visibly different.
Incomplete outcome data (attrition)	Attrition of outcome data was 10%, 27% and 55% respectively, but withdrawal were not specified.
Selective reporting (reporting bias)	The clinical trial registry offered insufficient information
Other bias	No other sources of bias were identified.
Tuğcu 2007	
Random sequence generation (selection bias)	"Randomised into three groups in order of appearance" which is insufficient, and author was not contacted.
Allocation concealment (selection bias)	"Randomised into three groups in order of appearance" which is insufficient, and author was not contacted.
Blinding of participants and personnel (performance bias)	"Placebo tablets compounded of lactose had a similar, appearance to doxazosin tablets. Ibuprofen was given at a low dose, with an effort to avoid side-effects during the long period of treatment". The interventions were visibly different.
Blinding of outcome assessment (detection bias)	No information on blinding of participants and author was not contacted.
Incomplete outcome data (attrition)	"Eighty-three of the initial 90 patients were eligible for evaluation after 6 mo, and 79 patients were eligible at the end of 12 mo" which is unclear how many patients were evaluated at each time point, and author was not contacted.
Selective reporting (reporting bias)	No protocol available and author was not contacted.
Other bias	No other sources of bias were identified.
Kaplan 2004	
Random sequence generation (selection bias)	No information available and author was not contacted.
Allocation concealment (selection bias)	No information available and author was not contacted.
Blinding of participants and personnel (performance bias)	The study was described as single-blind, but it is not specified who was blinded.
Blinding of outcome assessment (detection bias)	The study is described as single-blind; but it is not specified who was blinded.
Incomplete outcome data (attrition)	Balanced attrition, at 3-month follow-up, missing outcome data < 10%.
Selective reporting (reporting bias)	No protocol available and author was not contacted.
Other bias	No other sources of bias were identified.

Macchione 2019	
Random sequence generation (selection bias)	"Using a pseudo-random number generator software (R statistical software, version 3.2.5)".
Allocation concealment (selection bias)	"Using a pseudo-random number generator software (R statistical software, version 3.2.5)".
Blinding of participants and personnel (performance bias)	"A placebo arm was not included in this trial, and the lack of placebo-arm was evaluated in the results analysis"
Blinding of outcome assessment (detection bias)	"A placebo arm was not included in this trial, and the lack of placebo-arm was evaluated in the results analysis"
Incomplete outcome data (attrition)	"3 patients refused to be treated with phytotherapic drugs, 8 patients were lost and 12 patients showed positive semen culture".
Selective reporting (reporting bias)	No protocol available and author was not contacted.
Other bias	No other sources of bias were identified.
Cheah 2003	
Random sequence generation (selection bias)	"Random number table".
Allocation concealment (selection bias)	No information available and author was not contacted.
Blinding of participants and personnel (performance bias)	Double-blind study. Blinding of personnel not described and author was not contacted.
Blinding of outcome assessment (detection bias)	Participants were blinded with the use of placebo.
Incomplete outcome data (attrition)	14% of participants in each group did not complete assessments.
Selective reporting (reporting bias)	No protocol available and author was not contacted.
Other bias	No other sources of bias were identified.
Nickel 2004	
Random sequence generation (selection bias)	No information available.
Allocation concealment (selection bias)	No information available.
Blinding of participants and personnel (performance bias)	No information about blinding of personnel.
Blinding of outcome assessment (detection bias)	Participants were blinded with the use of placebo.
Incomplete outcome data (attrition)	Only 64/76 participants were available for outcome assessment without sufficient information.

Selective reporting (reporting bias)	No protocol available.
Other bias	No other sources of bias were identified.
Kim 2011b	
Random sequence generation (selection bias)	"Patients were enrolled to subgroup by the table of random sampling digit in each medical center" and author could not provide additional data.
Allocation concealment (selection bias)	No information available and author could not provide additional data.
Blinding of participants and personnel (performance bias)	"We did not have placebo group, because the effects of each drugs such as a-blocker, antibiotics and non-steroidal anti-inflammatory drugs were widely known to the standard drug for CP/CPPS treatment".
Blinding of outcome assessment (detection bias)	Open-label study.
Incomplete outcome data (attrition)	"Other anti-inflammatory drugs or antibiotics while the study was in progress" and "Two patients opted for transurethral resection of the prostate".
Selective reporting (reporting bias)	No protocol available and author could not provide additional data.
Other bias	Some baseline characteristics were missing and author could not provide additional data.
Nickel 2003	
Random sequence generation (selection bias)	"The medications were prepackaged according to a computer-generated, randomized allocation schedule".
Allocation concealment (selection bias)	"The medications were prepackaged according to a computer-generated, randomized allocation schedule".
Blinding of participants and personnel (performance bias)	Double-blind study. Blinding of personnel not described.
Blinding of outcome assessment (detection bias)	Participants were blinded with the use of placebo.
Incomplete outcome data (attrition)	1 participant withdrew from the study.
Selective reporting (reporting bias)	No protocol available.
Other bias	No other sources of bias were identified.
Wang 2016	
Random sequence generation (selection bias)	Computer-generated random blocks.

Allocation concealment (selection bias)	No information available and author was not contacted.
Blinding of participants and personnel (performance bias)	Open-label study.
Blinding of outcome assessment (detection bias)	Open-label study.
Incomplete outcome data (attrition)	Outcome data were available for all participants.
Selective reporting (reporting bias)	Based on clinical trial registry, predefined outcomes were reported.
Other bias	No other sources of bias were identified.
Nickel 2011b	
Random sequence generation (selection bias)	Randomisation schedule created by the sponsor without specifying.
Allocation concealment (selection bias)	Central pharmacy allocation.
Blinding of participants and personnel (performance bias)	Participants and personnel were blinded using placebo.
Blinding of outcome assessment (detection bias)	Participants were blinded using placebo.
Incomplete outcome data (attrition)	It is unclear if all participants had outcome data at 12 weeks follow-up.
Selective reporting (reporting bias)	Based on clinical trial registry, predefined outcomes were reported.
Other bias	No other sources of bias were identified.
Maurizi 2019	
Random sequence generation (selection bias)	No information available and author was not contacted.
Allocation concealment (selection bias)	No information available and author was not contacted.
Blinding of participants and personnel (performance bias)	Only the patients were blinded to the type of treatment.
Blinding of outcome assessment (detection bias)	“The patients enrolled were blinded to the type of treatment”.
Incomplete outcome data (attrition)	Overall outcome data were available for all participants.
Selective reporting (reporting bias)	No protocol available and author was not contacted.

Other bias	No other sources of bias were identified.
Cai 2014	
Random sequence generation (selection bias)	"Single sequence of random assignments (simple randomization) was performed using a pseudo- random number generator software".
Allocation concealment (selection bias)	No concerned information and and author was not contacted.
Blinding of participants and personnel (performance bias)	"This was not a blinded study".
Blinding of outcome assessment (detection bias)	"This was not a blinded study".
Incomplete outcome data (attrition)	Missing data for 1/41 and 2/46 participants in each study arm due to lost to follow-up.
Selective reporting (reporting bias)	No protocol available and author was not contacted.
Other bias	No other sources of bias were identified.
Morgia 2010	
Random sequence generation (selection bias)	"Patients were divided by a computerized, centralized randomization into two equally sized groups".
Allocation concealment (selection bias)	"Patients were divided by a computerized, centralized randomization into two equally sized groups".
Blinding of participants and personnel (performance bias)	"Double-Blind". Blinding of personnel not described and author could not provide additional data.
Blinding of outcome assessment (detection bias)	"Double-Blind". Blinding of personnel not described and author could not provide additional data.
Incomplete outcome data (attrition)	"All enrolled patients completed this study."
Selective reporting (reporting bias)	Primary outcome was presented without variability measurement and some parts were only presented graphically.
Other bias	Number of participants in each arm are not available and author could not provide additional data.

eTable 5. Mixed comparisons

Study ID	NIH-CPSI total scores	NIH-CPSI pain scores	NIH-CPSI urinary scores	NIH-CPSI quality of life scores	Adverse events
Placebo vs. Alfuzosin	-0.62 (-7.21, 5.82) DC: -0.6 (-2.68, 1.48)	-0.3 (-4.91, 4.29) DC: -0.3 (-1.36, 0.76)	-0.21 (-3.13, 2.58) DC: -0.2 (-0.82, 0.42)	0.02 (-1.99, 1.99) DC: 0 (-0.36, 0.36)	1.01 (0.23, 4.39) DC: 1 (0.59, 1.7)
Placebo vs. Bromelain	6.85 (-0.98, 14.74)	1.1 (-5.11, 7.38)	-	-	2.72 (0.1, 144.7)
Placebo vs. Celecoxib	-3.2 (-9.7, 3.26) DC: -3.2(-4.65,-1.75)	-2.62 (-7.35, 2.13) DC: -2.6 (-3.87, -1.33)	0.1 (-2.83, 3.04) DC: 0.1 (-0.73, 0.93)	-1.22 (-3.37, 0.96) DC: -1.2 (-2.08, -0.32)	2.44 (0.13, 123.8) DC: 0.48 (0.04, 5.62)
Placebo vs. Ciprofloxacin	-3.59 (-9.72, 2.83) DC: -2.8 (-5.28, -0.32)	-2.07 (-6.32, 2.34) DC: -1.4 (-2.92, 0.12)	-1.02 (-3.61, 1.64) DC: -0.8 (-1.61, 0.01)	-0.93 (-2.98, 1.03) DC: -0.7 (-1.49, 0.09)	1.34 (0.32, 6.53) DC: 1.09 (0.48, 2.45)
Placebo vs. Doxazosin	-11.47 (-17.87, -5.05)* DC: -11.4 (-11.93, -10.87)*	-4.65 (-9.07, -0.14)* DC: -4.6 (-5.05, -4.15)*	-2.79 (-5.68, -0.11)* DC: -2.8 (-3.29, -2.31)*	-3.99(-5.95, -1.95)* DC: -4 (-4.38, -3.62)*	1.57 (0.28, 9.87) DC: 0.66 (0.21, 2.07)
Placebo vs. Dutasteride	-2.52 (-8.77, 3.72) DC: -2.5 (-3.54, -1.46)*	-1.04 (-5.67, 3.31) DC: -1 (-1.59, -0.41)*	-0.71 (-3.5, 2.13) DC: -0.7 (-1.08, -0.32)*	-0.69 (-2.83, 1.44) DC: -0.7 (-1.07, -0.33)*	-
Placebo vs. Finasteride	-3.46 (-8.89, 1.76) DC: -2.1 (-2.59, -1.61)*	-1.94 (-6.34, 2.23) DC: -1.9 (-2.17, -1.63)*	-0.61 (-3.54, 2.21) DC: -0.6 (-0.75, -0.45)*	-0.29 (-2.34, 1.84) DC: -0.3 (-0.5, -0.1)*	0.6 (0.1, 3.47) DC: 1.63 (0.46, 5.82)
Placebo vs. Ibuprofen	2.62 (-5.27, 10.45)	0.73 (-5.86, 7.18)	-	-	10.09 (0.57, 435.5)
Placebo vs. Levofloxacin	-5.06 (-10.51, 0.32) DC: -2.5 (-4.98, -0.02)*	0.41 (-2.91, 3.81) DC: 0.1 (-1.21, 1.41)	-0.38(-2.46,1.85) DC: -1 (-1.82, -0.18)*	0.49 (-1.07, 2.2) DC: -0.3 (-0.77, 0.17)	1.32 (0.26, 7.17) DC: 0.83 (0.26, 2.6)
Placebo vs. Pollen extract	-3.48 (-8.38, 1.4) DC: -2.5 (-4.43, -0.57)*	-1.58 (-6.21, 2.82) DC: -1.6 (-2.76, -0.44)*	-0.13(-3.01,2.68) DC: -0.1 (-0.63, 0.43)	-0.78 (-2.98, 1.35) DC: -0.8 (-1.53, -0.07)*	0.94 (0.21, 4.13) DC: 1.15 (0.44, 3.03)
Placebo vs. Pentosan polysulfate sodium	-2.63 (-9.8, 4) DC: -2.7 (-5.62, 0.22)	-1 (-5.58, 3.67) DC: -1 (-2.53, 0.53)	-0.7(-3.62,2.27) DC: -1.68 (-0.7, 0.28)	-0.97 (-3.11, 1.24) DC: -4.1 (-4.8, -3.4)*	3.22 (0.53, 21.65) DC: 0.32 (0.1, 1.1)
Placebo vs. Pregabalin	-2.18 (-8.6, 4.21) DC: -2.2 (-3.42, -0.98)*	-0.9 (-5.5, 3.48) DC: -0.9 (-1.56, -0.24)*	-0.49(-3.35,2.18) DC: -0.5 (-0.89, -0.11)*	-0.51 (-2.64, 1.61) DC: -1 (-1.9, -0.1)*	-
Placebo vs. Profluss	-2.82 (-11.78, 6.06)	-	-	-2.23 (-4.75, 0.26)	1.2 (0.05, 33.76)
Placebo vs. Quercetin	-3.31 (-8.85, 1.79) DC: -6.6 (-9.82, -3.38)*	-3.69 (-8.43, 0.99) DC: -3.7 (-5.28, -2.12)*	-1 (-3.96, 2.02) DC: -1 (-2.29, 0.29)	0.49 (-1.07, 2.2) DC: -0.5 (-0.93, -0.07)*	1.97 (0.16, 36.1) DC: 0.46 (0.04, 5.75)
Placebo vs. Saw palmetto	1.78 (-4.31, 7.89)	-0.17 (-6.57, 6.04)	-	-	0.93 (0.07, 12.07)
Placebo vs. Silodosin	-2.66 (-7.63, 2.05) DC: -2.66 (-4.92, -0.40)*	-0.91 (-4.1, 2.5) DC: -0.96 (-2.12, 0.20)	-0.7 (-2.76, 1.41) DC: -1.46 (-0.70, 0.06)	-1.09 (-2.69, 0.59) DC: -1.11 (-1.88, -0.34)*	6.51 (1.07, 63.43)* DC: 0.19 (0.04, 0.91)
Placebo vs. Tamsulosin	-2.2 (-6.62, 2.35) DC: -2.48 (-4.89, -0.07)*	-1.77 (-4.43, 0.93) DC: -1.80 (-3.75, 0.16)	-1.03 (-2.75, 0.81) DC: -1.20 (-0.78, -0.36)*	-0.4 (-1.78, 0.85) DC: -0.10 (-0.42, 0.22)	2.19 (0.73, 7.98) DC: 0.38 (0.05, 3.15)
Placebo vs. Terazosin	-1.44 (-6.97, 4.03) DC: -4.1 (-6.66, -1.54)*	-1.66 (-6.24, 2.96) DC: -1.7 (-3.25, -0.15)*	-1.3 (-4.17, 1.63) DC: -2.18 (-1.3, -0.42)*	-1.31 (-3.48, 0.87) DC: -0.1 (-0.45, 0.25)	2.64 (0.55, 13.32) DC: 0.39 (0.15, 0.98)*

Placebo vs. Ciprofloxacin plus Solifenacin	-7.44 (-16.27, 1.65)	-3.58 (-9.89, 2.63)	-2.81 (-6.82, 1.21)	-1.54 (-4.49, 1.49)	-
Placebo vs. Doxazosin plus Ibuprofen plus Thiocolchicoside	-11.77 (-17.68, -5.77)* DC: -11.7 (-12.17, -11.23)*	-4.76 (-9.15, -0.41)* DC: -4.8 (-5.21, -4.39)*	-3.28 (-6.22, -0.42)* DC: -3.3 (-3.79, -2.81)*	-3.58 (-5.62, -1.56)* DC: -1.3 (-2.19, -0.41)*	3.18 (0.53, 19.3) DC: 0.33 (0.11, 1.05)
Placebo vs. Tamsulosin plus Ciprofloxacin	-1.7 (-6.96, 3.91) DC: -0.7 (-2.91, 1.51)	-1.13 (-4.43, 2.36) DC: -0.2 (-1.52, 1.12)	-0.68 (-2.95, 1.66) DC: -0.6 (-1.35, 0.15)	-0.57 (-2.28, 1.08) DC: -3.6 (-3.95, -3.25)*	1.13 (0.27, 5.22) DC: 1.3 (0.57, 2.94)
Placebo vs. Tamsulosin plus Dapoxetine	-4.76 (-12.64, 3.08)	-	-1.4 (-4.69, 1.98)	-2.22 (-4.69, 0.3)	5.95 (0.67, 67.34)
Placebo vs. Tamsulosin plus Diclofenac	-3.17 (-10.22, 4.07)	-2.36 (-7.04, 2.63)	-1.05 (-3.99, 2.02)	-0.17 (-2.46, 2.09)	-
Placebo vs. Tamsulosin plus Levofloxacin	-6.66 (-13.91, 0.66)	-6.31 (-10.21, -2.36)*	-4.45 (-6.88, -1.85)*	-2.8 (-4.56, -0.94)*	1.79 (0.04, 48.17)
Alfuzosin vs. Bromelain	7.32 (-2.44, 17.36)	1.37 (-6.39, 9.3)	-	-	2.65 (0.07, 177.4)
Alfuzosin vs. Celecoxib	-2.73 (-11.92, 6.3)	-2.3 (-8.62, 4.3)	0.31 (-3.81, 4.54)	-1.23 (-4.22, 1.68)	2.53 (0.09, 161.1)
Alfuzosin vs. Ciprofloxacin	-3 (-11.83, 6.07)	-1.78 (-7.74, 4.65)	-0.82 (-4.65, 3.2)	-0.97 (-3.91, 1.88)	1.35 (0.17, 12.19)
Alfuzosin vs. Doxazosin	-10.83 (-19.98, -1.37)*	-4.31 (-10.84, 2.13)	-2.59 (-6.75, 1.45)	-4.01 (-6.88, -1.14)*	1.56 (0.16, 16.43)
Alfuzosin vs. Dutasteride	-1.94 (-11.2, 7.03)	-0.69 (-7.24, 5.67)	-0.49 (-4.33, 3.55)	-0.73 (-3.62, 2.19)	-
Alfuzosin vs. Finasteride	-2.86 (-11.59, 5.34)	-1.66 (-7.89, 4.87)	-0.4 (-4.26, 3.45)	-0.29 (-3.28, 2.73)	0.6 (0.06, 5.73)
Alfuzosin vs. Ibuprofen	3.26 (-7.25, 13.4)	0.96 (-6.68, 8.82)	-	-	10.16 (0.41, 619.7)
Alfuzosin vs. Levofloxacin	-4.48 (-12.83, 3.97)	0.68 (-4.95, 6.42)	-0.18 (-3.74, 3.52)	0.48 (-2.02, 3.08)	1.34 (0.15, 12.16)
Alfuzosin vs. Pollen extract	-2.89 (-11.07, 5.25)	-1.3 (-7.65, 5.1)	0.09 (-3.97, 4.01)	-0.79 (-3.77, 2.19)	0.93 (0.12, 7.61)
Alfuzosin vs. Pentosan polysulfate sodium	-2.03 (-11.74, 7.14)	-0.71 (-7.3, 5.68)	-0.49 (-4.75, 3.81)	-1 (-3.99, 2.1)	3.23 (0.31, 36.75)
Alfuzosin vs.	-1.58 (-10.79, 7.54)	-0.58 (-6.98, 5.87)	-0.3 (-4.21, 3.61)	-0.52 (-3.44, 2.49)	-

Pregabalin					
Alfuzosin vs. Profluss	-2.17 (-13.4, 8.58)	-	-	-	1.21 (0.03, 45.91)
Alfuzosin vs. Quercetin	-2.74 (-11.24, 5.51)	-3.38 (-9.95, 3.36)	-0.79 (-4.77, 3.33)	-2.25 (-5.54, 0.88)	1.99 (0.11, 49.95)
Alfuzosin vs. Saw palmetto	2.4 (-6.58, 11.27)	0.03 (-7.84, 8.1)	-	-	0.9 (0.05, 17.01)
PAlfuzosin vs. Silodosin	-2.05 (-10.2, 6.15)	-0.63 (-6.13, 5.01)	-0.49 (-4, 3.1)	-1.09 (-3.61, 1.56)	6.51 (0.65, 93.29)
Alfuzosin vs. Tamsulosin	-1.7 (-9.27, 6.48)	-1.49 (-6.71, 3.96)	-0.83 (-4.13, 2.67)	-0.4 (-2.76, 1.88)	2.17 (0.35, 16.86)
Alfuzosin vs. Terazosin	-0.9 (-9.16, 7.41)	-1.39 (-7.78, 5.17)	-1.11 (-4.97, 2.91)	-1.33 (-4.29, 1.73)	2.57 (0.3, 25.03)
Alfuzosin vs. Ciprofloxacin plus Solifenacin	-6.81 (-17.92, 3.99)	-3.3 (-10.74, 4.46)	-2.61 (-7.42, 2.44)	-1.54 (-5.08, 2)	-
Alfuzosin vs. Doxazosin plus Ibuprofen plus Thiocolchicoside	-11.08 (-20.19, -2.44)*	-4.5 (-10.5, 1.7)	-3.06 (-7.12, 1)	-3.61 (-6.46, -0.84)*	3.16 (0.31, 33.37)
Alfuzosin vs. Tamsulosin plus Ciprofloxacin	-1.14 (-9.3, 7.43)	-0.8 (-6.25, 5.01)	-0.49 (-4.09, 3.34)	-0.58 (-3.11, 1.98)	1.14 (0.15, 9.8)
Alfuzosin vs. Tamsulosin plus Dapoxetine	-4.18 (-14.34, 5.69)	-	-1.2 (-5.47, 3.09)	-2.23 (-5.4, 0.9)	5.82 (0.43, 110.3)
Alfuzosin vs. Tamsulosin plus Diclofenac	-2.66 (-12.02, 7.46)	-2.04 (-8.56, 4.68)	-0.87 (-4.95, 3.41)	-0.21 (-3.12, 2.79)	-
Alfuzosin vs. Tamsulosin plus Levofloxacin	-6.05 (-16.34, 3.64)	-5.97 (-12.14, 0.1)	-4.25 (-7.87, -0.22)*	-2.85 (-5.54, -0.18)*	1.82 (0.03, 66.07)
Bromelain vs. Celecoxib	-10.14 (-20.1, 0.24)	-3.72 (-11.58, 4.26)	-	-	0.89 (0.01, 173.7)
Bromelain vs. Ciprofloxacin	-10.38 (-20.41, -0.06)*	-3.18 (-11.09, 4.54)	-	-	0.5 (0.01, 19.53)
Bromelain vs. Doxazosin	-18.2 (-28.42, -8.26)*	-5.69 (-13.37, 1.95)	-	-	0.54 (0.01, 25.9)
Bromelain vs. Dutasteride	-9.28 (-19.13, 0.58)	-2.15 (-10.27, 5.54)	-	-	-
Bromelain vs.	-10.45 (-19.33, -1.39)*	-3.01 (-10.49, 4.4)	-	-	0.21 (0, 9.2)

Finasteride					
Bromelain vs. Ibuprofen	-4.21 (-13.09, 4.94)	-0.39 (-6.66, 5.99)	-	-	3.95 (0.04, 358.7)
Bromelain vs. Levofloxacin	-11.88 (-21.41, -2)*	-0.7 (-7.77, 6.4)	-	-	0.5 (0.01, 20.21)
Bromelain vs. Pollen extract	-10.33 (-16.55, -3.99)* DC: -10.3 (-11.28, -9.32)*	-2.71 (-7.16, 1.71) DC: -2.7 (-2.95, -2.45)*	-	-	0.35 (0.01, 7) DC: 2.19 (0.19, 25.3)
Bromelain vs. Pentosan polysulfate sodium	-9.49 (-19.88, 0.89)	-2.07 (-9.98, 5.76)	-	-	1.22 (0.02, 58.3)
Bromelain vs. Pregabalin	-9.02 (-19.09, 1.18)	-1.96 (-9.53, 5.76)	-	-	-
Bromelain vs. Profluss	-9.68 (-20.32, 1.18)	-	-	-	0.43 (0, 44.32)
Bromelain vs. Quercetin	-10.15 (-18.28, -2.19)	-4.81 (-12.9, 3.2)	-	-	0.73 (0.01, 64.35)
Bromelain vs. Saw palmetto	-5.07 (-13.61, 3.86)	-1.24 (-10.26, 7.45)	-	-	0.33 (0, 20.51)
Bromelain vs. Silodosin	-9.4 (-18.95, -0.64)*	-2 (-9.16, 5.01)	-	-	2.43 (0.03, 130.3)
Bromelain vs. Tamsulosin	-9.03 (-17.75, 0.16)	-2.88 (-9.84, 4.03)	-	-	0.82 (0.01, 27.23)
Bromelain vs. Terazosin	-8.24 (-17.86, 1.25)	-2.79 (-10.82, 5.08)	-	-	0.94 (0.02, 40.97)
Bromelain vs. Ciprofloxacin plus Solifenacin	-14.21 (-26.17, -2.19)*	-4.7 (-13.37, 3.87)	-	-	-
Bromelain vs. Doxazosin plus Ibuprofen plus Thiocolchicoside	-18.5 (-28.43, -8.73)*	-5.85 (-13.49, 1.86)	-	-	1.13 (0.02, 51.93)
Bromelain vs. Tamsulosin plus Ciprofloxacin	-8.55 (-18, 1.12)	-2.23 (-9.38, 5.11)	-	-	0.43 (0.01, 16.09)
Bromelain vs. Tamsulosin plus Dapoxetine	-11.65 (-22.59, -0.28)*	-	-	-	2.2 (0.02, 140.6)
Bromelain vs. Tamsulosin plus Diclofenac	-10.07 (-20.16, 0.77)	-3.45 (-11.33, 4.62)	-	-	-
Bromelain vs.	-13.5 (-24.25, -2.66)*	-7.38 (-14.88, 0.2)	-	-	0.63 (0, 65.71)

Tamsulosin plus Levofloxacin					
Celecoxib vs. Ciprofloxacin	-0.38 (-9.19, 8.73)	0.53 (-5.7, 6.9)	-1.11 (-5.05, 3.02)	0.26 (-2.57, 3.33)	0.54 (0.01, 14.93)
Celecoxib vs. Doxazosin	-8.1 (-17.2, 0.73)	-2.03 (-8.7, 4.77)	-2.87 (-7.02, 1.1)	-2.78 (-5.75, 0.23)	0.63 (0.01, 17.42)
Celecoxib vs. Dutasteride	0.73 (-8.22, 9.62)	1.55 (-4.92, 7.77)	-0.8 (-4.76, 3.16)	0.5 (-2.5, 3.72)	-
Celecoxib vs. Finasteride	-0.28 (-8.71, 7.72)	0.67 (-5.87, 7.16)	-0.73 (-4.8, 3.37)	0.91 (-1.98, 4.01)	0.24 (0, 7.96)
Celecoxib vs. Ibuprofen	5.83 (-4.64, 15.87)	3.32 (-4.6, 11.3)	-	-	4.23 (0.03, 544)
Celecoxib vs. Levofloxacin	-1.88 (-10.17, 6.74)	2.98 (-2.8, 8.74)	-0.51 (-4.05, 3.39)	1.68 (-1, 4.58)	0.54 (0.01, 16.38)
Celecoxib vs. Pollen extract	-0.28 (-8.41, 7.92)	1.01 (-5.45, 7.34)	-0.23 (-4.38, 3.84)	0.43 (-2.63, 3.52)	0.39 (0, 9.72)
Celecoxib vs. Pentosan polysulfate sodium	0.54 (-9.13, 9.73)	1.58 (-5.02, 8.1)	-0.8 (-5.06, 3.24)	0.26 (-2.88, 3.36)	1.29 (0.02, 47.15)
Celecoxib vs. Pregabalin	1.05 (-7.95, 10.09)	1.77 (-4.77, 7.89)	-0.62 (-4.67, 3.33)	0.71 (-2.26, 3.69)	-
Celecoxib vs. Profluss	0.35 (-10.16, 11.36)	-	-	-	0.47 (0, 36.92)
Celecoxib vs. Quercetin	-0.13 (-8.53, 8.34)	-1.06 (-7.57, 5.31)	-1.11 (-5.46, 3.05)	-1.02 (-4.32, 2.23)	0.81 (0.01, 50.84)
Celecoxib vs. Saw palmetto	5 (-3.64, 13.72)	2.41 (-5.69, 10.03)	-	-	0.36 (0, 17.85)
Celecoxib vs. Silodosin	0.62 (-7.72, 8.64)	1.66 (-3.85, 7.5)	-0.81 (-4.42, 2.79)	0.15 (-2.6, 2.89)	2.65 (0.04, 107.1)
Celecoxib vs. Tamsulosin	1.03 (-6.88, 8.91)	0.8 (-4.5, 6.37)	-1.13 (-4.57, 2.27)	0.84 (-1.68, 3.31)	0.89 (0.02, 23.21)
Celecoxib vs. Terazosin	1.82 (-6.82, 10.22)	0.96 (-5.73, 7.6)	-1.39 (-5.37, 2.54)	-0.12 (-3.05, 2.99)	1.08 (0.02, 30.09)
Celecoxib vs. Ciprofloxacin plus Solifenacin	-4.25 (-15.3, 6.76)	-0.99 (-8.76, 6.73)	-2.92 (-7.76, 2.12)	-0.32 (-3.93, 3.43)	-
Celecoxib vs. Doxazosin plus Ibuprofen plus Thiocolchicoside	-8.49 (-17.57, 0.44)	-2.21 (-8.63, 4.54)	-3.41 (-7.54, 0.58)	-2.38 (-5.33, 0.66)	1.3 (0.02, 36.26)
Celecoxib vs.	1.51 (-6.71, 10.09)	1.49 (-4.18, 7.29)	-0.78 (-4.48, 2.91)	0.65 (-2.09, 3.45)	0.45 (0.01, 13.45)

Tamsulosin plus Ciprofloxacin					
Celecoxib vs. Tamsulosin plus Dapoxetine	-1.63 (-11.8, 8.39)	-	-1.52 (-5.79, 2.96)	-0.98 (-4.29, 2.23)	2.37 (0.03, 110.6)
Celecoxib vs. Tamsulosin plus Diclofenac	0.08 (-9.31, 9.73)	0.24 (-6.34, 7.11)	-1.18 (-5.51, 3.18)	1.05 (-2.02, 4.19)	-
Celecoxib vs. Tamsulosin plus Levofloxacin	-3.49 (-13.26, 6.41)	-3.67 (-9.85, 2.48)	-4.58 (-8.19, -0.49)*	-1.62 (-4.44, 1.34)	0.71 (0, 59.57)
Ciprofloxacin vs. Doxazosin	-7.83 (-16.71, 0.83)	-2.58 (-9.09, 3.74)	-1.72 (-5.8, 1.98)	-3.07 (-5.81, -0.25)*	1.15 (0.11, 11.34)
Ciprofloxacin vs. Dutasteride	1.13 (-7.99, 10.02)	1.02 (-5.43, 7.37)	0.31 (-3.59, 4.17)	0.23 (-2.54, 3.05)	-
Ciprofloxacin vs. Finasteride	0.15 (-8.3, 7.74)	0.13 (-6.2, 6.22)	0.4 (-3.55, 4.24)	0.66 (-2.28, 3.58)	0.44 (0.04, 4.18)
Ciprofloxacin vs. Ibuprofen	6.26 (-4.25, 16.23)	2.76 (-4.92, 10.65)	-	-	7.57 (0.28, 406.9)
Ciprofloxacin vs. Levofloxacin	-1.47 (-9.83, 6.85)	2.5 (-2.61, 7.52)	0.64 (-2.57, 3.77)	1.42 (-0.9, 3.87)	0.98 (0.11, 8.21)
Ciprofloxacin vs. Pollen extract	0.14 (-7.81, 7.96)	0.47 (-5.69, 6.73)	0.92 (-3.11, 4.8)	0.15 (-2.78, 3.11)	0.7 (0.08, 5.33)
Ciprofloxacin vs. Pentosan polysulfate sodium	0.94 (-8.81, 9.68)	1.12 (-5.38, 7.15)	0.33 (-3.71, 4.27)	-0.02 (-3.04, 2.89)	2.42 (0.2, 29.28)
Ciprofloxacin vs. Pregabalin	1.39 (-7.58, 10.39)	1.17 (-5.01, 7.7)	0.5 (-3.56, 4.19)	0.45 (-2.6, 3.35)	-
Ciprofloxacin vs. Profluss	0.78 (-10.16, 11.72)	-	-	-	0.85 (0.02, 31.74)
Ciprofloxacin vs. Quercetin	0.27 (-8.55, 8.38)	-1.66 (-8, 4.59)	0.01 (-4.13, 3.92)	-1.27 (-4.63, 1.92)	1.42 (0.07, 35.92)
Ciprofloxacin vs. Saw palmetto	5.32 (-3.35, 14.19)	1.84 (-5.94, 9.58)	-	-	0.66 (0.03, 13.74)
Ciprofloxacin vs. Silodosin	0.94 (-7.35, 8.87)	1.09 (-4.19, 6.59)	0.32 (-3, 3.62)	-0.16 (-2.7, 2.51)	4.89 (0.46, 67.62)
Ciprofloxacin vs. Tamsulosin	1.36 (-4.79, 7.3) DC: 1.8 (-0.9, 4.5)	0.27 (-3.9, 4.63) DC: 0.7 (-0.95, 2.35)	0 (-2.48, 2.48) DC: 0.4 (-0.41, 1.21)	0.53 (-1.43, 2.42) DC: 0.6 (-0.14, 1.34)	1.6 (0.36, 7.57) DC: 0.92 (0.41, 2.06)
Ciprofloxacin vs. Terazosin	2.06 (-6.29, 10.41)	0.35 (-6.09, 6.71)	-0.27 (-4.27, 3.68)	-0.35 (-3.31, 2.66)	1.91 (0.22, 16.81)
Ciprofloxacin vs.	-3.87 (-10.24, 2.54)	-1.54 (-6.08, 2.98)	-1.8 (-4.66, 1.09)	-0.59 (-2.71, 1.64)	-

Ciprofloxacin plus Solifenacin	DC: -3.9 (-5.53, -2.27)*	DC: -1.5 (-2.34, -0.66)*	DC: -1.8 (-2.38, -1.22)*	DC: 0.6 (-0.27, 1.47)	
Ciprofloxacin vs. Doxazosin plus Ibuprofen plus Thiocolchicoside	-8.14 (-17.03, 0.72)	-2.74 (-8.94, 3.49)	-2.25 (-6.23, 1.65)	-2.64 (-5.54, 0.16)	2.32 (0.21, 22.03)
Ciprofloxacin vs. Tamsulosin plus Ciprofloxacin	1.93 (-4.42, 8.23) DC: 2.1 (-0.56, 4.76)	0.95 (-3.59, 5.4) DC: 1.2 (-0.45, 2.85)	0.35 (-2.37, 3.02) DC: 0.2 (-0.57, 0.97)	0.36 (-1.67, 2.32) DC: -0.6 (-1.34, 0.14)	0.83 (0.18, 4.28) DC: 1.19 (0.52, 2.71)
Ciprofloxacin vs. Tamsulosin plus Dapoxetine	-1.17 (-10.26, 7.62)	-	-0.41 (-4.25, 3.29)	-1.26(-4.06, 1.48)	4.34 (0.44, 54.66)
Ciprofloxacin vs. Tamsulosin plus Diclofenac	0.36 (-7.53, 8.51)	-0.25 (-6.02, 5.49)	-0.03 (-3.5, 3.5)	0.72 (-1.86, 3.44)	-
Ciprofloxacin vs. Tamsulosin plus Levofloxacin	-3.02 (-12.91, 6.42)	-4.23 (-9.28, 0.95)	-3.45 (-6.64, -0.18)*	-1.89 (-4.23, 0.48)	1.34 (0.03, 38.98)
Doxazosin vs. Dutasteride	8.92 (0.04, 17.6)*	3.58 (-2.69, 9.89)	2.06 (-1.73, 6.18)	3.3 (0.3, 6.15)*	-
Doxazosin vs. Finasteride	7.92 (-0.25, 16.23)	2.67 (-3.54, 8.91)	2.15 (-1.81, 6.25)	3.69 (0.87, 6.59)*	0.37 (0.03, 4.63)
Doxazosin vs. Ibuprofen	14.02 (3.97, 24.53)*	5.36 (-2.34, 12.98)	-	-	6.74 (0.22, 405.2)
Doxazosin vs. Levofloxacin	6.37 (-2.03, 14.92)	5.06 (-0.48, 10.57)	2.4 (-1.13, 6.07)	4.47 (1.86, 7.22)*	0.85 (0.07, 9.67)
Doxazosin vs. Pollen extract	7.96 (0.03, 15.87)*	3.04 (-3.33, 9.48)	2.65 (-1.26, 6.78)	3.2 (0.16, 6.19)*	0.6 (0.06, 5.94)
Doxazosin vs. Pentosan polysulfate sodium	8.75 (-0.71, 18.17)	3.63 (-2.93, 10.21)	2.07 (-1.83, 6.2)	3.05 (0.07, 5.98)*	2.07 (0.15, 28.21)
Doxazosin vs. Pregabalin	9.19 (0.43, 18.61)*	3.77 (-2.8, 9.96)	2.27 (-1.64, 6.35)	3.49 (0.51, 6.43)*	-
Doxazosin vs. Profluss	8.57 (-2.57, 19.91)	-	-	-	0.76 (0.02, 35.34)
Doxazosin vs. Quercetin	8.14 (-0.12, 16.2)	0.96 (-5.7, 7.55)	1.77 (-2.25, 5.84)	1.76 (-1.48, 4.82)	1.3 (0.05, 38.7)
Doxazosin vs. Saw palmetto	13.2 (4.12, 22.38)*	4.42 (-3.45, 12.1)	-	-	0.59 (0.03, 12.99)
Doxazosin vs. Silodosin	8.72 (0.92, 16.85)*	3.67 (-1.99, 9.26)	2.08 (-1.3, 5.72)	2.9 (0.24, 5.53)*	4.28 (0.29, 69.9)

Doxazosin vs. Tamsulosin	9.22 (1.52, 17.04)*	2.82 (-2.41, 8.32)	1.75 (-1.52, 5.31)	3.6 (1.15, 5.92)*	1.39 (0.17, 12.24)
Doxazosin vs. Terazosin	10.06 (1.5, 18.39)*	2.88 (-3.47, 9.4)	1.5 (-2.48, 5.59)	2.68 (-0.38, 5.61)	1.68 (0.15, 17.97)
Doxazosin vs. Ciprofloxacin plus Solifenacin	3.91 (-7.16, 14.9)	1.09 (-6.79, 9.03)	-0.05 (-4.62, 4.96)	2.49 (-1.12, 5.86)	-
Doxazosin vs. Doxazosin plus Ibuprofen plus Thiocolchicoside	-0.33 (-6.44, 6.01) DC: -0.3 (-0.81, 0.21)	-0.18 (-4.52, 4.42) DC: -0.2 (-0.68, 0.28)	-0.5 (-3.33, 2.34) DC: -0.5 (-0.96, -0.04)*	0.41 (-1.72, 2.4) DC: 0.6 (-0.26, 1.46)	1.99 (0.36, 10.24) DC: 0.51 (0.18, 1.42)
Doxazosin vs. Tamsulosin plus Ciprofloxacin	9.74 (1.39, 18.04)*	3.49 (-1.95, 9.37)	2.08 (-1.27, 5.81)	3.44 (0.79, 6.02)*	0.72 (0.08, 7.01)
Doxazosin vs. Tamsulosin plus Dapoxetine	6.63 (-3.71, 16.86)	-	1.38 (-2.94, 5.65)	1.79 (-1.39, 4.92)	3.73 (0.25, 83.02)
Doxazosin vs. Tamsulosin plus Diclofenac	8.28 (-1.34, 17.76)	2.26 (-4.21, 9.14)	1.73 (-2.35, 5.84)	3.81 (0.82, 6.78)*	-
Doxazosin vs. Tamsulosin plus Levofloxacin	4.75 (-5.14, 14.3)	-1.69 (-7.56, 4.36)	-1.69 (-5.4, 2.33)	1.16 (-1.62, 3.88)	1.12 (0.02, 52.18)
Dutasteride vs. Finasteride	-0.96 (-9.45, 7.06)	-0.9 (-7.04, 5.26)	0.11 (-4.05, 4.11)	0.39 (-2.45, 3.31)	-
Dutasteride vs. Ibuprofen	5.16 (-4.75, 15.24)	1.76 (-5.86, 9.78)	-	-	-
Dutasteride vs. Levofloxacin	-2.61 (-10.68, 5.81)	1.41 (-4.19, 7.25)	0.32 (-3.19, 3.97)	1.19 (-1.34, 3.86)	-
Dutasteride vs. Pollen extract	-0.98 (-8.87, 6.67)	-0.59 (-6.61, 5.85)	0.6 (-3.68, 4.43)	-0.11 (-3.1, 2.96)	-
Dutasteride vs. Pentosan polysulfate sodium	-0.19 (-9.55, 9.03)	0.08 (-6.62, 6.48)	0.03 (-4.01, 4.1)	-0.26 (-3.29, 2.79)	-
Dutasteride vs. Pregabalin	0.3 (-8.66, 9.25)	0.17 (-6.35, 6.86)	0.19 (-3.62, 4.06)	0.2 (-2.67, 3.11)	-
Dutasteride vs. Profluss	-0.27 (-10.99, 10.29)	-	-	-	-
Dutasteride vs. Quercetin	-0.88 (-9.05, 6.85)	-2.59 (-9.11, 3.9)	-0.28 (-4.43, 3.71)	-1.51 (-4.88, 1.68)	-
Dutasteride vs.	4.32 (-4.38, 12.72)	0.84 (-6.67, 8.29)	-	-	-

Saw palmetto					
Dutasteride vs. Silodosin	-0.19 (-8.13, 7.6)	0.08 (-5.28, 5.82)	0.01 (-3.58, 3.5)	-0.37 (-3.02, 2.18)	-
Dutasteride vs. Tamsulosin	0.26 (-7.47, 8.11)	-0.76 (-5.9, 4.56)	-0.33 (-3.59, 3.13)	0.29 (-2.22, 2.73)	-
Dutasteride vs. Terazosin	1.05 (-7.38, 9.36)	-0.62 (-7, 5.95)	-0.57 (-4.7, 3.42)	-0.59 (-3.54, 2.33)	-
Dutasteride vs. Ciprofloxacin plus Solifenacin	-4.95 (-16.02, 6.48)	-2.55 (-10.18, 5.14)	-2.09 (-6.92, 2.6)	-0.82 (-4.38, 2.69)	-
Dutasteride vs. Doxazosin plus Ibuprofen plus Thiocolchicoside	-9.18 (-17.74,-0.46)	-3.76 (-9.94, 2.34)	-2.56 (-6.67, 1.54)	-2.88 (-5.82, -0.02)*	-
Dutasteride vs. Tamsulosin plus Ciprofloxacin	0.86 (-7.42, 9.02)	-0.06 (-5.64, 5.68)	0 (-3.55, 3.76)	0.12 (-2.6, 2.76)	-
Dutasteride vs. Tamsulosin plus Dapoxetine	-2.31 (-12.19, 7.7)	-	-0.72 (-4.88, 3.97)	-1.51 (-4.67, 1.62)	-
Dutasteride vs. Tamsulosin plus Diclofenac	-0.73 (-9.87, 9.13)	-1.36 (-7.53, 5.46)	-0.35 (-4.5, 3.87)	0.52 (-2.65, 3.47)	-
Dutasteride vs. Tamsulosin plus Levofloxacin	-4.17 (-13.65, 5.77)	-5.25 (-11, 0.63)	-3.75 (-7.46, 0.18)	-2.14 (-4.91, 0.64)	-
Finasteride vs. Ibuprofen	6.17 (-2.62, 14.86)	2.61 (-4.44, 10.35)	-	-	17.28 (0.64, 963.8)
Finasteride vs. Levofloxacin	-1.55 (-9.24, 6.34)	2.37 (-3.26, 8.02)	0.22 (-3.09, 3.87)	0.78 (-1.85, 3.55)	2.24 (0.22, 25.47)
Finasteride vs. Pollen extract	-0.03 (-6.14, 6.23)	0.37 (-5.52, 6.48)	0.49 (-3.43, 4.46)	-0.49 (-3.62, 2.53)	1.59 (0.18, 13.69)
Finasteride vs. Pentosan polysulfate sodium	0.7 (-7.82, 9.24)	0.96 (-5.43, 7.52)	-0.09 (-3.97, 4.12)	-0.69 (-3.73, 2.38)	5.28 (0.46, 77.46)
Finasteride vs. Pregabalin	1.27 (-6.77, 9.53)	1.06 (-5.35, 7.45)	0.09 (-3.71, 4.03)	-0.2 (-3.25, 2.73)	-
Finasteride vs. Profluss	0.67 (-7.76, 9.15)	-	-	-	2.05 (0.11, 42.51)
Finasteride vs. Quercetin	0.11 (-6.81, 7.19)	-1.71 (-8.01, 4.59)	-0.38 (-4.51, 3.94)	-1.95(-5.19,1.15)	3.43 (0.15, 101)

Finasteride vs. Saw palmetto	5.25 (-0.2, 10.75) DC: 6.9 (4.79, 9.01)*	1.76 (-2.76, 6.32) DC: 1.8 (1.51, 2.09)*	-	-	1.51 (0.18, 14.47) DC: 0.64 (0.1, 4.14)
Finasteride vs. Silodosin	0.85 (-6.27, 8.23)	1 (-4.2, 6.58)	-0.1 (-3.6, 3.33)	-0.78 (-3.54, 1.83)	11.3 (0.85, 188.1)
Finasteride vs. Tamsulosin	1.23 (-5.42, 8.39)	0.14 (-4.88, 5.48)	-0.42 (-3.69, 3.03)	-0.09 (-2.7, 2.32)	3.72 (0.46, 31.38)
Finasteride vs. Terazosin	2.01 (-5.51, 9.61)	0.26 (-6.15, 6.7)	-0.66 (-4.66, 3.29)	-1.01 (-4.03, 1.97)	4.48 (0.42, 48.48)
Finasteride vs. Ciprofloxacin plus Solifenacin	-3.94 (-14.19, 6.44)	-1.63 (-9.4, 6.08)	-2.22 (-6.97, 2.72)	-1.2 (-4.88, 2.28)	-
Finasteride vs. Doxazosin plus Ibuprofen plus Thiocolchicoside	-8.21 (-16.21, 0.14)	-2.82 (-8.98, 3.46)	-2.67 (-6.82, 1.68)	-3.29 (-6.31, -0.32)*	5.3 (0.45, 60.16)
Finasteride vs. Tamsulosin plus Ciprofloxacin	1.76 (-5.46, 9.61)	0.83 (-4.71, 6.54)	-0.08 (-3.7, 3.84)	-0.28 (-3.05, 2.31)	1.89 (0.2, 19.7)
Finasteride vs. Tamsulosin plus Dapoxetine	-1.25 (-10.76, 8.2)	-	-0.81 (-5.2, 3.64)	-1.92 (-5.21, 1.27)	10.13 (0.62, 196.4)
Finasteride vs. Tamsulosin plus Diclofenac	0.24 (-8.27, 9.58)	-0.43 (-6.67, 6.32)	-0.47 (-4.51, 3.83)	0.09 (-3.04, 3.15)	-
Finasteride vs. Tamsulosin plus Levofloxacin	-3.24 (-12.49, 6.04)	-4.36 (-10.18, 1.43)	-3.86 (-7.53, 0.15)	-2.53 (-5.33, 0.26)	2.98 (0.05, 134.4)
Ibuprofen vs. Levofloxacin	-7.73 (-17.03, 1.96)	-0.36 (-7.93, 7.07)	-	-	0.13 (0, 3.85)
Ibuprofen vs. Pollen extract	-6.12 (-12.46, 0.16) DC: -6.1 (-6.78, -5.42)*	-2.31 (-7.05, 2.09) DC: -2.2 (-2.41, -1.99)*	-	-	0.1 (0, 1.07) DC: 7.18 (0.84, 61.1)
Ibuprofen vs. Pentosan polysulfate sodium	-5.34 (-15.72, 5.08)	-1.77 (-9.48, 5.95)	-	-	0.32 (0, 10.29)
Ibuprofen vs. Pregabalin	-4.81 (-15, 5.34)	-1.57 (-9.24, 6.19)	-	-	-
Ibuprofen vs. Profluss	-5.54 (-15.94, 5.09)	-	-	-	0.11 (0, 7.62)
Ibuprofen vs. Quercetin	-5.99 (-14.33, 2.09)	-4.4 (-12.63, 3.36)	-	-	0.19 (0, 10.21)
Ibuprofen vs. Saw	-0.94 (-9.6, 7.55)	-0.92 (-9.89, 7.67)	-	-	0.09 (0, 3.7)

palmetto					
Ibuprofen vs. Silodosin	-5.36 (-14.77, 3.92)	-1.65 (-8.88, 5.66)	-	-	0.66 (0.01, 23.46)
Ibuprofen vs. Tamsulosin	-4.85 (-13.86, 4.43)	-2.51 (-9.47, 4.63)	-	-	0.21 (0, 5.15)
Ibuprofen vs. Terazosin	-4.12 (-13.63, 5.4)	-2.41 (-10.19, 5.31)	-	-	0.26 (0, 7.47)
Ibuprofen vs. Ciprofloxacin plus Solifenacin	-9.96 (-21.83, 2.13)	-4.32 (-13.5, 4.24)	-	-	-
Ibuprofen vs. Doxazosin plus Ibuprofen plus Thiocolchicoside	-14.36 (-24.58, -3.9)*	-5.56 (-13.24, 1.98)	-	-	0.31 (0.01, 8.66)
Ibuprofen vs. Tamsulosin plus Ciprofloxacin	-4.36 (-13.99, 5.18)	-1.81 (-9.07, 5.49)	-	-	0.11 (0, 3.16)
Ibuprofen vs. Tamsulosin plus Dapoxetine	-7.49 (-18.61, 3.73)	-	-	-	0.57 (0.01, 26.47)
Ibuprofen vs. Tamsulosin plus Diclofenac	-5.87 (-16.31, 4.81)	-3.1 (-11.05, 5.2)	-	-	-
Ibuprofen vs. Tamsulosin plus Levofloxacin	-9.31 (-19.88, 1.76)	-7.03 (-14.63, 0.47)	-	-	0.16 (0, 14.95)
Levofloxacin vs. Pollen extract	1.61 (-6.04, 8.92)	-1.93 (-7.5, 3.47)	0.25 (-3.46, 3.8)	-1.28 (-4.13, 1.45)	0.72 (0.07, 6)
Levofloxacin vs. Pentosan polysulfate sodium	2.35 (-6.81, 10.95)	-1.36 (-7.37, 4.62)	-0.32 (-4.09, 3.32)	-1.46 (-4.19, 1.24)	2.44 (0.21, 28.07)
Levofloxacin vs. Pregabalin	2.85 (-5.51, 11.33)	-1.3 (-6.8, 4.42)	-0.15 (-3.74, 3.11)	-0.99 (-3.72, 1.62)	-
Levofloxacin vs. Profluss	2.26 (-8.55, 12.73)	-	-	-	0.94 (0.02, 35.26)
Levofloxacin vs. Quercetin	1.77 (-6.01, 9.25)	-4.07 (-9.84, 1.73)	-0.62 (-4.42, 2.98)	-2.7 (-5.71, 0.09)	1.47 (0.07, 40.94)
Levofloxacin vs. Saw palmetto	6.85 (-1.74, 14.94)	-0.58 (-7.93, 6.57)	-	-	0.69 (0.03, 13.67)
Levofloxacin vs. Silodosin	2.37 (-5.07, 9.74)	-1.31 (-6.17, 3.28)	-0.31 (-3.25, 2.56)	-1.6 (-3.89, 0.69)	5.06 (0.4, 76.99)

Levofloxacin vs. Tamsulosin	2.84 (-4.41, 9.76)	-2.2 (-5.45, 1.28) DC: -1.3 (-2.46, -0.14)*	-0.65 (-2.9, 1.47) DC: -0.6 (-1.66, 0.46)	-0.88 (-2.61, 0.58) DC: 0.4 (0.07, 0.73)*	1.64 (0.25, 11.23) DC: 1 (0.02, 52.73)
Levofloxacin vs. Terazosin	3.64 (-1.61, 8.92) DC: 5.9 (4.15, 7.65)*	-2.06 (-7.87, 3.84)	-0.92 (-4.6, 2.64)	-1.8 (-4.55, 0.81)	1.99 (0.19, 20.1)
Levofloxacin vs. Ciprofloxacin plus Solifenacin	-2.38 (-12.74, 8.2)	-3.95 (-11.02, 2.86)	-2.41 (-6.68, 1.72)	-2.01 (-5.34, 1.22)	-
Levofloxacin vs. Doxazosin plus Ibuprofen plus Thiocolchicoside	-6.64 (-14.86, 1.56)	-5.2 (-10.76, 0.4)	-2.89 (-6.73, 0.51)	-4.05 (-6.82, -1.55)*	2.38 (0.19, 26.34)
Levofloxacin vs. Tamsulosin plus Ciprofloxacin	3.36 (-3.85, 11.19)	-1.53 (-5.72, 2.97)	-0.29 (-3.12, 2.42)	-1.07 (-3.16, 0.98)	0.83 (0.1, 7.69)
Levofloxacin vs. Tamsulosin plus Dapoxetine	0.3 (-9.5, 9.9)	-	-1.02 (-4.64, 2.55)	-2.73 (-5.48, -0.02)*	4.46 (0.31, 78.02)
Levofloxacin vs. Tamsulosin plus Diclofenac	1.9 (-7.04, 10.84)	-2.8 (-7.74, 2.85)	-0.68 (-4.05, 2.62)	-0.68 (-3.21, 1.7)	-
Levofloxacin vs. Tamsulosin plus Levofloxacin	-1.62 (-7.96, 4.54) DC: -0.5 (-2.41, 1.41)	-6.67 (-10.48, -2.86)* DC: -7.9 (-8.96, -6.84)*	-4.1 (-6.49, -1.52)* DC: -5.1 (-6.02, -4.18)*	-3.33 (-5.23, -1.59)* DC: -1.6 (-2.45, -0.75)*	1.33 (0.03, 46.72) DC: 1 (0.02, 52.73)
Pollen extract vs. Pentosan polysulfate sodium	0.71 (-7.53, 9.04)	0.61 (-5.75, 6.99)	-0.59 (-4.75, 3.77)	-0.19 (-3.24, 3)	3.42 (0.31, 39.21)
Pollen extract vs. Pregabalin	1.2 (-6.63, 9.46)	0.69 (-5.71, 7.15)	-0.41 (-4.25, 3.55)	0.28 (-2.74, 3.48)	-
Pollen extract vs. Profluss	0.62 (-7.74, 9.19)	-	-	-	1.25 (0.04, 40.04)
Pollen extract vs. Quercetin	0.16 (-5.09, 5.17) DC: 2.4 (1.81, 2.99)*	-2.15 (-8.81, 4.44)	-0.87(-5,3.2)	-1.44 (-4.83, 1.85)	2.03 (0.14, 46.18) DC: 0.62 (0.02, 22.02)
Pollen extract vs. Saw palmetto	5.24 (-0.23, 10.88) DC: 3.7 (1.95, 5.45)*	1.37 (-6.45, 9.12)	-	-	0.98 (0.06, 15.42) DC: 1.07 (0.02, 55.41)
Pollen extract vs. Silodosin	0.79 (-6.06, 7.72)	0.68 (-4.87, 6.29)	-0.58 (-4.17, 2.94)	-0.28 (-2.97, 2.59)	7.05 (0.63, 97.62)
Pollen extract vs. Tamsulosin	1.24 (-5.28, 8.19)	-0.23 (-5.37, 5.22)	-0.91 (-4.28, 2.71)	0.39 (-2.1, 2.87)	2.33 (0.37, 16.69)
Pollen extract vs. Terazosin	2.04 (-5.17, 9.34)	-0.09 (-6.41, 6.53)	-1.19 (-5.03, 2.84)	-0.53 (-3.66, 2.57)	2.8 (0.29, 25.61)
Pollen extract vs.	-3.95 (-13.76, 6.6)	-1.96 (-9.52, 5.48)	-2.7 (-7.49, 2.35)	-0.76 (-4.43, 2.99)	-

Ciprofloxacin plus Solifenacin					
Pollen extract vs. Doxazosin plus Ibuprofen plus Thiocolchicoside	-8.29 (-16.13, -0.62)*	-3.22 (-9.47, 3.06)	-3.17 (-7.12, 0.94)	-2.81 (-5.81, 0.14)	3.4 (0.31, 36.53)
Pollen extract vs. Tamsulosin plus Ciprofloxacin	1.78 (-5.35, 9.11)	0.51 (-4.99, 6.13)	-0.54 (-4.13, 3.32)	0.21 (-2.53, 2.94)	1.23 (0.16, 10.16)
Pollen extract vs. Tamsulosin plus Dapoxetine	-1.34 (-10.57, 8.11)	-	-1.3 (-5.5, 3.19)	-1.45 (-4.66, 1.94)	6.62 (0.44, 118)
Pollen extract vs. Tamsulosin plus Diclofenac	0.24 (-8.01, 9.06)	-0.76 (-7.07, 6.13)	-0.94 (-5.06, 3.32)	0.6 (-2.57, 3.74)	-
Pollen extract vs. Tamsulosin plus Levofloxacin	-3.21 (-11.71, 5.68)	-4.72 (-10.68, 1.31)	-4.34 (-8.2, -0.46)*	-2.06 (-4.78, 0.81)	1.91 (0.03, 69.37)
Pentosan polysulfate sodium vs. Pregabalin	0.48 (-8.72, 9.92)	0.11 (-6.42, 6.52)	0.19 (-3.94, 4.27)	0.47 (-2.49, 3.46)	-
Pentosan polysulfate sodium vs. Profluss	-0.06 (-11.25, 10.98)	-	-	-	0.38 (0.01, 16.66)
Pentosan polysulfate sodium vs. Quercetin	-0.67 (-9.31, 7.85)	-2.69 (-9.44, 4.06)	-0.31 (-4.46, 3.82)	-1.26 (-4.67, 1.99)	0.61 (0.03, 18.28)
Pentosan polysulfate sodium vs. Saw palmetto	4.57 (-4.63, 13.64)	0.85 (-7.36, 8.55)	-	-	0.28 (0.01, 7.7)
Pentosan polysulfate sodium vs. Silodosin	0.01 (-8.3, 8.43)	0.06 (-5.52, 5.83)	-0.01 (-3.56, 3.6)	-0.11 (-2.83, 2.6)	2.04 (0.15, 33.15)
Pentosan polysulfate sodium vs. Tamsulosin	0.45 (-7.51, 9.05)	-0.82 (-6.09, 4.48)	-0.33 (-3.76, 3.13)	0.58 (-2.06, 3.12)	0.67 (0.08, 6.35)
Pentosan polysulfate sodium vs. Terazosin	1.29 (-7.46, 10.07)	-0.65 (-7.1, 6.1)	-0.62 (-4.77, 3.5)	-0.32 (-3.41, 2.62)	0.81 (0.07, 9.47)
Pentosan polysulfate sodium	-4.73 (-15.77, 6.66)	-2.6 (-10.32, 5.21)	-2.1 (-7.09, 2.93)	-0.54 (-4.25, 3.19)	-

vs. Ciprofloxacin plus Solifenacin					
Pentosan polysulfate sodium vs. Doxazosin plus Ibuprofen plus Thiocolchicoside	-8.95 (-18.28, 0.31)	-3.79 (-10.04, 2.39)	-2.59 (-6.68, 1.4)	-2.64 (-5.69, 0.38)	0.96 (0.07, 12.58)
Pentosan polysulfate sodium vs. Tamsulosin plus Ciprofloxacin	0.97 (-7.45, 10.05)	-0.09 (-5.9, 5.63)	0.01 (-3.75, 3.64)	0.4 (-2.38, 3.17)	0.35 (0.03, 3.78)
Pentosan polysulfate sodium vs. Tamsulosin plus Dapoxetine	-2.03 (-12.31, 8.22)	-	-0.73 (-5.03, 3.69)	-1.24 (-4.5, 2.16)	1.82 (0.1, 38.87)
Pentosan polysulfate sodium vs. Tamsulosin plus Diclofenac	-0.46 (-9.92, 9.51)	-1.35 (-7.68, 5.14)	-0.39 (-4.57, 4.06)	0.77 (-2.41, 3.81)	-
Pentosan polysulfate sodium vs. Tamsulosin plus Levofloxacin	-3.98 (-13.95, 6.19)	-5.29 (-11.34, 0.75)	-3.78 (-7.57, 0.25)	-1.85 (-4.72, 1.04)	0.55 (0.01, 24.87)
Pregabalin vs. Profluss	-0.63 (-11.61, 10.39)	-	-	-	-
Pregabalin vs. Quercetin	-1.14 (-9.65, 7.22)	-2.83 (-9.45, 3.8)	-0.5 (-4.49, 3.79)	-1.74 (-4.97, 1.52)	-
Pregabalin vs. Saw palmetto	4.01 (-5.06, 12.67)	0.71 (-7.21, 8.34)	-	-	-
Pregabalin vs. Silodosin	-0.5 (-8.64, 7.72)	-0.06 (-5.57, 5.68)	-0.22 (-3.51, 3.27)	-0.58 (-3.25, 2.06)	
Pregabalin vs. Tamsulosin	0.02 (-7.72, 7.77)	-0.91 (-5.95, 4.32)	-0.52 (-3.69, 2.86)	0.09 (-2.35, 2.54)	-
Pregabalin vs. Terazosin	0.8 (-7.86, 8.77)	-0.84 (-7.02, 5.63)	-0.77 (-4.66, 3.29)	-0.8 (-3.79, 2.16)	-
Pregabalin vs. Ciprofloxacin plus Solifenacin	-5.34 (-16.09, 6.11)	-2.65 (-10.62, 4.83)	-2.31 (-7, 2.67)	-1.03 (-4.56, 2.75)	-
Pregabalin vs. Doxazosin plus Ibuprofen plus	-9.53 (-18.57, -0.5)*	-3.96 (-10.35, 2.61)	-2.78 (-6.71, 1.08)	-3.11 (-5.94, -0.14)*	-

Thiocolchicoside					
Pregabalin vs. Tamsulosin plus Ciprofloxacin	0.51 (-7.88, 8.94)	-0.24 (-5.83, 5.39)	-0.17 (-3.47, 3.57)	-0.08 (-2.88, 2.66)	-
Pregabalin vs. Tamsulosin plus Dapoxetine	-2.6 (-12.76, 7.51)	-	-0.87 (-5.05, 3.46)	-1.72 (-4.89, 1.54)	-
Pregabalin vs. Tamsulosin plus Diclofenac	-0.88 (-10.82, 8.45)	-1.51 (-7.68, 5.21)	-0.57 (-4.46, 3.54)	0.31 (-2.76, 3.38)	-
Pregabalin vs. Tamsulosin plus Levofloxacin	-4.52 (-14.24, 5.62)	-5.41 (-11.14, 0.57)	-3.96 (-7.52, -0.17)*	-2.34 (-5.09, 0.59)	-
Profluss vs. Quercetin	-0.49 (-10.2, 8.67)	-	-	-	1.7 (0.03, 127.4)
Profluss vs. Saw palmetto	4.63 (-1.88, 11.08) DC: 4.6 (2.99, 6.21)*	-	-	-	0.77 (0.1, 5.36) DC: 1.28 (0.32, 5.06)
Profluss vs. Silodosin	0.14 (-9.97, 10.35)	-	-	-	5.65 (0.14, 274.4)
Profluss vs. Tamsulosin	0.59 (-9.44, 10.73)	-	-	-	1.83 (0.06, 58.54)
Profluss vs. Terazosin	1.38 (-9.1, 11.73)	-	-	-	2.26 (0.05, 82.54)
Profluss vs. Ciprofloxacin plus Solifenacin	-4.58 (-17, 7.61)	-	-	-	-
Profluss vs. Doxazosin plus Ibuprofen plus Thiocolchicoside	-8.87 (-19.61, 2.11)	-	-	-	2.6 (0.06, 108)
Profluss vs. Tamsulosin plus Ciprofloxacin	1.14 (-9.04, 12.01)	-	-	-	0.94 (0.03, 36.73)
Profluss vs. Tamsulosin plus Dapoxetine	-1.97 (-13.62, 9.98)	-	-	-	4.92 (0.11, 281.3)
Profluss vs. Tamsulosin plus Diclofenac	-0.36 (-12, 11.02)	-	-	-	-
Profluss vs. Tamsulosin plus	-3.85 (-15.15, 7.53)	-	-	-	1.47 (0.01, 147.9)

Levofloxacin					
Quercetin vs. Saw palmetto	5.12 (-1.97, 12.35)	3.46 (-4.03, 10.94)	-	-	0.45 (0.01, 18.21)
Quercetin vs. Silodosin	0.6 (-6.24, 8.04)	2.79 (-2.8, 8.48)	0.27 (-3.36, 4)	1.14(-1.8,4.21)	3.4 (0.11, 89.08)
Quercetin vs. Tamsulosin	1.1 (-5.76, 8.33)	1.92 (-3.57, 7.29)	-0.03 (-3.45, 3.44)	1.83(-0.89,4.67)	1.13 (0.05, 19.11)
Quercetin vs. Terazosin	1.9 (-5.69, 9.61)	2.06 (-4.44, 8.57)	-0.3 (-4.34, 3.95)	0.93(-2.21,4.28)	1.36 (0.05, 25.81)
Quercetin vs. Ciprofloxacin plus Solifenacin	-4.17 (-14.37, 6.7)	0.13 (-7.53, 7.88)	-1.79 (-6.79, 3.24)	0.68(-3.22,4.68)	-
Quercetin vs. Doxazosin plus Ibuprofen plus Thiocolchicoside	-8.36 (-16.13, -0.35)*	-1.07 (-7.58, 5.14)	-2.28 (-6.48, 1.8)	-1.33(-4.59,1.93)	1.58 (0.05, 36.59)
Quercetin vs. Tamsulosin plus Ciprofloxacin	1.68 (-5.89, 9.74)	2.58 (-2.93, 8.32)	0.32 (-3.45, 4.18)	1.65(-1.25,4.76)	0.58 (0.02, 11.41)
Quercetin vs. Tamsulosin plus Dapoxetine	-1.51 (-11.04, 8.39)	-	-0.41 (-4.79, 4.13)	0.01(-3.49,3.61)	3.04 (0.08, 115.3)
Quercetin vs. Tamsulosin plus Diclofenac	0.11 (-8.39, 9.27)	1.33 (-5.25, 8.03)	-0.05 (-4.27, 4.4)	2.07(-1.32,5.56)	-
Quercetin vs. Tamsulosin plus Levofloxacin	-3.41 (-12.06, 5.8)	-2.59 (-8.73, 3.42)	-3.43 (-7.32, 0.44)	-0.62(-3.59,2.68)	0.88 (0.01, 66.7)
Saw palmetto vs. Silodosin	-4.5 (-12.55, 3.33)	-0.77 (-7.5, 6.31)	-	-	7.45 (0.31, 222.3)
Saw palmetto vs. Tamsulosin	-4.01 (-11.47, 3.61)	-1.57 (-8.28, 5.28)	-	-	2.44 (0.14, 41.61)
Saw palmetto vs. Terazosin	-3.3 (-11.4, 4.88)	-1.48 (-9.09, 6.5)	-	-	2.86 (0.13, 59.78)
Saw palmetto vs. Ciprofloxacin plus Solifenacin	-9.22 (-19.61, 1.53)	-3.36 (-12.25, 5.24)	-	-	-
Saw palmetto vs. Doxazosin plus Ibuprofen plus Thiocolchicoside	-13.45 (-22.36, -5.02)*	-4.57 (-12.04, 3.25)	-	-	3.43(0.15,80.62)

Saw palmetto vs. Tamsulosin plus Ciprofloxacin	-3.52 (-11.36, 4.79)	-0.92 (-7.9, 6.26)	-	-	1.26 (0.06, 25.23)
Saw palmetto vs. Tamsulosin plus Dapoxetine	-6.54 (-16.14, 3.48)	-	-	-	6.55 (0.24, 229.1)
Saw palmetto vs. Tamsulosin plus Diclofenac	-4.99 (-14.28, 4.75)	-2.18 (-9.88, 5.95)	-	-	-
Saw palmetto vs. Tamsulosin plus Levofloxacin	-8.42 (-17.77,1.29)	-6.09 (-13.29, 1.08)	-	-	1.96 (0.02, 139.2)
Silodosin vs. Tamsulosin	0.48 (-6.17, 7.41)	-0.85 (-5.1, 3.49)	-0.34 (-3, 2.46)	0.68 (-1.43, 2.71)	0.34 (0.03, 2.82)
Silodosin vs. Terazosin	1.17 (-5.91, 8.39)	-0.74 (-6.4, 4.99)	-0.59 (-4.09, 2.99)	-0.22 (-2.98, 2.5)	0.4 (0.02, 4.29)
Silodosin vs. Ciprofloxacin plus Solifenacin	-4.79 (-15.03, 5.91)	-2.66 (-9.75, 4.35)	-2.1 (-6.58, 2.26)	-0.43 (-3.81, 2.78)	-
Silodosin vs. Doxazosin plus Ibuprofen plus Thiocolchicoside	-9.11 (-16.79, -1.27)*	-3.83 (-9.59, 1.42)	-2.59 (-6.24, 0.9)	-2.51 (-5.15, 0.06)	0.47 (0.03, 6.44)
Silodosin vs. Tamsulosin plus Ciprofloxacin	0.96 (-6.23, 8.63)	-0.19 (-4.83, 4.62)	-0.02 (-2.92, 3.16)	0.53 (-2, 2.84)	0.18 (0.01, 1.75)
Silodosin vs. Tamsulosin plus Dapoxetine	-2.18 (-11.59, 7.12)	-	-0.7 (-4.76, 3.29)	-1.13 (-4.04, 1.78)	0.9 (0.04, 20.68)
Silodosin vs. Tamsulosin plus Diclofenac	-0.61 (-8.93, 8.7)	-1.42 (-7.09, 4.31)	-0.37 (-3.89, 3.39)	0.9 (-1.92, 3.62)	-
Silodosin vs. Tamsulosin plus Levofloxacin	-4.04 (-12.78, 4.78)	-5.39 (-10.4, -0.29)*	-3.75 (-6.96, -0.36)*	-1.73 (-4.2, 0.65)	0.26 (0, 12.07)
Tamsulosin vs. Terazosin	0.71 (-6.33, 8.1)	0.13 (-5.11, 5.32)	-0.28 (-3.65, 3.03)	-0.93 (-3.45, 1.71)	1.19 (0.16, 8)
Tamsulosin vs. Ciprofloxacin plus Solifenacin	-5.25 (-14.2, 3.55)	-1.8 (-8.16, 4.28)	-1.79 (-5.62, 2.07)	-1.13 (-3.99, 1.79)	-
Tamsulosin vs.	-9.53 (-17.26, -1.93)*	-3 (-8.37, 2.08)	-2.25 (-5.79, 1.07)	-3.19 (-5.58, -0.79)*	1.43 (0.14, 12.02)

Doxazosin plus Ibuprofen plus Thiocolchicoside	DC: 0.09 (-3.28, 0.11)				
Tamsulosin vs. Tamsulosin plus Ciprofloxacin	0.52 (-4.13, 4.93)	0.68 (-2.42, 3.78) DC: 0.31 (-0.59, 1.21)	0.34 (-1.62, 2.31) DC: 0.53 (-0.25, 1.03)	-0.18 (-1.59, 1.29) DC: -0.34 (-0.92, 0.24)	0.52 (0.12, 2.32) DC: 1.3 (0.57, 2.94)
Tamsulosin vs. Tamsulosin plus Dapoxetine	-2.62 (-9.03, 3.62) DC: -2.6 (-3.65, -1.55)*	-	-0.39 (-3.24, 2.42) DC: -0.4 (-0.82, 0.02)	-1.82 (-3.95, 0.32) DC: -0.6 (-1.26, 0.06)	2.68 (0.43, 21.44) DC: 0.41 (0.12, 1.48)
Tamsulosin vs. Tamsulosin plus Diclofenac	-0.95 (-7, 5.09) DC: -1.2 (-2.78, 0.38)	-0.55 (-4.77, 3.8) DC: -0.8 (-1.83, 0.23)	-0.03 (-2.7, 2.62) DC: 0.1 (-0.48, 0.68)	0.22 (-1.67, 2.23) DC: -1.8 (-2.12, -1.48)*	-
Tamsulosin vs. Tamsulosin plus Levofloxacin	-4.5 (-13.11, 4.14)	-4.48 (-7.73, -1.43)* DC: -4.44 (-8.66, -0.23)*	-3.44 (-5.33, -1.31)* DC: -5.88 (-3.13, -0.39)*	-2.43 (-3.87, -0.95)* DC: -2.15 (-2.83, -1.46)*	0.82 (0.02, 19.74) DC: 1 (0.06, 16.5)
Terazosin vs. Ciprofloxacin plus Solifenacin	-6.03 (-16.36, 4.57)	-1.89 (-9.78, 5.9)	-1.54 (-6.43, 3.38)	-0.22 (-3.74, 3.49)	-
Terazosin vs. Doxazosin plus Ibuprofen plus Thiocolchicoside	-10.25 (-18.6, -1.88)*	-3.11 (-9.72, 3.22)	-1.95 (-6.15, 1.93)	-2.27 (-5.24, 0.68)	1.21 (0.11, 13.47)
Terazosin vs. Tamsulosin plus Ciprofloxacin	-0.26 (-7.87, 7.44)	0.55 (-5.15, 6.31)	0.62 (-3.1, 4.23)	0.75 (-2.06, 3.46)	0.43 (0.05, 3.89)
Terazosin vs. Tamsulosin plus Dapoxetine	-3.31 (-13.22, 6.25)	-	-0.11 (-4.66, 4.32)	-0.9 (-4.31, 2.28)	2.26 (0.15, 44.39)
Terazosin vs. Tamsulosin plus Diclofenac	-1.71 (-10.67, 7.21)	-0.66 (-7.16, 5.97)	0.23 (-3.93, 4.36)	1.11 (-2.05, 4.26)	-
Terazosin vs. Tamsulosin plus Levofloxacin	-5.19 (-11.34, 1) DC: -6.4 (-8.31, -4.49)*	-4.62 (-10.47, 1.33)	-3.16 (-6.87, 0.86)	-1.52 (-4.33, 1.34)	0.66 (0.01, 30.58)
Ciprofloxacin plus Solifenacin vs. Doxazosin plus Ibuprofen plus Thiocolchicoside	-4.19 (-15.42, 6.54)	-1.25 (-8.77, 6.76)	-0.5 (-5.3, 4.53)	-2.07 (-5.65, 1.39)	-
Ciprofloxacin plus Solifenacin vs.	5.73 (-3.38, 15.03)	2.51 (-3.8, 9.01)	2.12 (-1.97, 6.02)	0.95 (-1.99, 3.82)	-

Tamsulosin plus Ciprofloxacin					
Ciprofloxacin plus Solifenacin vs. Tamsulosin plus Dapoxetine	2.66 (-8.19, 13.42)	-	1.41 (-3.35, 6.13)	-0.67 (-4.22, 2.66)	-
Ciprofloxacin plus Solifenacin vs. Tamsulosin plus Diclofenac	4.31 (-5.94, 14.48)	1.25 (-5.96, 8.65)	1.75 (-2.69, 6.27)	1.32 (-1.97, 4.59)	-
Ciprofloxacin plus Solifenacin vs. Tamsulosin plus Levofloxacin	0.86 (-10.81, 12.14)	-2.71 (-9.16, 4.19)	-1.66 (-5.93, 2.76)	-1.31 (-4.5, 1.98)	-
Doxazosin plus Ibuprofen plus Thiocolchicoside vs. Tamsulosin plus Ciprofloxacin	10.02 (2.12, 18.3)*	3.68(-1.86,9.33)	2.63 (-1.07, 6.31)	3.03 (0.38, 5.67)*	0.36 (0.04, 3.86)
Doxazosin plus Ibuprofen plus Thiocolchicoside vs. Tamsulosin plus Dapoxetine	6.9 (-3.05, 16.96)	-	1.87 (-2.35, 6.38)	1.37 (-1.75, 4.6)	1.92 (0.11, 42.93)
Doxazosin plus Ibuprofen plus Thiocolchicoside vs. Tamsulosin plus Diclofenac	8.55 (-0.84, 18.33)	2.44 (-3.99, 8.9)	2.23 (-1.93, 6.32)	3.39 (0.34, 6.44)*	-
Doxazosin plus Ibuprofen plus Thiocolchicoside vs. Tamsulosin plus Levofloxacin	5.04 (-4.68, 14.48)	-1.47 (-7.37, 4.67)	-1.21 (-4.89, 2.76)	0.73 (-1.91, 3.53)	0.58 (0.01, 25.47)
Tamsulosin plus Ciprofloxacin vs. Tamsulosin plus Dapoxetine	-3.08 (-11.05, 4.5)	-	-0.74 (-4.11, 2.63)	-1.64 (-4.28, 0.98)	0.19 (0.02, 2.02)
Tamsulosin plus Ciprofloxacin vs. Tamsulosin plus Diclofenac	-1.48 (-7.37, 4.5) DC: -1.2 (-2.93, 0.53)	-1.22 (-5.5, 3.14) DC: -1 (-2.14, 0.14)	-0.38 (-3.1, 2.34) DC: -0.5(-1.07, 0.07)	0.38 (-1.65, 2.36) DC: -1.8 (-2.56, -1.04)*	-

Tamsulosin plus Ciprofloxacin vs. Tamsulosin plus Levofloxacin	-4.93 (-14.27, 3.94)	-5.18 (-9.59, -0.75)*	-3.78 (-6.41, -0.9)*	-2.26 (-4.22, -0.24)*	0.62 (0.02, 32.06)
Tamsulosin plus Dapoxetine vs. Tamsulosin plus Diclofenac	1.64 (-7.14, 10.32)	-	0.38 (-3.58, 4.1)	2.03 (-0.87, 4.94)	-
Tamsulosin plus Dapoxetine vs. Tamsulosin plus Levofloxacin	-1.94 (-12.53, 9.16)	-	-3.06 (-6.33, 0.56)	-0.62 (-3.21, 2.11)	0.3 (0, 12.92)
Tamsulosin plus Diclofenac vs. Tamsulosin plus Levofloxacin	-3.44 (-13.74, 6.52)	-3.96 (-9.49, 1.22)	-3.39 (-6.64, -0.06)*	-2.66 (-4.98, -0.21)	-

*Note: DC, direct comparison; -, not reported; *, Statistically significant.*

eTable 6: GRADE Summary of Findings

Patient or population: CP/CPPS patients
Interventions: oral pharmacological treatments
Comparator (reference): placebo
Outcome: NIH-CPSI total score
Setting(s): mixed intensive care unit setting

Total studies: 25 **Total Participants**:	**Relative effect * (95% CrI)**	**Anticipated absolute effect (95% CrI)**			**Certainty of the evidence**	**Number of participants**	**Ranking (%)**
		Placebo	**Intervention**	**Mean difference**			
Alfuzosin vs. Placebo	-	The MD was -6.5	The MD was -7.1	MD 0.6 lower (2.68 lower to 1.48 higher)	⨁⨁◯◯ **Low**	Direct evidence: 272 (1 trial)	20 (32.06)
Bromelain vs. Placebo	-	-	The MD was -3.1	MD 6.85 higher (0.98 lower to 14.74 higher)	⨁◯◯◯ **Very low**	No direct evidence. Indirect evidence only.	24 (3.28)
Celecoxib vs. Placebo	-	The MD was -4.8	The MD was -8.0	MD 3.2 lower (4.65 lower to 1.75 lower)	⨁⨁◯◯ **Low**	Direct evidence: 64 (1 trial)	10 (53.01)

Ciprofloxacin vs. Placebo	-	The MD was -3.4	The MD was -6.2	MD 2.8 lower (5.28 lower to 0.32 lower)	⊕⊕◯◯ **Low**	Direct evidence: 98 (1 trial)	8 (55.37)
Doxazosin vs. Placebo	-	The MD was -1.0	The MD was -12.4	MD 11.4 lower (11.93 lower to 10.87 lower)	⊕⊕◯◯ **Low**	Direct evidence: 53 (1 trial)	2 (93.97)
Dutasteride vs. Placebo	-	The MD was -2.7	The MD was -5.2	MD 2.5 lower (3.54 lower to 1.46 lower)	⊕⊕◯◯ **Low**	Direct evidence: 684 (1 trial)	15 (46.92)
Finasteride vs. Placebo	-	The MD was -0.8	The MD was -2.9	MD 2.1 lower (2.59 lower to 1.61 lower)	⊕⊕◯◯ **Low**	Direct evidence: 54 (1 trial)	7 (55.71)
Ibuprofen vs. Placebo	-	-	The MD was -6.2	MD 2.62 higher (5.27 lower to 10.45 higher)	⊕◯◯◯ **Very low**	No direct evidence. Indirect evidence only.	23 (15.00)
Levofloxacin vs. Placebo	-	The MD was -2.9	The MD was -5.4	MD 2.5 lower (4.98 lower to 0.02 lower)	⊕⊕◯◯ **Low**	Direct evidence: 80 (1 trial)	4 (67.87)

Pollen extract vs. Placebo	-	The MD was -5.2	The MD was -7.7	MD 2.5 lower (4.43 lower to 0.57 lower)	⊕⊕◯◯ **Low**	Direct evidence: 139 (1 trial)	6 (56.06)
Pentosan polysulfate sodium vs. Placebo	-	The MD was -3.2	The MD was -5.9	MD 2.7 lower (5.62 lower to 0.22 higher)	⊕⊕◯◯ **Low**	Direct evidence: 100 (1 trial)	13 (48.56)
Pregabalin vs. Placebo	-	The MD was -4.3	The MD was -6.5	MD 2.2 lower (3.42 lower to 0.98 lower)	⊕⊕◯◯ **Low**	Direct evidence: 324 (1 trial)	16 (44.30)
Profluss vs. Placebo	-	-	The MD was -6.8	MD 2,82 lower (11.78 lower to 6.06 higher)	⊕◯◯◯ **Very low**	No direct evidence. Indirect evidence only.	12 (49.81)
Quercetin vs. Placebo	-	The MD was -1.4	The MD was -8.0	MD 6.6 lower (9.82 lower to 3.38 lower)	⊕⊕◯◯ **Low**	Direct evidence: 30 (1 trial)	9 (54.46)
Saw palmetto vs. Placebo	-	-	The MD was 0.6	MD 1.78 higher (4.31 lower to 7.89 higher)	⊕◯◯◯ **Very low**	No direct evidence. Indirect evidence only.	22 (16.20)

Silodosin vs. Placebo	-	The MD was -8.5	The MD was -11.2	MD 2.66 lower (4.92 lower to 0.40 lower)	⊕⊕◯◯ **Low**	Direct evidence: 151 (1 trial)	14 (48.36)
Tamsulosin vs. Placebo	-	The MD was -3.7	The MD was -6.0	MD 2.48 lower (4.89 lower to 0.07 lower)	⊕⊕◯◯ **Low**	Direct evidence: 198 (2 trial)	17 (43.52)
Terazosin vs. Placebo	-	The MD was -10.2	The MD was -14.3	MD 4.1 lower (6.66 lower to 1.54 lower)	⊕⊕◯◯ **Low**	Direct evidence: 100 (1 trial)	19 (37.60)
Ciprofloxacin plus Solifenacin vs. Placebo	-	-	The MD was -7.7	MD 7.44 lower (16.27 lower to 1.65 higher)	⊕◯◯◯ **Very low**	No direct evidence. Indirect evidence only.	3 (78.54)
Doxazosin plus Ibuprofen plus Thiocolchicoside vs. Placebo	-	The MD was -1.0	The MD was -12.7	MD 11.7 lower (12.17 lower to 11.23 lower)	⊕⊕◯◯ **Low**	Direct evidence: 53 (1 trial)	1 (94.83)
Tamsulosin plus Ciprofloxacin vs. Placebo	-	The MD was -3.4	The MD was -4.1	MD 0.7 lower (2.91 lower to 1.51 higher)	⊕⊕◯◯ **Low**	Direct evidence: 98 (1 trial)	18 (39.04)

Tamsulosin plus Dapoxetine vs. Placebo	-	-	The MD was -14.3	MD 4.76 lower (12.64 lower to 3.08 higher)	⊕◯◯◯ **Very low**	No direct evidence. Indirect evidence only.	5 (64.39)
Tamsulosin plus Diclofenac vs. Placebo	-	-	The MD was -16.2	MD 3.17 lower (10.22 lower to 4.07 higher)	⊕◯◯◯ **Very low**	No direct evidence. Indirect evidence only.	11 (52.45)
Tamsulosin plus Levofloxacin vs. Placebo	-		-	MD 6.66 lower (13.91 lower to 0.66 higher)	⊕◯◯◯ **Very low**	No direct evidence. Indirect evidence only.	-
Placebo	Reference Comparator	-	-	-	-	-	21 (22.53)

Abbreviations: CP/CPPS = chronic prostatitis/chronic pelvic pain syndrome; NIH-CPSI = the National Institutes of Health Chronic Prostatitis Symptom Index; CrI = credible interval; MD = mean difference OR = odds ratio

NMA-SoF table definitions

* Network meta-analysis estimates are reported as odds ratio. CrI: credible interval (rather than confidence interval), since a Bayesian network meta-analysis has been conducted.

GRADE Working Group grades of evidence (or certainty in the evidence)


High quality: We are very confident that the true effect lies close to that of the estimate of the effect

Moderate quality: We are moderately confident in the effect estimate: The true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different


Low quality: Our confidence in the effect estimate is limited: The true effect may be substantially different from the estimate of the effect

Very low quality: We have very little confidence in the effect estimate: The true effect is likely to be substantially different from the estimate of effect


eFigure 1. The inconsistency assessment of network meta-analysis




(A) The inconsistency assessment for NIH-CPSI total scores




(B) The inconsistency assessment for NIH-CPSI pain scores



(C) The inconsistency assessment for NIH-CPSI urinary scores



(D) The inconsistency assessment for NIH-CPSI QoL scores



(E) The inconsistency assessment for adverse events