Overview PROMs HRQoL assessment IBM

Overview of applied PROMs for HRQoL assessment in the included quantitative studies

	Domains	Interpretation	Recommended by IMACS*
Generic subjective HRQoL			
36-Item Short-Form Health Survey (SF-36)	Physical functioning Role physical Bodily pain General health (=Physical Health Score, PCS) Vitality Social functioning Role-emotional Mental health (=Mental Health Score, MCS)	Range: 0-100 in two sub-scores (PCS, MCS); 100 is the best possible health status	x
12-Item Short-Form Health Survey (SF-12)	Physical functioning Role physical Bodily pain General health (=Physical Health Score, PCS) Vitality Social functioning Role-emotional Mental health (=Mental Health Score, MCS)	Range: 0-100 in two sub-scores (PCS, MCS); 100 is the best possible health status	
European quality of life 5D-5L questionnaire (EQ-5D-5L)	Mobility Self-care Usual activities Pain / discomfort Anxiety / depression	Index=1 is the best possible health status, Index<0 is the worst possible health status	
EQ Visual analogue scale (VAS)		Range 0-100 in one total score; 100 is the best possible health status	

	Domains	Interpretation	Recommended by IMACS*
Disease specific subjective HRQoL			
Individualized Neuromuscular Quality of Life questionnaire (INQoL)	Fatigue Weakness Pain Locking (myotonia) Emotion Social Independence Activity Body image Treatment effects and expectations INQoL index	Range 0-100 in 11 sub-scores and one total score; Higher percentage shows a greater symptom impact, respectively a worse health status	x
Subjective burden of mental illness			
Hospital Anxiety and Depression scale (HADS)	Anxiety and depression 14 Items, 7 each	Range: 0-21; Higher score indicates higher mental burden, Serious symptoms of anxiety or depression are indicated with a sub-score >8, Separate scores for depression and anxiety	
Beck Depression Inventory (BDI)	Depression 21 Items	Range 0-63; Higher score indicates higher depressive burden, Cut-off values: 0-10: no depression, clinically unremarkable or remitted 10-19: mild depressive syndrome 20-29: moderate depressive syndrome ≥ 30: severe depressive syndrome	

*IMACS: International Clinical Assessment and Studies Group; Rating according to Benveniste O, Rider LG. 213th ENMC International Workshop: Outcome measures and clinical trial readiness in idiopathic inflammatory myopathies, Heemskerk, The Netherlands, 18-20 September 2015. Neuromuscul Disord. 2016;26:523–34. doi:10.1016/j.nmd.2016.05.014.