




EBL
ETT
Mean Difference
Mean Difference
Study or Subgroup Mean SD Total Mean SD Total Weight IV, Random, 95% CI
IV, Random, 95% CI
7.7.1 higher risk
Abdelhalim 2014 0.1 0.7 20 2 3 20 5.8% −1.90 [−3.25, −0.55]
Elhendawy 2015 2.5 0.7 44 4.6 0.9 44 93.9% −2.10 [−2.44, −1.76]
Keohane 2013 3.5 6.1 8 9.3 17.1 15 0.1% −5.80 [−15.43, 3.83]
Subtotal (95% CI) 72 79 99.8% −2.09 [−2.42, −1.77]
Heterogeneity: Tau² = 0.00; Chi² = 0.65, df = 2 (P = 0.72); I² = 0%
Test for overall effect: Z = 12.55 (P < 0.00001)
7.7.2 lower risk
Sato 2011 2.4 0 12 2.7 0 22 Not estimable
Wells 2008 0.9 2 9 6.4 14.9 13 0.2% −5.50 [−13.70, 2.70]
Subtotal (95% CI) 21 35 0.2% −5.50 [−13.70, 2.70]
Heterogeneity: Not applicable
Test for overall effect: Z = 1.31 (P = 0.19)
Total (95% CI) 93 114 100.0% −2.10 [−2.42, −1.77]
Heterogeneity: Tau² = 0.00; Chi² = 1.31, df = 3 (P = 0.73); I² = 0%
Test for overall effect: Z = 12.60 (P < 0.00001)
Test for subgroup differences: Chi² = 0.66, df = 1 (P = 0.42), I² = 0%
−4
−2
0
2
4
Favours [EBL]
Favours [ETT]


Study or Subgroup	EBL Mean	EBL SD	EBL Total	ETT Mean	ETT SD	ETT Total	Weight	Mean Difference IV, Random, 95% CI
7.6.1 higher risk								
Abdelhalim 2014	2.4	1.4	20	1.4	0.9	20	22.6%	1.00 [0.27, 1.73]
Elhendawy 2015	3.6	0.2	44	3.1	0.2	44	39.9%	0.50 [0.42, 0.58]
Keohane 2013	3.1	0.6	8	1.9	0.4	15	30.8%	1.20 [0.74, 1.66]
Subtotal (95% CI)			72			79	93.5%	0.85 [0.32, 1.38]
Heterogeneity: $Tau^2 = 0.17$; $Chi^2 = 10.15$, df = 2 (P = 0.006); $I^2 = 80\%$								
Test for overall effect: Z = 3.14 (P = 0.002)								
7.6.2 lower risk								
Wells 2008	2.8	2.5	9	0.9	1.8	13	6.5%	1.90 [-0.00, 3.80]
Subtotal (95% CI)			9			13	6.5%	1.90 [-0.00, 3.80]
Heterogeneity: Not applicable								
Test for overall effect: Z = 1.96 (P = 0.05)								
Total (95% CI)			81			92	100.0%	0.92 [0.39, 1.45]
Heterogeneity: $Tau^2 = 0.16$; $Chi^2 = 12.14$, df = 3 (P = 0.007); $I^2 = 75\%$								
Test for overall effect: Z = 3.41 (P = 0.0006)								
Test for subgroup differences: $Chi^2 = 1.09$, df = 1 (P = 0.30), $I^2 = 7.9\%$								





EBL
ETT
Mean Difference
Mean Difference
Study or Subgroup Mean SD Total Mean SD Total Weight IV, Random, 95% CI
IV, Random, 95% CI
7.8.1 higher risk
Abdelhalim 2014 2.2 0.6 20 5.5 3.8 20 17.0% −3.30 [−4.99, −1.61]
Elhendawy 2015 2.9 0.8 44 3.5 0.9 44 26.3% −0.60 [−0.96, −0.24]
Keohane 2013 2.5 1 8 4.1 2.5 15 18.9% −1.60 [−3.04, −0.16]
Subtotal (95% CI) 72 79 62.2% −1.67 [−3.21, −0.12]
Heterogeneity: Tau² = 1.46; Chi² = 10.80, df = 2 (P = 0.005); I² = 81%
Test for overall effect: Z = 2.11 (P = 0.03)
7.8.2 lower risk
Sato 2011 3 0.5 12 2.3 0.4 22 26.4% 0.70 [0.37, 1.03]
Wells 2008 1.9 0.6 9 4.7 4.7 13 11.4% −2.80 [−5.38, −0.22]
Subtotal (95% CI) 21 35 37.8% −0.81 [−4.20, 2.59]
Heterogeneity: Tau² = 5.24; Chi² = 6.93, df = 1 (P = 0.008); I² = 86%
Test for overall effect: Z = 0.46 (P = 0.64)
Total (95% CI) 93 114 100.0% −1.15 [−2.30, −0.01]
Heterogeneity: Tau² = 1.26; Chi² = 52.23, df = 4 (P < 0.00001); I² = 92%
Test for overall effect: Z = 1.98 (P = 0.05)
Test for subgroup differences: Chi² = 0.20, df = 1 (P = 0.65), I² = 0%
−4
−2
0
2
4
Favours [EBL]
Favours [ETT]


Random sequence generation (selection bias)
Allocation concealment (selection bias)
Blinding of participants and personnel (performance bias)
Blinding of outcome assessment (detection bias)
Incomplete outcome data (attrition bias)
Selective reporting (reporting bias)
Other bias
0%
25%
50%
75%
100%
Low risk of bias
Unclear risk of bias
High risk of bias


A. Subgroup Analyses of Transfusion Requirements Based on Study Average Age, Sample Size, and Design



B. Subgroup Analyses of Transfusion Requirements Based on Interval Between Endoscopies



A. Subgroup Analyses of Hb Change Based on Study Average Age, Sample Size, and Design



B. Subgroup Analyses of Hb Change Based on Interval Between Endoscopies

