Annex 2

Publication risk of bias per disorder.

UI

Number of studies = 3 Root MSE = 0.6523

Std_Eff	Coef.	Std. Err.	t	*p* > |t|	[95% Conf. Interval]
slope	0.61339	0.100272	6.12	0.103	−0.6607–1.8875
bias	0.48164	0.766341	0.63	0.643	−9.2557–10.2189

Test of H0: no small-study effects *p* = 0.643

RESISTANCE FOR FUTURE NULL RESULTS (ROSENTHAL model)

The number of additional non-significant studies that are necessary to add a significant meta-analysis for becoming non-significant:73.

The number of unpublished studies 73 is >25 (k*5 + 10).

POP

Number of studies = 4 Root MSE = 1013

Std_Eff	Coef.	Std. Err.	t	p > |t|	[95% Conf. Interval]
slope	1.89124	0.353177	5.35	0.033	0.3716–3.4109
bias	−2.77292	1.319602	−2.10	0.170	−8.4508–2.9049

Test of H0: no small-study effects p = 0,170

RESISTANCE FOR FUTURE NULL RESULTS (ROSENTHAL model)

The number of additional non-significant studies that are necessary to add a significant meta-analysis for becoming non-significant:99.

The number of unpublished studies 99 is > 30 (k*5 + 10).

AI

Number of studies = 4 Root MSE = 1,618

Std_Eff	Coef.	Std. Err.	t	p > |t|	[95% Conf. Interval]
slope	0.52553	0.119651	4.39	0.048	0.0107–1.0403
bias	−1.34033	1.343679	−1.00	0,424	−7.1217–4.4410

Test of H0: no small-study effects p = 0.424

RESISTANCE FOR FUTURE NULL RESULTS (ROSENTHAL model)

The number of additional non-significant studies that are necessary to add a significant meta-analysis for becoming non-significant:65.

The number of unpublished studies 65 is >30 (k*5 + 10).