The World Journal of MEN's HEALTH

Supplement Table 3. Characteristics of ongoing studies

	NCT02321670
Study name	Scandinavian urethroplasty study. A multicentre, prospective, randomized study comparing bulbar urethroplasty with excision and primary anastomosis or with an onlay grafting procedure using buccal mucosa.
Methods	Study design: randomised parallel controlled trial (open label) Study dates: September 2015 to December 2020 Setting: inpatient, multinational/multicentre Country: Finland, Norway, Sweden
Participants	Inclusion criteria: 1. Male 18–70 years old; 2. Participant with bulbar urethral stricture, length estimated by urethrography or endoscopy ≤2 cm; 3. Participant is able and willing to sign informed consent; 4. Participant is able and willing to complete all study requirements Exclusion criteria: 1. Previous open urethroplasty; 2. Previous hypospadia surgery; 3. Previous surgery for congenital curvature or Mb Peyronie; 4. Previous pelvic irradiation therapy 5. Known grave psychiatric disorder; 6. Haemophilia or other clotting disorders that cause bleeding diathesis; 7. Use of medication to increase erectile function, such as PDE5-inhibitors and intracavernous injections, during the study; 8. Any condition or situation, which, in the investigator's opinion, puts the patient at significant risk, could confound the study results, or may interfere significantly with the patient's participation in the study Estimated enrollment: 300 participants
Interventions	Intervention: Excision of the stricture and end-to-end anastomosis of the urethra. Control: Incision of stricture and grafting procedure with buccal mucosa where the corpus spongiosum is not divided
Outcomes	Primary outcomes • Degree of erectile dysfunction after surgery (time frame: up to one year) measured by changes in IIEF-5 • Penile complications (time frame: up to one year) measured by questionnaire made for this study Secondary outcomes • Occurrence of failure within the follow up period (time frame: one year) • Occurrence of complications (time frame: from operation to one year)
Starting date	September 2015
Contact information	Ole Jacob Nilsen, MD; +47 23070000; ojnilsen@ous-hf.no Alexander Schultz, MD; +47 23070000; aschultz@ous-hf.no
Notes	Sponsor: Oslo University Hospital