

Experimental
Control
Std. Mean Difference
Std. Mean Difference
Study or Subgroup Mean SD Total Mean SD Total Weight IV, Fixed, 95% CI
IV, Fixed, 95% CI
1.7.1 perennial allergic rhinitis and seasonal allergic rhinitis
Giovannini2007 24 319.837 91 91.7 387.3103 92 27.8% -0.19 [-0.48, 0.10]
Subtotal (95% CI) 91 92 27.8% -0.19 [-0.48, 0.10]
Heterogeneity: Not applicable
Test for overall effect: Z = 1.28 (P = 0.20)
1.7.2 perennial allergic rhinitis
Ishida 2005 -36.7 780.4736 25 -72.6 712.4608 24 7.5% 0.05 [-0.51, 0.61]
Jan 2011 61.39 284.82 52 -150.5 632.76 51 15.4% 0.43 [0.04, 0.82]
Lin 2013 40.34 392.55 55 150.5 632.76 51 16.1% -0.21 [-0.59, 0.17]
Lue 2012 14.25 216.42 30 8.04 197.89 27 8.7% 0.03 [-0.49, 0.55]
Nishimura 2009(low dose) -19 278.9358 15 -38 457.7827 15 4.6% 0.05 [-0.67, 0.76]
Nishimura 2009(high dose) -101 2,709.536 15 -38 457.7827 15 4.6% -0.03 [-0.75, 0.68]
Subtotal (95% CI) 192 183 56.8% 0.07 [-0.13, 0.27]
Heterogeneity: Chi² = 5.43, df = 5 (P = 0.37); I² = 8%
Test for overall effect: Z = 0.67 (P = 0.50)
1.7.3 seasonal allergic rhinitis
Nagata 2010 7.2 760.0199 9 -57.5 478.4282 10 2.9% 0.10 [-0.80, 1.00]
Xiao 2006a 8.7 61.93489 20 40.6 278.5341 20 6.1% -0.15 [-0.78, 0.47]
Xiao 2006b 30.3 126.8871 22 52.2 234.1578 20 6.4% -0.12 [-0.72, 0.49]
Subtotal (95% CI) 51 50 15.4% -0.09 [-0.48, 0.30]
Heterogeneity: Chi² = 0.22, df = 2 (P = 0.90); I² = 0%
Test for overall effect: Z = 0.46 (P = 0.65)
Total (95% CI) 334 325 100.0% -0.03 [-0.18, 0.13]
Heterogeneity: Chi² = 7.81, df = 9 (P = 0.55); I² = 0%
Test for overall effect: Z = 0.35 (P = 0.72)
Test for subgroup differences: Chi² = 2.17, df = 2 (P = 0.34), I² = 7.7%
-2
-1
0
1
2
Favours [experimental]
Favours [control]