
In this file are included in total (n=6) supplementary material tables and (n=2) supplementary material figures.

Supplementary material Table 1 Search syntax used in the different databases

Database	Search terms	Hits
PubMed	("amyloidosis"[All Fields] OR "amyloidosis"[Mesh] OR "amyloid"[All Fields]) AND ("prealbumin"[MeSH Terms] OR "prealbumin"[All Fields] OR "TTR"[All Fields] OR "ATTR"[All Fields] OR "ATTRwt"[All Fields] OR "transthyretin"[All Fields]) AND ("cardiomyopathies"[MeSH Terms] OR "cardiomyopathies"[All Fields] OR "cardiomyopathy"[All Fields] OR ("heart failure"[MeSH Terms] OR ("heart"[All Fields] AND "failure"[All Fields]) OR "heart failure"[All Fields] OR "cardiac failure"[All Fields]))	846
EMBASE	(amyloidosis OR 'amyloidosis'/exp OR amyloid) AND ('prealbumin'/exp OR prealbumin OR ttr OR attr OR attrwt OR transthyretin) AND ('cardiomyopathies'/exp OR cardiomyopathies OR cardiomyopathy OR ((((heart AND 'failure'/exp OR heart) AND failure OR heart) AND failure OR cardiac) AND failure))	2276
CENTRAL	(amyloidosis OR amyloid) AND (prealbumin OR ttr OR attr OR attrwt OR transthyretin) AND (cardiomyopathies OR cardiomyopathy OR (heart AND failure OR heart failure OR cardiac failure))	79

Supplementary material Figure 1 PRISMA flow diagram



*Tanskanen et al. 2008 was added after search of the references of the included studies

Supplementary material Table 2 List of included studies

#	Title	Author (year)
1	First Australian registry of patients with wild type cardiac amyloidosis (ATTRwt): Queensland experience	Burrage M., et al. 2019
2	Senile cardiac amyloidosis	Wilson et al. 2010
3	Cardiac Amyloid Imaging with 18F-Florbetaben PET: A Pilot Study.	Law et al. 2016
4	Mechanisms of heart failure in transthyretin vs. light chain amyloidosis.	Binder et al . 2019
5	When and how do patients with cardiac amyloidosis die?	Escher et al. 2019
6	Riociguat-new therapeutic option in the management of cardiac amyloidosis	Franz et al.2016
7	Abnormal Coronary Flow Velocity Reserve and Decreased Myocardial Contractile Reserve Are Main Factors in Relation to Physical Exercise Capacity in Cardiac Amyloidosis.	Clemmense n et al. 2017
8	Diagnostic delay in wild type transthyretin cardiac amyloidosis – A clinical challenge	Ladefoged et al. 2020
9	Apical sparing pattern of left ventricular myocardial (99m)Tc-HMDP uptake in patients with transthyretin cardiac amyloidosis	Van Der Gucht et al. 2017
10	Causes and Consequences of Longitudinal LV Dysfunction Assessed by 2D Strain Echocardiography in Cardiac Amyloidosis	Ternacle et al. 2016
11	Atrial fibrillation and subtype of atrial fibrillation in cardiac amyloidosis: clinical and echocardiographic features, impact on mortality	Sanchis et al. 2019
12	Identification of prognostic markers in transthyretin and AL cardiac amyloidosis	Damy et al. 2016
13	Pharyngo-laryngeal involvement in systemic amyloidosis with cardiac involvement: a prospective observational study	Bartier et al. 2019
14	Pilot study for left ventricular imaging phenotype of patients over 65 years old with heart failure and preserved ejection fraction: the high prevalence of amyloid cardiomyopathy	Bennani et al. 2016
15	Prevalence, Severity, and Prognostic Value of Sleep Apnea Syndromes in Cardiac Amyloidosis	Bodez et al. 2016

#	Title	Author (year)
16	Prognostic value of right ventricular systolic function in cardiac amyloidosis	Bodez et al. 2016
17	Renal Infarction and Its Consequences for Renal Function in Patients With Cardiac Amyloidosis	Dang et al. 2019
18	Usefulness of (99m)Tc-HMDP scintigraphy for the etiologic diagnosis and prognosis of cardiac amyloidosis.	Galat et al. 2015
19	Comparison of different types of cardiac amyloidosis by cardiac magnetic resonance imaging.	Kristen et al. 2015
20	Carpal tunnel syndrome and spinal canal stenosis: harbingers of transthyretin amyloid cardiomyopathy?	aus dem Siepen et al. 2019
21	Genotype and phenotype heterogeneity of transthyretin-associated amyloidosis: A report from the German Amyloidosis referral center	Purrucker et al. 2014
22	Green tea extract as a treatment for patients with wild-type transthyretin amyloidosis: an observational study.	aus dem Siepem et al. 2015
23	Non-invasive risk stratification of patients with transthyretin amyloidosis.	Kristen et al. 2014
24	Predictors of survival stratification in patients with wild-type cardiac amyloidosis.	aus dem Siepen et al. 2018
25	Risk stratification in wild-type transthyretin amyloidosis	Kristen et al. 2016
26	Skeletal scintigraphy indicates disease severity of cardiac involvement in patients with senile systemic amyloidosis	Kristen et al. 2011
27	Coexistence of Degenerative Aortic Stenosis and Wild-Type Transthyretin-Related Cardiac Amyloidosis	Longhi et al. 2015
28	Disease profile and differential diagnosis of hereditary transthyretin-related amyloidosis with exclusively cardiac phenotype: an Italian perspective.	Rapezzi et al. 2013
29	Atrial fibrillation in amyloidotic cardiomyopathy: prevalence, incidence, risk factors and prognostic role.	Longhi et al. 2015
30	Different NT-proBNP circulating levels for different types of cardiac amyloidosis.	Perfetto et al. 2016

#	Title	Author (year)
31	Gender-related risk of myocardial involvement in systemic amyloidosis.	Rapezzi et al. 2008
32	Phenotypic profile of Ile68Leu transthyretin amyloidosis: an underdiagnosed cause of heart failure	Gagliardi et al. 2018
33	Systemic cardiac amyloidoses: disease profiles and clinical courses of the 3 main types.	Rapezzi et al. 2009
34	The etiology of amyloidosis influences the pathophysiology and outcome of heart failure in amyloidotic cardiomyopathy	Quarta et al. 2013
35	Usefulness and limitations of (99m)Tc-3, 3NRdiphosphono-1, 2-propanodicarboxylic acid scintigraphy in the aetiological diagnosis of amyloidotic cardiomyopathy	Rapezziet al. 2011
36	Comparison of the standard and speckle tracking echocardiographic features of wild-type and mutated transthyretin cardiac amyloidoses.	Minamisawa et al. 2016
37	Carpal tunnel syndrome: a common initial symptom of systemic wild-type ATTR (ATTRwt) amyloidosis.	Nakagawa et al. 2016
38	First nationwide survey on systemic wild-type ATTR amyloidosis in Japan	Sekijima et al. 2017
39	High prevalence of ATTR amyloidosis in endomyocardial biopsy-proven cardiac amyloidosis patients	Nakagawa et al. 2013
40	Clinicopathological features of senile systemic amyloidosis: an ante- and post-mortem study.	Ueda et al. 2011
41	Diagnostic utility of cardiac troponin T level in patients with cardiac amyloidosis.	Takashio et al. 2017
42	Wild-type ATTR amyloidosis may be associated with unexpected death among the elderly.	Shiozaki et al. 2019
43	Reduced trans-mitral A-wave velocity predicts the presence of wild-type transthyretin amyloidosis in elderly patients with left ventricular hypertrophy	Yamamura et al. 2017
44	Characterization of wild-type transthyretin amyloidosis among women: Preliminary results from an international multicenter study	Quarta et al. 2017
45	Clinical characteristics of wild-type transthyretin cardiac amyloidosis: disproving myths.	Gonzalez et al. 2017
46	Description of the health-related quality of life of transthyretin amyloid cardiomyopathy subjects from the transthyretinamyloidosis outcomes survey	Grogan et al. 2017

#	Title	Author (year)
47	Impact of genotype and phenotype on cardiac biomarkers in patients with transthyretin amyloidosis - Report from the Transthyretin Amyloidosis Outcome Survey (THAOS)	Kristen et al. 2017
48	Left atrial structure and function in cardiac amyloidosis.	Nochioka et al. 2017
49	Left Ventricular Structure and Function in Transthyretin-Related Versus Light-Chain Cardiac Amyloidosis	Quarta et al. 2014
50	Nonbiopsy Diagnosis of Cardiac Transthyretin Amyloidosis.	Gillmore et al. 2016
51	Features of transthyretin cardiomyopathy patients with 6 specific cardiac genotypes in the THAOS registry	Schumacher et al. 2015
52	Survival following heart transplantation in patients with transthyretin amyloid cardiomyopathy	Selby et al. 2014
53	Tafamidis Treatment for Patients with Transthyretin Amyloid Cardiomyopathy	Maurer et al. 2018
54	THAOS - The Transthyretin Amyloidosis Outcomes Survey: initial report on clinical manifestations in patients with hereditary and wild-type transthyretin amyloidosis.	Coelho et al. 2013
55	Transthyretin cardiac amyloidosis in continental Western Europe: an insight through the Transthyretin Amyloidosis Outcomes Survey (THAOS).	Damy et al. 2019
56	Prevalence of Positive 99 mTc-DPD Scintigraphy as an Indicator ofthe Prevalence of Wild-type Transthyretin Amyloidosisin the Elderly	Hyue et al. 2019
57	Wild-type transthyretin amyloidosis as a cause of heart failure with preserved ejection fraction.	Gonzalez et al. 2015
58	A new staging system for cardiac transthyretin amyloidosis.	Gillmore et al. 2017
59	Clinical utility of T1 mapping in cardiac ATTR amyloidosis-diagnostic performance and prognostic capability	Martinez-Naharro et al. 2017
60	Differential Myocyte Responses in Patients with Cardiac Transthyretin Amyloidosis and Light-Chain Amyloidosis: A Cardiac MR Imaging Study.	Fontana et al. 2015
61	Genetic diagnosis in ATTR amyloidosis; A single UK centre 26 year experience	Rowczenio et al. 2017

#	Title	Author (year)
62	Native T1 and Extracellular Volume in Transthyretin Amyloidosis	Martinez et al. 2019
63	Native T1 fails to predict outcome in ATTR amyloidosis but ECV does	Norrington et al. 2017
64	Natural History, Quality of Life, and Outcome in Cardiac Transthyretin Amyloidosis	Thirusha et al. 2019
65	Echocardiographic phenotype and prognosis in transthyretin cardiac amyloidosis	Chacko et al. 2020
66	Senile Systemic Amyloidosis: Clinical Features at Presentation	Pinney et al. 2013
67	Spotting senile systemic amyloidosis: Why we miss it	Fontana et al. 2015
68	Transthyretin amyloidosis cardiomyopathy: Genotype, phenotype, quality of life and outcome in 997 patients in the UK	Lane et al. 2017
69	Utility and limitations of 3,3-diphosphono-1, 2-propanodicarboxylic acid scintigraphy in systemic amyloidosis	Hutt et al. 2014
70	A comparison of electrocardiographic findings among the three main types of cardiac amyloid	Cyrille et al. 2013
71	Association between ruptured distal biceps tendon and wild-type transthyretin cardiac amyloidosis	Geller et al. 2017
72	Comparison of cardiac amyloidosis due to wild-type and V122I transthyretin in older adults referred to an academic medical center	Givens et al. 2013
73	Diflunisal is associated with improved mortality in patients with wild-type attr cardiac amyloidosis: The bu amyloidosis center experience	Mints et al. 2019
74	Diflusinal tolerability in transthyretin cardiac amyloidosis: a single center's experience	Ikram et al. 2018
75	Genetic variation of the transthyretin gene in wild-type transthyretin amyloidosis (ATTRwt)	Sikora et al. 2015
76	Genotype and Phenotype of Transthyretin Cardiac Amyloidosis: THAOS (Transthyretin Amyloid Outcome Survey).	Maurer et al. 2016
77	Heart Failure Resulting From AgeNRRelated Cardiac Amyloid Disease Associated With WildNRType Transthyretin: A Prospective, Observational Cohort Study.	Connors et al. 2016

#	Title	Author (year)
78	Left Ventricular Amyloid Deposition in Patients with Heart Failure and Preserved Ejection Fraction	Mohammed et al. 2014
79	Machine learning predicts mortality better than biomarker staging in wild-type transthyretin cardiac amyloidosis	Singh et al. 2017
80	Matrix metalloproteinases and their tissue inhibitors in cardiac amyloidosis: relationship to structural, functional myocardial changes and to light chain amyloid deposition.	Biolo et al. 2008
81	Monoclonal gammopathy of undetermined significance in systemic transthyretin amyloidosis (ATTR).	Phull et al. 2018
82	Natural History of Wild-Type Transthyretin Cardiac Amyloidosis and Risk Stratification Using a Novel Staging System.	Grogan et al. 2016
83	Outcomes After Cardiac Transplant for Wild Type Transthyretin Amyloidosis.	Rosenbaum et al. 2018
84	Prospective evaluation of the morbidity and mortality of wild-type and V122I mutant transthyretin amyloid cardiomyopathy: the Transthyretin Amyloidosis Cardiac Study (TRACS).	Ruberg et al. 2012
85	Ruptured biceps tendon: A novel non-cardiac clue to TTR cardiac amyloidosis	Geller et al. 2016
86	Senile systemic amyloidosis presenting with heart failure: a comparison with light chain-associated amyloidosis.	Ng et al. 2005
87	Stabilization of Cardiac Function With Diflunisal in Transthyretin (ATTR) Cardiac Amyloidosis.	Lohrmann et al. 2019
88	Clinical features and survival in senile systemic amyloidosis: comparison to familial transthyretin cardiomyopathy.	Connors et al. 2011
89	Tafamidis in Transthyretin Amyloid Cardiomyopathy	Maurer et al. 2015
90	The myocardial contraction fraction is superior to ejection fraction in predicting survival in patients with AL cardiac amyloidosis.	Tendler et al. 2014
91	The premortem recognition of systemic senile amyloidosis with cardiac involvement.	Kyle et al. 1996
92	The prognostic significance of central hemodynamics in patients with cardiac amyloidosis.	Russo et al. 2013

#	Title	Author (year)
93	Hip and knee arthroplasty are common among patients with transthyretin cardiac amyloidosis, occurring years before cardiac amyloid diagnosis: can we identify affected patients earlier?	Rubin J, et al. 2017
94	Safety and efficacy of a TTR specific antisense oligonucleotide in patients with transthyretin amyloid cardiomyopathy.	Benson et al. 2017
95	Subtype-Specific Interactions and Prognosis in Cardiac Amyloidosis.	Sperry et al. 2016
96	Use of Ventilatory Efficiency Slope as a Marker for Increased Mortality in Wild-Type Transthyretin Cardiac Amyloidosis	Yunis et al. 2019
97	Utility of cardiac transplant for wild type transthyretin amyloidosis	Rosenbaum et al. 2017
98	Val122Ile familial transthyretin (TTR) amyloid cardiomyopathy in African-American patients has a worse prognosis than wild-type TTR amyloid cardiomyopathy	Falk et al. 2016
99	Transthyretin Cardiac Amyloidosis as Diagnosed by 99mTc-PYP Scanning in Patients with Acute Heart Failure and Preserved Ejection Fraction.	Lo Presti et al. 2019
100	Treatment With Tafamidis Slows Disease Progression in Early-Stage Transthyretin Cardiomyopathy.	Sultan et al. 2017
101	Use of Serum Transthyretin as a Prognostic Indicator and Predictor of Outcome in Cardiac Amyloid Disease Associated With Wild-Type Transthyretin.	Hanson et al. 2018
102	Wild type transthyretin cardiac amyloidosis: Clinical characteristics, echocardiographic findings, and predictors of outcome	Grogan et al. 2015
103	Yield of noncardiac biopsy for the diagnosis of transthyretin cardiac amyloidosis.	Fine et al. 2014
104	Senile systemic amyloidosis affects 25% of the very aged and associates with genetic variation in alpha2-macroglobulin and tau: A population-based autopsy study	Tanskanen et al. 2008
105	Occult Transthyretin Cardiac Amyloid in Severe Calcific Aortic Stenosis: Prevalence and Prognosis in Patients Undergoing Surgical Aortic Valve Replacement.	Treibel et al. 2016

Supplementary material Table 3 Characteristics of identified unique studies

#	Title	Publication year	Year of data collection	Country	Prospective	Autopsy	Mean age	Total patients (n)	Male (%)	Atrial Fibrillation (%))	CTS (%)
1	First Australian registry of patients with wild type cardiac amyloidosis (ATTRwt): Queensland experience	2019	2011 - 2016	Australia	No	No	77	35	100	37.5	NR
2	Senile cardiac amyloidosis	2010	2005 - 2008	Australia	No	No	76	6	100	83.3	NR
3	Mechanisms of heart failure in transthyretin vs. light chain amyloidosis.	2019	NR - 2016	Austria	Yes	No	NA	62	82.3	58.8	NR
4	When and how do patients with cardiac amyloidosis die?	2019	2000 - 2018	Austria	No	Yes	74.6	48	91.7	50	NR
5	Abnormal Coronary Flow Velocity Reserve and Decreased Myocardial Contractile Reserve Are	2017	2014 - 2016	Denmark	Yes	No	78	10	100	NR	NR

#	Title	Publication year	Year of data collection	Country	Prospective	Autopsy	Mean age	Total patients (n)	Male (%)	Atrial Fibrillation (%))	CTS (%)
6	Main Factors in Relation to Physical Exercise Capacity in Cardiac Amyloidosis. Diagnostic delay in wild type transthyretin cardiac amyloidosis – A clinical challenge	2020	2017 - 2019	Denmark	No	No	81.7	50	88	NR	36
7	Atrial fibrillation and subtype of atrial fibrillation in cardiac amyloidosis: clinical and echocardiographic features, impact on mortality	2019	2008 - 2018	France	No	No	81	97	85.6	NR	NR
8	Identification of prognostic markers in transthyretin and AL cardiac amyloidosis	2016	2005 - 2016	France	Yes	No	NA	23	78	40	48

#	Title	Publication year	Year of data collection	Country	Prospective	Autopsy	Mean age	Total patients (n)	Male (%)	Atrial Fibrillation (%))	CTS (%)
9	Pharyngo-laryngeal involvement in systemic amyloidosis with cardiac involvement: a prospective observational study	2019	2016 - 2018	France	Yes	No	80	40	77.5	NR	NR
10	Carpal tunnel syndrome and spinal canal stenosis: harbingers of transthyretin amyloid cardiomyopathy?	2019	2005 - 2016	Germany	No	No	74	253	92	NR	62
11	Atrial fibrillation in amyloidotic cardiomyopathy: prevalence, incidence, risk factors and prognostic role.	2015	2000 - 2012	Italy	No	No	76.9	45	84.4	NR	NR
12	Different NT-proBNP circulating levels for different types of cardiac	2016	2006 - 2013	Italy	No	No	81	18	83.3	33	NR

#	Title	Publication year	Year of data collection	Country	Prospective	Autopsy	Mean age	Total patients (n)	Male (%)	Atrial Fibrillation (%))	CTS (%)
	amyloidosis. Carpal tunnel syndrome: a common initial symptom of systemic wild-type ATTR (ATTRwt) amyloidosis.										
13	Clinicopathological features of senile systemic amyloidosis: an ante- and post-mortem study.	2016	2008 - 2015	Japan	No	No	74.5	31	77.4	42	54.8
14	Diagnostic utility of cardiac troponin T level in patients with cardiac amyloidosis.	2011	2001 - 2008 (post-mortem); 2005 - 2010 (ante-mortem)	Japan	No	Yes	NA	17	88.2	NR	NR
15	Wild-type ATTR amyloidosis may be associated with unexpected death among the elderly.	2017	2010 - 2016	Japan	No	No	77	40	92.5	53	NR
16		2019	2006 - 2018	Japan	No	Yes	NA	32	43.75	NR	NR

#	Title	Publication year	Year of data collection	Country	Prospective	Autopsy	Mean age	Total patients (n)	Male (%)	Atrial Fibrillation (%))	CTS (%)
17	Prevalence of Positive 99 mTc-DPD Scintigraphy as an Indicator of the Prevalence of Wild-type Transthyretin Amyloidosis in the Elderly	2019	2014 - 2017	South Korea	No	No	80.7	6	83.33	NR	NR
18	Wild-type transthyretin amyloidosis as a cause of heart failure with preserved ejection fraction.	2015	NR	Spain	Yes	No	86	16	50	NR	NR
19	Echocardiographic phenotype and prognosis in transthyretin cardiac amyloidosis	2020	2000 - 2019	UK	Yes	No	77.5	766	94.52	NR	NR
20	Clinical features and survival in senile systemic amyloidosis: comparison to familial	2011	1994 - 2009	US	No	No	NA	82	96.34	NR	6

#	Title	Publication year	Year of data collection	Country	Prospective	Autopsy	Mean age	Total patients (n)	Male (%)	Atrial Fibrillation (%))	CTS (%)
	transthyretin cardiomyopathy. Hip and knee arthroplasty are common among patients with transthyretin cardiac amyloidosis, occurring years before cardiac amyloid diagnosis: can we identify affected patients earlier? Safety and efficacy of a TTR specific antisense oligonucleotide in patients with transthyretin amyloid cardiomyopathy. Subtype-Specific Interactions and Prognosis in Cardiac Amyloidosis. Transthyretin Cardiac										
21		2017	NR	US	Yes	No	75	108	92.59	NR	59.3
22		2017	NR	US	Yes	No	NA	7	100	NR	NR
23		2016	2002 - 2014	US	No	No	75.9	75	88	77	NR
24		2019	2016 - 2018	US	No	No	82	19	84.21	63	NR

#	Title	Publication year	Year of data collection	Country	Prospective	Autopsy	Mean age	Total patients (n)	Male (%)	Atrial Fibrillation (%))	CTS (%)
25	Amyloidosis as Diagnosed by 99mTc-PYP Scanning in Patients with Acute Heart Failure and Preserved Ejection Fraction. Treatment With Tafamidis Slows Disease Progression in Early-Stage Transthyretin Cardiomyopathy. Use of Serum Transthyretin as a Prognostic Indicator and Predictor of Outcome in Cardiac Amyloid Disease Associated With Wild-Type Transthyretin.	2017	2008 - 2010	US	Yes	No	76.4	49	95.92	NR	NR
26	Wild type transthyr	2018	2009 - 2016	US	No	No	NA	116	97.41	NR	NR
27		2015	1968-2013	US	No	Yes	77	428	88.32	NR	NR

#	Title	Publication year	Year of data collection	Country	Prospective	Autopsy	Mean age	Total patients (n)	Male (%)	Atrial Fibrillation (%))	CTS (%)
	etin cardiac amyloidosis: Clinical characteristics, echocardiographic findings, and predictors of outcome										
28	Senile systemic amyloidosis affects 25% of the very aged and associates with genetic variation in alpha2-macroglobulin and tau: A population-based autopsy study	2008	1991-1999	Finland	No	Yes	93.8	63	22.2	NR	NR

Supplementary material Table 4 Quality assessment of individual studies

Checkpoint → Study ↓	1	2	3	6	7	9	10	11	12	13	21	22
Burrage et al.	Y	N	Y	Y	Y	Y	N	UD	UD	Y	Y	UD
Wilson et al.	Y	Y	Y	Y	N	Y	N	Y	Y	Y	UD	UD
Binder et al.	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
Escher et al.	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	N	Y
Clemmensen et al.	Y	Y	Y	Y	Y	N	N	Y	Y	Y	Y	Y
Ladefoged et al.	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
Sanchis et al.	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
Damy et al.	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	N	Y
Bartier et al.	Y	Y	Y	Y	Y	N	Y	Y	Y	Y	Y	Y
Aus dem Siepen et al.	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
Longhi et al.	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
Perfetto et al.	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
Nakagawa et al.	N	Y	N	Y	Y	Y	Y	Y	Y	UD	Y	Y
Ueda et al.	N	Y	Y	Y	N	Y	N	Y	UD	UD	Y	Y
Takashio et al.	Y	Y	Y	Y	Y	Y	N	Y	UD	Y	Y	Y
Shiozaki et al.	Y	Y	Y	Y	N	Y	N	Y	UD	Y	Y	Y
Hyue et al.	N	Y	N	Y	N	Y	N	UD	UD	Y	Y	UD

Checkpoint → Study ↓	1	2	3	6	7	9	10	11	12	13	21	22
Gonzalez et al.	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
Chacko et al.	Y	Y	Y	Y	Y	N	Y	Y	Y	Y	Y	Y
Connors et al.	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
Rubin et al.	Y	Y	Y	Y	Y	N	Y	Y	UD	Y	Y	Y
Benson et al.	Y	Y	Y	Y	Y	N	Y	UD	Y	UD	Y	UD
Sperry et al.	Y	Y	Y	Y	Y	Y	Y	Y	N	Y	Y	Y
Lo Presti et al.	Y	Y	Y	Y	Y	Y	Y	Y	UD	Y	Y	Y
Sultan et al.	Y	Y	Y	Y	Y	N	Y	UD	UD	UD	N	UD
Hanson et al.	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
Grogan et al.	Y	Y	Y	Y	N	Y	N	Y	UD	Y	Y	UD
Tanskanen et al.	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y

Abbreviations: Y: Yes, N: No, UD: Unable to determine

Supplementary material Table 5 Downs and Black checklist

#	Item
1	Is the hypothesis/aim/objective of the study clearly described?
2	Are the main outcomes to be measured clearly described in the Introduction or Methods section?
3	Are the characteristics of the patients included in the study clearly described?
4	Are the intervention(s) of interest clearly described?
5	Are the distributions of principal confounders in each group of subjects to be compared clearly described?
6	Are the main findings of the study clearly described?
7	Does the study provide estimates of the random variability in the data for the main outcomes?
8	Have all important adverse events that may be a consequence of the intervention been reported?
9	Have the characteristics of patients lost to follow-up been described?
10	Have actual probability values been reported (e.g., 0.035 rather than <0.05) for the main outcomes except where the probability value is less than 0.001?
11	Were the subjects asked to participate in the study representative of the entire population from which they were recruited?
12	Were those subjects who were prepared to participate representative of the entire population from which they were recruited?
13	Were the staff, places, and facilities where the patients were treated, representative of the treatment the majority of patients receive?
14	Was an attempt made to blind study subjects to the intervention they have received?
15	Was an attempt made to blind those measuring the main outcomes of the intervention?
16	If any of the results of the study were based on "data dredging," was this made clear?
17	In trials and cohort studies, do the analyses adjust for different lengths of follow-up of patients, or in case-control studies, is the period between the intervention and outcome the same for cases and controls?
18	Were the statistical tests used to assess the main outcomes appropriate?
19	Was compliance with the intervention/s reliable?
20	Were the main outcome measures used accurate (valid and reliable)?
21	Were the patients in different intervention groups (trials and cohort studies), or were the cases and controls (case-control studies) recruited from the same population?

22	Were study subjects in different intervention groups (trials and cohort studies), or were the cases and controls (case-control studies) recruited over the same period?
23	Were study subjects randomized to intervention groups?
24	Was the randomized intervention assignment concealed from both patients and health care staff until recruitment was complete and irrevocable?
25	Was there an adequate adjustment for confounding in the analyses from which the main findings were drawn?
26	Were losses of patients to follow-up taken into account?
27	Did the study have sufficient power to detect a clinically important effect where the probability value for a difference is due to chance is less than 5%?

Supplementary material Table 6 Differences in baseline characteristics by gender

Study	Findings
Ladefoged et al. (2020)	The average age of females was significantly higher than that of males (86.4 ± 2.6 years vs. 81.1 ±0.9 years, respectively; p<0.01)
Tansaknen et al. (2008)	The severity of SSA was significantly associated with male gender (p=0.022). Additionally, females with SSA are reported to die at an older age compared to males.
Ueda et al. (2011)	Females were older than males at death, 92.5 (±13.4) vs. 78.5 (±9.7)
Shiozaki et al. (2019)	Females were older than males at death, 88.4 (±6) vs. 85.8 (±10.5) and had higher left ventricular free wall thickness, 1.67 (±0.35) vs. 1.49 (±0.15), respectively.
aus dem Siepen F., et al. (2018)	Significant differences were presented between females and males in the NYHA and EGFR scores 3 (±0.4) vs. 2.5 (±0.7), p=0.006 and 86 (±33.2) vs. 66.7 (±21.4), p=0.014, respectively
Kristen et al. (2013)	The three females with SSA were ten years older than the male study patients.
Gonzalez-Lopez et al. (2017)	Female patients were significantly older than male patients at the time of symptoms' onset and diagnosis (82.3 vs. 76.1; p=0.002) and (83.9 vs. 77.4; p=0.001), respectively. Additionally, women were diagnosed non-invasively more frequently compared to males and presented better LVEF (75% vs. 29%; p<0.001) and 59 vs. 51; p=0.002), respectively.

Supplementary material Figure 2 Funnel plot

