*Braz. J. Nephrol. (J. Bras. Nefrol.)*

https://doi.org/10.1590/2175-8239-JBN-2021-0283

# Supplementary Material to "Adenine phosphoribosyl transferase deficiency lead renal allograft dysfunction among kidney transplant recipients: a systematic review"

**Supplement A** - Study and patient characteristics.

| | Author | Type of study | No of Cases | Sex | Age at diagnosis (Years) | Age at Transplantation (Years) | Patient follow up (months) | Family history of kidney disease | Patient history of kidney disease | Status of kidney disease | Duration of dialysis before transplant (Years) |
|---|---|---|---|---|---|---|---|---|---|---|---|
| **1** | Rajput et al, 2020, India | Case series | 2 | M | 43 | 43 | 2 | NA | kidney stones in the past | ESRD | 1 |
| | | | | M | 31 | 31 | 2 | NA | History of CKD due to bilateral kidney stones | Haemodialysis | 1 |
| **2** | Bagai et al, 2019, India | Case report | 1 | M | 44 | 44 | NA | History of nephrolithiasis | CKD | ESRD | 1 |
| **3** | Li et al, 2019, Australia | Case report | 1 | M | 47 | 52 | 6 | No | Obesity, hypertension, atrial fibrillation | Dialysis | 4 |
| **4** | George et al, 2017, Turkey | Case report | 1 | F | 41 | 41 | NA | Yes, her mother and 1 sister had frequent urinary stones | Significant for recurrent nephrolithiasis and urinary tract infections since childhood | ESRD | NA |
| **5** | Nanumuko et al, 2017, Japan | Case report | 1 | M | 28 | 28 | 4 | No | Several episodes of urinary stone excretion in childhood | Haemodialysis | 5 |
| **6** | Brilland et al, 2015, France | Case report | 1 | F | 25 | 30 | 18 | NA | Recurrent urinary infections, one episode of renal colic in childhood | ESRD | 3 |
| **7** | Kaartinen et al, 2014, Finland | Case report | 1 | M | 63 | ($1^{st}$) 60, ($2^{nd}$) 63 | 11 | Grandfather had urinary stones | Yes | ESRD | 2 |
| **8** | Quaglia et al, 2014, Italy | Case series | 2 | F | 67 | 67 | NA | NA | NA | ESRD | NA |
| | | | | F | NA | 48 | 36 | No | No | ESRD | 1 |
| **9** | Zaiden et al, 2014, France (5), Italy (2) | Case series | 7 | F | 28 | 28 | 132 | NA | CKD due to chronic tubulointerstitial nephropathy | CKD | NA |

| | Author | Type of study | No of Cases | Sex | Age at diagnosis (Years) | Age at Transplantation (Years) | Patient follow up (months) | Family history of kidney disease | Patient history of kidney disease | Status of kidney disease | Duration of dialysis before transplant (Years) |
|---|---|---|---|---|---|---|---|---|---|---|---|
| | | | | M | 41 | 38 | 6 | NA | CKD due to unknown cause | CKD | NA |
| | | | | M | 48 | 46 | 40 | NA | CKD due to unknown cause | CKD | NA |
| | | | | M | 48 | 48 | 24 | NA | CKD due to Oxalate nephropathy | Haemodialysis | NA |
| | | | | F | 49 | 46 | 30 | NA | CKD due to unknown cause | CKD | NA |
| | | | | F | 58 | 58 | 24 | NA | CKD due to chronic tubulointerstitial nephropathy | CKD | NA |
| | | | | M | 64 | 64 | 8 | NA | CKD due to unknown cause | Haemodialysis | NA |
| **10** | Sharma et al, 2012, USA | Case report | 1 | M | 80 | 80 | 1 | NA | Seizure disorder, and multiple episodes of nephrolithiasis prior 43 years of age | ESRD | 3 |
| **11** | Bertram et al, 2010, Germany | Case report | 1 | M | 55 | 39 (1st), 55 (2nd), 56 (3rd) | 9 | Yes | Recurrent nephrolithiasis | ESRD | NA |
| **12** | Micheli et al, 2010, Italy | Case report | 1 | M | 48 | 43 | 2 | No | Recurrent renal staghorn lithiasis on both sides since the age of 17 | Haemodialysis | 1 |
| **13** | Nasr et al, 2010, USA | Case series | 3 | F | 42 | 39 | 12 | Younger sibling is a case of renal failure and deficiency of APRT | History of type 2 diabetes and depression, severe renal failure and anemia | Haemodialysis | 3 |
| | | | | M | 41 | 41 | 4 | Elder sibling is a case of renal failure and deficiency of APRT | Hypertension and diet-controlled type 2 diabetes mellitus, acute and chronic tubulointerstitial nephritis and arteriolar hyalinosis. | Haemodialysis | NA |
| | | | | M | 55 | 54 | 18 | Negative | Type 2 diabetes for 8 years, hypertension and obesity. At the age of 18, he reported kidney stones | Dialysis | 1 |
| **14** | Stratta et al, 2010, Italy | Case report | 1 | F | 67 | 67 | 12 | Family history was negative except for hypertension | Repeated renal colic with recurrent episodes of spontaneous elimination of small kidney stones since childhood | ESRD | 2 |

| | Author | Type of study | No of Cases | Sex | Age at diagnosis (Years) | Age at Transplantation (Years) | Patient follow up (months) | Family history of kidney disease | Patient history of kidney disease | Status of kidney disease | Duration of dialysis before transplant (Years) |
|---|---|---|---|---|---|---|---|---|---|---|---|
| **15** | Cassidy et al, 2004, London | Case report | 1 | M | 23 | 23 | 8 | Mother and father are carriers of APRT deficiency | Chronic renal failure of unknown cause | CKD | NA |
| **16** | Eller et al, 2004, NA | Case report | 1 | M | 11 | NA | 7 | NA | History of nephrolithiasis, CKD | Haemodialysis | NA |
| **17** | Benedetto et al, 2001, USA | Case report | 1 | M | 44 | 43 | 19 | NA | Two episodes of urolithiasis | ESRD | NA |
| **18** | Brown et al, 1998, Canada | Case report | 1 | F | 47 | 47 | 4 | NA | Episode of pyelonephritis | ESRD | 2 |
| **19** | De jong et al, 1996, Netherland | Case report | 1 | M | 56 | 56 | 6 | NA | History of recurrent radiolucent and radiopaque urolithiasis of supposed uric acid composition | Haemodialysis | NA |
| **20** | Gagne et al, 1994, Canada | Case report | 1 | M | 51 | 42 | 108 | NA | History of urolithiasis at the age of 8 years. | ESRD | 5 |

M = Male, F = Female. ESRD = End Stage Renal Disease. NA = Not available. CKD = chronic kidney disease