Online supplementary file 5: Sensitivity analyses of the all-cause mortality by comparison between MCS and non-MCS

		Risk Ratio (95%CI)
Short-term mortality		
	Inconsistent length of follow-up	
	In hospital	1.37 (0.50-3.78)
	30 days	0.70 (0.39-1.25)
	Different types of MCS use	
	IABP vs non-IABP	0.92 (0.57-1.47)
	ECLS vs non-ECLS	0.33 (0.01-1.47)
	Excluding high risk bias studies	
	Zhou et al[29], Arias et al[33] and Korbinian et al[35]	0.85 (0.51-1.45)
	Excluding each of the studies	
	SEMPER FI[25]	0.92 (0.57-1.48)
	Zhou et al[29]	0.80 (0.53-1.21)
	IABP-SHOCK II[8]	0.89 (0.45-1.74)
	CRISP AMI[28]	0.95 (0.59-1.55)
	Gu et al[30]	1.02 (0.66-1.58)
	Prondzinsky et al[31]	0.80 (0.48-1.34)
	Arias et al[33]	0.98 (0.58-1.66)
	Vijayalakshmi et al[34]	0.86 (0.55-1.34)
	Korbinian et al[35]	0.92 (0.57-1.47)
Long-term mortality		
	Inconsistent length of follow-up	
	6 months	0.72 (0.43-1.20)
	12 months	0.90 (0.35-2.31)
	Different types of MCS use	
	IABP vs non-IABP	0.87 (0.60-1.25)
	ECLS vs non-ECLS	0.50 (0.18-1.41)
	Excluding high risk bias studies	
	Zhou et al[29] and Korbinian et al[35]	0.72 (0.43-1.20)
	Excluding each of the studies	
	SEMPER FI[25]	0.82 (0.56-1.20)
	Zhou et al[29]	0.70 (0.45-1.09)
	IABP-SHOCK II[9]	0.68 (0.40-1.17)
	CRISP AMI[28]	0.89 (0.65-1.23)
	Gu et al[30]	0.92 (0.66-1.28)
	Korbinian et al[35]	0.87 (0.61-1.25)

CI indicates confidence interval.