Supplementary Table 2: Risk of bias assessment for RCTs [a] and quasi-experiments [b]

RCTs	**Risk of bias (High, low, unclear)**					
Source	Randomisation sequence Allocation	Concealment	Performance bias	Ascertainment bias [c]	Incomplete outcome data	selective reporting
Bassey et al. 1987	Unclear	Unclear	High	Low	High	Low
Sharpe et al. 1992	High	Unclear	High	High	High	Low
Granacher et al. 2011	High	Unclear	High	Low	Unclear	Low
Hughes et al. 2011	Low	Unclear	High	High	High	Low
Strijk et al. 2012	Low	Low	High	Low	High	High
Palumbo et al. 2012	Unclear	Unclear	High	low	High	High
Cook et al. 2015	Low	Low	High	High	High	Low
Low 2015	Unclear	Unclear	High	high	Unclear	High
Kocur et al. 2017	Unclear	Low	High	Low	High	Low
Chopp-Hurley et al. 2017	Low	Low	High	Low	Low	Low
Bergman et al. 2018	Low	unclear	High	Low	Low	Low
Qi et al. 2019	unclear	Low	High	low	High	Low
Non-RCT: Confounding						
Chen et al., 2016	High	Low	High	High	High	Low
Arao, 2007	High	Low	High	Low	High	Low

[a] Higgins JPT, Altman DG, Gotzsche PC, et al. Cochrane Bias Methods Group Cochrane Statistical Methods Group. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ* 2011;343:d5928. doi:10.1136/bmj.d5928

[b] Sterne JA, Hernán MA, Reeves BC, et al. ROBINS-I: a tool for assessing risk of bias in non-randomised studies of interventions. BMJ. 2016 Oct 12;355:i4919. doi: 10.1136/bmj.i4919. PMID: 27733354; PMCID: PMC5062054.

[c] Studies that had included objective physical activity outcomes and subjective outcomes were classified as low