S20 File. Subgroup analysis



BD – bipolar disorder, SCZ – schizophrenia, SMI – mixed severe mental illness, CHD – coronary heart disease, CVA – cerebrovascular accident, CVD – all circulatory disease, HF – heart failure, HR – hazard ratio, SMR – standardised mortality ratio, FU – follow-up

Fig A: Forest plots showing studies reporting CVD mortality risk for severe mental illness vs controls for a) age groups ≤60; b) older age groups or all ages





SCZ – schizophrenia, HR – hazard ratio, SMR – standardised mortality ratio, OR – odds ratio, CHD – coronary heart disease, CVA – cerebrovascular accident, CVD – all circulatory disease, HF – heart failure

Fig B: Forest plots showing relative risk of CVD mortality in those with vs without schizophrenia, studies included in meta-analysis for a) males; b) females



BD – bipolar disorder, HR – hazard ratio, SMR – standardised mortality ratio, OR – odds ratio, CHD – coronary heart disease, CVA – cerebrovascular accident, CVD – all circulatory disease

Fig C: Forest plots showing relative risk of CVD mortality in those with vs without bipolar disorder, studies included in meta-analysis for a) males; b) females





SMI – severe mental illness, SMR – standardised mortality ratio, CVD – all circulatory disease

Fig D: Forest plots showing relative risk of CVD mortality in those with vs without mixed SMI, studies included in meta-analysis for a) males; b) females



BD – bipolar disorder, SCZ – schizophrenia, CHD – coronary heart disease, CVA – cerebrovascular accident, CVD – major cardiovascular events, HR – hazard ratio, OR – odds ratio, SIR – standardised incidence ratio,

Fig E: Forest plots of CVD incidence for a) studies with low risk of bias, b) all studies included in meta-analysis, schizophrenia and bipolar disorder