# A systematic review of behavioural and exercise interventions for the prevention and management of chemotherapy-induced peripheral neuropathy symptoms

Journal of Cancer Survivorship
Tanay MAL, Armes J, Moss-Morris R, Rafferty AM and Robert G
Corresponding author: Tanay MAL mary.tanay@kcl.ac.uk

Online material

Table S-1. Exemplar of the search protocol for Medline.

1. chemotherapy.mp.
2. peripheral neuropathy.mp.
3. neuropathy.mp.
4. 2 or 3
5. 1 and 4
6. intervention.mp.
7. program$.mp. or exp *Programs/
8. system.mp.
9. exercise.mp
10. exp *Physical Activity as Topic
11. follow-up.mp.
12. model.mp.
13. 6 or 7 or 8 or 9 or 10 or 11 or 12
14. psycholog*.mp.
15. exp Behavior Therapy/ or exp Behavior/ or behavio#r.mp.
16. education*.mp.
17. exp *Patient Education as Topic/ or psychoeducation*.mp.
18. self-management.mp. or exp Self Care/ or exp Self-Management/
19. motivational interview*.mp. or Motivational Interviewing /
20. 14 or 15 or 16 or 17 or 18 or 19
21. 5 and 13 and 20
22. limit 21 to (english language and yr="2000 -Current" and english and humans and cancer and "humans only (remove records about animals)")

Table S-2. Results of quality assessment of studies using the EPHPP a Quality Assessment Tool for Quantitative Studies (Thomas et al. 2004)

| STUDIES INCLUDED | CRITERIA | | | | | | |
|---|---|---|---|---|---|---|---|
| | Selection bias | Study design | Confounders | Blinding Process | Data collection methods | Sample attrition | Overall |
| **Behavioural intervention studies** | | | | | | | |
| Given et al. (2008) | Moderate | Strong | Moderate | Weak | Moderate | Moderate | Moderate |
| Tofthagen et al. (2016) | Moderate | Moderate | Weak | Moderate | Strong | Moderate | Moderate |
| Knoerl et al. (2018a) | Moderate | Moderate | Moderate | Moderate | Strong | Strong | Strong |
| Knoerl et al. (2018b) | Strong | Strong | Moderate | Strong | Strong | Moderate | Strong |
| Kolb et al. (2018) | Moderate | Strong | Weak | Moderate | Strong | Moderate | Moderate |
| Knoerl et al. (2019) | Moderate | Strong | Moderate | Moderate | Strong | Strong | Strong |
| **Exercise intervention studies** | | | | | | | |
| Wonders et al. (2013) | Moderate | Moderate | Moderate | Moderate | Weak | Weak | Weak |
| Streckman et al. (2014) | Strong | Moderate | Moderate | Weak | Moderate | Weak | Weak |
| Tofthagen et al. (2014) | Weak | Moderate | Weak | Weak | Moderate | Weak | Weak |
| Fernandes & Kumar (2016) | Weak | Moderate | Weak | Weak | Weak | Weak | Weak |
| Schwenk et al. (2016) | Moderate | Strong | Moderate | Moderate | Strong | Moderate | Moderate |
| Kleckner et al. (2018) | Moderate | Strong | Strong | Moderate | Strong | Strong | Strong |
| Vollmers et al. (2018) | Moderate | Weak | Moderate | Moderate | Strong | Moderate | Moderate |
| Zimmer et al (2018) | Moderate | Moderate | Moderate | Moderate | Strong | Strong | Moderate |
| McCrary et al. (2019) | Moderate | Moderate | Moderate | Moderate | Strong | Strong | Moderate |
| Kneis et al. (2019) | Strong | Moderate | Moderate | Moderate | Strong | Strong | Moderate |
| Bland et al. (2019) | Strong | Moderate | Strong | Weak | Moderate | Strong | Moderate |
| Hammond et al. (2020) | Moderate | Weak | Weak | Moderate | Weak | Weak | Weak |
| Bahar-Ozdemir et al. (2020) | Moderate | Moderate | Weak | Moderate | Moderate | Moderate | Moderate |