Supplementary Table III.

1994 Macdonnel	Macdonell RAL, Triggs WJ, Leikauskas J, Bourque M, Robb K, Day BJ, et al. Functional electrical stimulation to the affected lower limb and recovery after cerebral infarction. J Stroke Cerebrovasc Dis 1994;4:155–60. https://doi.org/10.1016/S1052-3057(10)80178-8.
1999 Kwakkel	Kwakkel G, Wagenaar RC, Twisk JW, Lankhorst GJ, Koetsier JC. Intensity of leg and arm training after primary middle-cerebral-artery stroke: a randomised trial. The Lancet 1999;354:191–6. https://doi.org/10.1016/S0140-6736(98)09477-X.
2001 Cheng	Cheng PT, Wu SH, Liaw MY, Wong AM, Tang FT. Symmetrical body-weight distribution training in stroke patients and its effect on fall prevention. Arch Phys Med Rehabil 2001;82:1650–4. https://doi.org/10.1053/apmr.2001.26256.
2003 Duncan	Duncan P, Studenski S, Richards L, Gollub S, Lai SM, Reker D, et al. Randomized Clinical Trial of Therapeutic Exercise in Subacute Stroke. Stroke 2003;34:2173–80. https://doi.org/10.1161/01.STR.0000083699.95351.F2.
2003 Moreland	Moreland JD, Goldsmith CH, Huijbregts MP, Anderson RE, Prentice DM, Brunton KB, et al. Progressive resistance strengthening exercises after stroke: a single-blind randomized controlled trial. Arch Phys Med Rehabil 2003;84:1433–40. https://doi.org/10.1016/S0003-9993(03)00360-5.
2004 Newsman	Newsam CJ, Baker LL. Effect of an electric stimulation facilitation program on quadriceps motor unit recruitment after stroke. Arch Phys Med Rehabil 2004;85:2040–5. https://doi.org/10.1016/j.apmr.2004.02.029.
2004 Richards	Richards CL, Malouin F, Bravo G, Dumas F, Wood-Dauphinee S. The Role of Technology in Task-Oriented Training in Persons with Subacute Stroke: A Randomized Controlled Trial. Neurorehabil Neural Repair 2004;18:199–211. https://doi.org/10.1177/1545968304269397.
2005 Park	Park J, White AR, James MA, Hemsley AG, Johnson P, Chambers J, et al. Acupuncture for Subacute Stroke Rehabilitation: A Sham-Controlled, Subject- and Assessor-Blind, Randomized Trial. Arch Intern Med 2005;165:2026. https://doi.org/10.1001/archinte.165.17.2026.
2005 Van Vliet	van Vliet PM. Comparison of Bobath based and movement science based treatment for stroke: a randomised controlled trial. J Neurol Neurosurg Psychiatry 2005;76:503–8. https://doi.org/10.1136/jnnp.2004.040436.
2005 Yan	Yan T, Hui-Chan CWY, Li LSW. Functional Electrical Stimulation Improves Motor Recovery of the Lower Extremity and Walking Ability of Subjects With First Acute Stroke: A Randomized Placebo-Controlled Trial. Stroke 2005;36:80–5. https://doi.org/10.1161/01.STR.0000149623.24906.63.
2005 Yang	Yang Y-R, Yen J-G, Wang R-Y, Yen L-L, Lieu F-K. Gait outcomes after additional backward walking training in patients with stroke: a randomized controlled trial. Clin Rehabil 2005;19:264–73. https://doi.org/10.1191/0269215505cr860oa.
2006 Tong	Tong RK, Ng MF, Li LS. Effectiveness of gait training using an electromechanical gait trainer, with and without functional electric stimulation, in subacute stroke: a randomized controlled trial. Arch Phys Med Rehabil 2006;87:1298–304. https://doi.org/10.1016/j.apmr.2006.06.016.
2006 Yavuzer	Yavuzer G, Geler-Külcü D, Sonel-Tur B, Kutlay S, Ergin S, Stam HJ. Neuromuscular electric stimulation effect on lower-extremity motor recovery and gait kinematics of patients with stroke: a randomized controlled trial. Arch Phys Med Rehabil 2006;87:536–40. https://doi.org/10.1016/j.apmr.2005.12.041.
2007 Dean	Dean CM, Channon EF, Hall JM. Sitting training early after stroke improves sitting ability and quality and carries over to standing up but not to walking: a randomised trial. Aust J Physiother 2007;53:97–102.
2007 Sutbeyaz	Sütbeyaz S, Yavuzer G, Sezer N, Koseoglu BF. Mirror therapy enhances lower-extremity motor recovery and motor functioning after stroke: a randomized controlled trial. Arch Phys Med Rehabil 2007;88:555–9. https://doi.org/10.1016/j.apmr.2007.02.034.
2007 Tseng	Tseng C-N, Chen CC-H, Wu S-C, Lin L-C. Effects of a range-of-motion exercise programme. J Adv Nurs 2007;57:181–91. https://doi.org/10.1111/j.1365-2648.2006.04078.x.
2007 Wang	Wang T, Wang X, Wang H, He X, Su J, Zhu Y, et al. Effects of ULEM apparatus on motor function of patients with stroke. Brain Inj 2007;21:1203–8. https://doi.org/10.1080/02699050701633064.

2007 Yavuzer	Yavuzer G, Oken O, Atay MB, Stam HJ. Effect of sensory-amplitude electric stimulation on motor recovery and gait kinematics after stroke: a randomized controlled study. Arch Phys Med Rehabil 2007;88:710–4. https://doi.org/10.1016/j.apmr.2007.02.030.
2007 Yen	Yen C-L, Wang R-Y, Liao K-K, Huang C-C, Yang Y-R. Gait Training-Induced Change in Corticomotor Excitability in Patients With Chronic Stroke. Neurorehabil Neural Repair 2008;22:22–30. https://doi.org/10.1177/1545968307301875.
2008 Eser	Eser F, Yavuzer G, Karakus D, Karaoglan B. The effect of balance training on motor recovery and ambulation after stroke: a randomized controlled trial. Eur J Phys Rehabil Med 2008;44:19–25.
2008 Ferrante	Ferrante S, Pedrocchi A, Ferrigno G, Molteni F. Cycling induced by functional electrical stimulation improves the muscular strength and the motor control of individuals with post-acute stroke. Eur J Phys Rehabil Med 2008;44:159–67.
2008 Gok	Gok H, Geler-Kulcu D, Alptekin N, Dincer G. Efficacy of treatment with a kinaesthetic ability training device on balance and mobility after stroke: a randomized controlled study. Clin Rehabil 2008;22:922–30. https://doi.org/10.1177/0269215508090673.
2009 Mesci	Mesci N, Ozdemir F, Kabayel DD, Tokuc B. The effects of neuromuscular electrical stimulation on clinical improvement in hemiplegic lower extremity rehabilitation in chronic stroke: A single-blind, randomised, controlled trial. Disabil Rehabil 2009;31:2047–54. https://doi.org/10.3109/09638280902893626.
2009 Yan	Yan T, Hui-Chan CWY. Transcutaneous electrical stimulation on acupuncture points improves muscle function in subjects after acute stroke: a randomized controlled trial. J Rehabil Med 2009;41:312–6. https://doi.org/10.2340/16501977-0325.
2010 Cooke	Cooke EV, Tallis RC, Clark A, Pomeroy VM. Efficacy of functional strength training on restoration of lower-limb motor function early after stroke: phase I randomized controlled trial. Neurorehabil Neural Repair 2010;24:88–96. https://doi.org/10.1177/1545968309343216.
2010 Goljar	Goljar N, Burger H, Rudolf M, Stanonik I. Improving balance in subacute stroke patients: a randomized controlled study. Int J Rehabil Res 2010;33:205–10. https://doi.org/10.1097/MRR.0b013e328333de61.
2010 Ivey	Ivey FM, Hafer-Macko CE, Ryan AS, Macko RF. Impaired Leg Vasodilatory Function After Stroke: Adaptations With Treadmill Exercise Training. Stroke 2010;41:2913–7. https://doi.org/10.1161/STROKEAHA.110.599977.
2010 Jonsdottir	Jonsdottir J, Cattaneo D, Recalcati M, Regola A, Rabuffetti M, Ferrarin M, et al. Task-Oriented Biofeedback to Improve Gait in Individuals With Chronic Stroke: Motor Learning Approach. Neurorehabil Neural Repair 2010;24:478–85. https://doi.org/10.1177/1545968309355986.
2010 Paoloni	Paoloni M, Mangone M, Scettri P, Procaccianti R, Cometa A, Santilli V. Segmental muscle vibration improves walking in chronic stroke patients with foot drop: a randomized controlled trial. Neurorehabil Neural Repair 2010;24:254–62. https://doi.org/10.1177/1545968309349940.
2010 Tihanyi	Tihanyi J, Di Giminiani R, Tihanyi T, Gyulai G, Trzaskoma L, Horváth M. Low resonance frequency vibration affects strength of paretic and non-paretic leg differently in patients with stroke. Acta Physiol Hung 2010;97:172–82. https://doi.org/10.1556/APhysiol.97.2010.2.3.
2010 Tung	Tung F-L, Yang Y-R, Lee C-C, Wang R-Y. Balance outcomes after additional sit-to-stand training in subjects with stroke: a randomized controlled trial. Clin Rehabil 2010;24:533–42. https://doi.org/10.1177/0269215509360751.
2011 Galvin	Galvin R, Cusack T, O'Grady E, Murphy TB, Stokes E. Family-Mediated Exercise Intervention (FAME): Evaluation of a Novel Form of Exercise Delivery After Stroke. Stroke 2011;42:681–6. https://doi.org/10.1161/STROKEAHA.110.594689.
2011 Hesse	Hesse S, Welz A, Werner C, Quentin B, Wissel J. Comparison of an intermittent high-intensity vs continuous low-intensity physiotherapy service over 12 months in community-dwelling people with stroke: a randomized trial. Clin Rehabil 2011;25:146–56. https://doi.org/10.1177/0269215510382148.

SUPPLEMENTARY DIGITAL MATERIAL 3

Supplementary Table III.—References of the included studies.

2011 Kim	Kim CS, Gong W, Kim SG. The Effects of Lower Extremitiy Muscle Strengthening Exercise and Treadmill Walking Exercise on the Gait and Balance of Stroke Patients. J Phys Ther Sci 2011;23:405–8. https://doi.org/10.1589/jpts.23.405.
2011 Sungkarat	Sungkarat S, Fisher BE, Kovindha A. Efficacy of an insole shoe wedge and augmented pressure sensor for gait training in individuals with stroke: a randomized controlled trial. Clin Rehabil 2011;25:360–9. https://doi.org/10.1177/0269215510386125.
2011 Varoqui	Varoqui D, Froger J, Pélissier J-Y, Bardy BG. Effect of coordination biofeedback on (re)learning preferred postural patterns in post-stroke patients. Motor Control 2011;15:187–205. https://doi.org/10.1123/mcj.15.2.187.
2012 Batchelor	Batchelor FA, Hill KD, Mackintosh SF, Said CM, Whitehead CH. Effects of a Multifactorial Falls Prevention Program for People With Stroke Returning Home After Rehabilitation: A Randomized Controlled Trial. Arch Phys Med Rehabil 2012;93:1648–55. https://doi.org/10.1016/j.apmr.2012.03.031.
2012 Chang	Chang WH, Kim MS, Huh JP, Lee PKW, Kim Y-H. Effects of robot-assisted gait training on cardiopulmonary fitness in subacute stroke patients: a randomized controlled study. Neurorehabil Neural Repair 2012;26:318–24. https://doi.org/10.1177/1545968311408916.
2012 Chumbler	Chumbler NR, Quigley P, Li X, Morey M, Rose D, Sanford J, et al. Effects of Telerehabilitation on Physical Function and Disability for Stroke Patients: A Randomized, Controlled Trial. Stroke 2012;43:2168–74. https://doi.org/10.1161/STROKEAHA.111.646943.
2012 Globas	Globas C, Becker C, Cerny J, Lam JM, Lindemann U, Forrester LW, et al. Chronic Stroke Survivors Benefit From High-Intensity Aerobic Treadmill Exercise: A Randomized Control Trial. Neurorehabil Neural Repair 2012;26:85–95. https://doi.org/10.1177/1545968311418675.
2012 Mohapatra	Mohapatra S, Eviota AC, Ringquist KL, Muthukrishnan SR, Aruin AS. Compelled Body Weight Shift Technique to Facilitate Rehabilitation of Individuals with Acute Stroke. ISRN Rehabil 2012;2012:1–7. https://doi.org/10.5402/2012/328018.
2013 Hsu	Hsu H-W, Lee C-L, Hsu M-J, Wu H-C, Lin R, Hsieh C-L, et al. Effects of noxious versus innocuous thermal stimulation on lower extremity motor recovery 3 months after stroke. Arch Phys Med Rehabil 2013;94:633–41. https://doi.org/10.1016/j.apmr.2012.11.021.
2013 MacKay-Lyons	Mackay-Lyons M, McDonald A, Matheson J, Eskes G, Klus M-A. Dual effects of body-weight supported treadmill training on cardiovascular fitness and walking ability early after stroke: a randomized controlled trial. Neurorehabil Neural Repair 2013;27:644–53. https://doi.org/10.1177/1545968313484809.
2013 Rao	Rao N, Zielke D, Keller S, Burns M, Sharma A, Krieger R, et al. Pregait balance rehabilitation in acute stroke patients. Int J Rehabil Res 2013;36:112–7. https://doi.org/10.1097/MRR.0b013e328359a2fa.
2013 Sekhar	Sekhar PKC, Madhavi K, Srikumari V. Efficacy of Isokinetic Strength Training and Balance Exercises on Lower Limb Muscles in Subjects with Stroke. Indian J Physiother Occup Ther - Int J 2013;7:112. https://doi.org/10.5958/j.0973-5674.7.4.132.
2013 Sheffler	Sheffler LR, Taylor PN, Gunzler DD, Buurke JH, Ijzerman MJ, Chae J. Randomized controlled trial of surface peroneal nerve stimulation for motor relearning in lower limb hemiparesis. Arch Phys Med Rehabil 2013;94:1007–14. https://doi.org/10.1016/j.apmr.2013.01.024.
2013 Zhang	Zhang Y, Jin H, Ma D, Fu Y, Xie Y, Li Z, et al. Efficacy of Integrated Rehabilitation Techniques of Traditional Chinese Medicine for Ischemic Stroke: A Randomized Controlled Trial. Am J Chin Med 2013;41:971–81. https://doi.org/10.1142/S0192415X13500651.
2014 Cha	Cha H-K, Ji S-G, Kim M-K, Chang J-S. Effect of Transcranial Direct Current Stimulation of Function in Patients with Stroke. J Phys Ther Sci 2014;26:363–5. https://doi.org/10.1589/jpts.26.363.
2014 Chen	Chen I-H, Yang Y-R, Chan R-C, Wang R-Y. Turning-Based Treadmill Training Improves Turning Performance and Gait Symmetry After Stroke. Neurorehabil Neural Repair 2014;28:45–55. https://doi.org/10.1177/1545968313497102.
2014 De Nunzio	De Nunzio AM, Zucchella C, Spicciato F, Tortola P, Vecchione C, Pierelli F, et al. Biofeedback rehabilitation of posture and weightbearing distribution in stroke: a center of foot pressure analysis. Funct Neurol 2014;29:127–34.

2014 Dragin	Dragin AS, Konstantinović LM, Veg A, Schwirtlich LB. Gait training of poststroke patients assisted by the Walkaround (body postural support). Int J Rehabil Res Int Z Rehabil Rev Int Rech Readaptation 2014;37:22–8. https://doi.org/10.1097/MRR.0b013e328363ba30.
2014 Hung	Hung J-W, Chou C-X, Hsieh Y-W, Wu W-C, Yu M-Y, Chen P-C, et al. Randomized Comparison Trial of Balance Training by Using Exergaming and Conventional Weight-Shift Therapy in Patients With Chronic Stroke. Arch Phys Med Rehabil 2014;95:1629–37. https://doi.org/10.1016/j.apmr.2014.04.029.
2014 Ji	Ji SG, Kim MK. The effects of mirror therapy on the gait of subacute stroke patients: a randomized controlled trial. Clin Rehabil 2015;29:348–54. https://doi.org/10.1177/0269215514542356.
2014 McEwen	McEwen D, Taillon-Hobson A, Bilodeau M, Sveistrup H, Finestone H. Virtual reality exercise improves mobility after stroke: an inpatient randomized controlled trial. Stroke 2014;45:1853–5. https://doi.org/10.1161/STROKEAHA.114.005362.
2014 Suh	Suh HR, Han HC, Cho H. Immediate therapeutic effect of interferential current therapy on spasticity, balance, and gait function in chronic stroke patients: a randomized control trial. Clin Rehabil 2014;28:885–91. https://doi.org/10.1177/0269215514523798.
2014 van Nunen	van Nunen MPM, Gerrits KHL, Konijnenbelt M, Janssen TWJ, de Haan A. Recovery of walking ability using a robotic device in subacute stroke patients: a randomized controlled study. Disabil Rehabil Assist Technol 2015;10:141–8. https://doi.org/10.3109/17483107.2013.873489.
2014 Yang	Yang H-C, Lee C-L, Lin R, Hsu M-J, Chen C-H, Lin J-H, et al. Effect of biofeedback cycling training on functional recovery and walking ability of lower extremity in patients with stroke. Kaohsiung J Med Sci 2014;30:35–42. https://doi.org/10.1016/j.kjms.2013.07.006.
2014 You	You G, Liang H, Yan T. Functional electrical stimulation early after stroke improves lower limb motor function and ability in activities of daily living. NeuroRehabilitation 2014;35:381–9. https://doi.org/10.3233/NRE-141129.
2015 Chang	Chang MC, Kim DY, Park DH. Enhancement of Cortical Excitability and Lower Limb Motor Function in Patients With Stroke by Transcranial Direct Current Stimulation. Brain Stimulat 2015;8:561–6. https://doi.org/10.1016/j.brs.2015.01.411.
2015 da Silva Ribeiro	da Silva Ribeiro NM, Ferraz DD, Pedreira É, Pinheiro Í, da Silva Pinto AC, Neto MG, et al. Virtual rehabilitation via Nintendo Wii® and conventional physical therapy effectively treat post-stroke hemiparetic patients. Top Stroke Rehabil 2015;22:299–305. https://doi.org/10.1179/1074935714Z.0000000017.
2015 Ding	Ding X-D, Zhang G-B, Chen H-X, Wang W, Song J-H, Fu D-G. Color Doppler ultrasound-guided botulinum toxin type a injection combined with an ankle foot brace for treating lower limb spasticity after a stroke. Eur Rev Med Pharmacol Sci 2015;19:406–11.
2015 Fujino	Fujino Y, Amimoto K, Fukata K, Ishihara S, Makita S, Takahashi H. Does training sitting balance on a platform tilted 10° to the weak side improve trunk control in the acute phase after stroke? A randomized, controlled trial. Top Stroke Rehabil 2016;23:43–9. https://doi.org/10.1179/1945511915Y.0000000010.
2015 Gürcan	Gurcan A, Selcuk B, Onder B, Akyuz M, Akbal Yavuz A. Evaluation of Clinical and Electrophysiological Effects of Electrical Stimulation on Spasticity of Plantar Flexor Muscles in Patients with Stroke. Türkiye Fiz Tip Ve Rehabil Derg 2015;61:307–13. https://doi.org/10.5152/tftrd.2015.38981.
2015 Hwang	Hwang D-Y, Lee H-J, Lee G-C, Lee S-M. Treadmill training with tilt sensor functional electrical stimulation for improving balance, gait, and muscle architecture of tibialis anterior of survivors with chronic stroke: A randomized controlled trial. Technol Health Care 2015;23:443–52. https://doi.org/10.3233/THC-150903.
2015 Kawakami	Kawakami K, Miyasaka H, Nonoyama S, Hayashi K, Tonogai Y, Tanino G, et al. Randomized controlled comparative study on effect of training to improve lower limb motor paralysis in convalescent patients with post-stroke hemiplegia. J Phys Ther Sci 2015;27:2947–50. https://doi.org/10.1589/jpts.27.2947.
2015 Kim (a)	Kim C-Y, Lee J-S, Kim H-D, Kim J, Lee I-H. Lower extremity muscle activation and function in progressive task-oriented training on the supplementary tilt table during stepping-like movements in

	patients with acute stroke hemiparesis. J Electromyogr Kinesiol 2015;25:522–30. https://doi.org/10.1016/j.jelekin.2015.03.004.
2015 Kim (b)	Kim C-Y, Lee J-S, Kim H-D, Kim J-S. The effect of progressive task-oriented training on a supplementary tilt table on lower extremity muscle strength and gait recovery in patients with hemiplegic stroke. Gait Posture 2015;41:425–30. https://doi.org/10.1016/j.gaitpost.2014.11.004.
2015 Lee	Lee YH, Park SH, Yoon ES, Lee C-D, Wee SO, Fernhall B, et al. Effects of combined aerobic and resistance exercise on central arterial stiffness and gait velocity in patients with chronic poststroke hemiparesis. Am J Phys Med Rehabil 2015;94:687–95. https://doi.org/10.1097/PHM.0000000000000233.
2015 Mao	Mao Y-R, Lo WL, Lin Q, Li L, Xiao X, Raghavan P, et al. The Effect of Body Weight Support Treadmill Training on Gait Recovery, Proximal Lower Limb Motor Pattern, and Balance in Patients with Subacute Stroke. BioMed Res Int 2015;2015:175719. https://doi.org/10.1155/2015/175719.
2015 Ochi	Ochi M, Wada F, Saeki S, Hachisuka K. Gait training in subacute non-ambulatory stroke patients using a full weight-bearing gait-assistance robot: A prospective, randomized, open, blinded-endpoint trial. J Neurol Sci 2015;353:130–6. https://doi.org/10.1016/j.jns.2015.04.033.
2015 Oostra	Oostra K, Oomen A, Vanderstraeten G, Vingerhoets G. Influence of motor imagery training on gait rehabilitation in sub-acute stroke: A randomized controlled trial. J Rehabil Med 2015;47:204–9. https://doi.org/10.2340/16501977-1908.
2015 Park	Park K-H, Kim D-Y, Kim T-H. The effect of step climbing exercise on balance and step length in chronic stroke patients. J Phys Ther Sci 2015;27:3515–8. https://doi.org/10.1589/jpts.27.3515.
2015 Sheffler	Sheffler LR, Taylor PN, Bailey SN, Gunzler DD, Buurke JH, IJzerman MJ, et al. Surface Peroneal Nerve Stimulation in Lower Limb Hemiparesis: Effect on Quantitative Gait Parameters. Am J Phys Med Rehabil 2015;94:341–57. https://doi.org/10.1097/PHM.0000000000000269.
2015 Shen	Shen D; Huang H; Yuan H; Ye S; Li M; Gu J; Wang Z. Therapeutic efficacy of intensified walk training under the electrocardiogram telemetry in stroke induced lower limb dysfunction patients with heart failure. Int J Clin Exp Med 2015 Sep 308916599-16605 2015.
2015 Wang	Wang Y, Meng F, Zhang Y, Xu M, Yue S. Full-movement neuromuscular electrical stimulation improves plantar flexor spasticity and ankle active dorsiflexion in stroke patients: a randomized controlled study. Clin Rehabil 2016;30:577–86. https://doi.org/10.1177/0269215515597048.
2015 Zou	Zou J, Wang Z, Qu Q, Wang L. Resistance training improves hyperglycemia and dyslipidemia, highly prevalent among nonelderly, nondiabetic, chronically disabled stroke patients. Arch Phys Med Rehabil 2015;96:1291–6. https://doi.org/10.1016/j.apmr.2015.03.008.
2016 Chen	Chen L, Fang J, Ma R, Gu X, Chen L, Li J, et al. Additional effects of acupuncture on early comprehensive rehabilitation in patients with mild to moderate acute ischemic stroke: a multicenter randomized controlled trial. BMC Complement Altern Med 2016;16:226. https://doi.org/10.1186/s12906-016-1193-y.
2016 Du	Du J, Tian L, Liu W, Hu J, Xu G, Ma M, et al. Effects of repetitive transcranial magnetic stimulation on motor recovery and motor cortex excitability in patients with stroke: a randomized controlled trial. Eur J Neurol 2016;23:1666–72. https://doi.org/10.1111/ene.13105.
2016 Han	Han EY, Im SH, Kim BR, Seo MJ, Kim MO. Robot-assisted gait training improves brachial-ankle pulse wave velocity and peak aerobic capacity in subacute stroke patients with totally dependent ambulation: Randomized controlled trial. Medicine (Baltimore) 2016;95:e5078. https://doi.org/10.1097/MD.0000000000005078.
2016 In	In T, Lee K, Song C. Virtual Reality Reflection Therapy Improves Balance and Gait in Patients with Chronic Stroke: Randomized Controlled Trials. Med Sci Monit 2016;22:4046–53. https://doi.org/10.12659/MSM.898157.
2016 Salhab	Salhab G, Sarraj AR, Saleh S. Mirror therapy combined with functional electrical stimulation for rehabilitation of stroke survivors' ankle dorsiflexion. Annu Int Conf IEEE Eng Med Biol Soc IEEE Eng Med Biol Soc Annu Int Conf 2016;2016:4699–702. https://doi.org/10.1109/EMBC.2016.7591776.

2016 Sandberg	Sandberg K, Kleist M, Falk L, Enthoven P. Effects of Twice-Weekly Intense Aerobic Exercise in Early Subacute Stroke: A Randomized Controlled Trial. Arch Phys Med Rehabil 2016;97:1244–53. https://doi.org/10.1016/j.apmr.2016.01.030.
2016 Zhang	Zhang Y, Wang Y-Z, Huang L-P, Bai B, Zhou S, Yin M-M, et al. Aquatic Therapy Improves Outcomes for Subacute Stroke Patients by Enhancing Muscular Strength of Paretic Lower Limbs Without Increasing Spasticity: A Randomized Controlled Trial. Am J Phys Med Rehabil 2016;95:840–9. https://doi.org/10.1097/PHM.0000000000000512.
2017 Kim J	Kim J, Yim J. Effects of an Exercise Protocol for Improving Handgrip Strength and Walking Speed on Cognitive Function in Patients with Chronic Stroke. Med Sci Monit 2017;23:5402–9. https://doi.org/10.12659/MSM.904723.
2017 An	An C-M, Jo S-O. Effects of Talocrural Mobilization with Movement on Ankle Strength, Mobility, and Weight-Bearing Ability in Hemiplegic Patients with Chronic Stroke: A Randomized Controlled Trial. J Stroke Cerebrovasc Dis Off J Natl Stroke Assoc 2017;26:169–76. https://doi.org/10.1016/j.jstrokecerebrovasdis.2016.09.005.
2017 Arya	Arya KN, Pandian S, Kumar V. Effect of activity-based mirror therapy on lower limb motor-recovery and gait in stroke: A randomised controlled trial. Neuropsychol Rehabil 2019;29:1193–210. https://doi.org/10.1080/09602011.2017.1377087.
2017 Cha	Cha H-G, Shin Y-J, Kim M-K. Effects of the Bad Ragaz Ring Method on muscle activation of the lower limbs and balance ability in chronic stroke: A randomised controlled trial. Hong Kong Physiother J 2017;37:39–45. https://doi.org/10.1016/j.hkpj.2017.02.001.
2017 Choi	Choi H-S, Shin W-S, Bang D-H, Choi S-J. Effects of Game-Based Constraint-Induced Movement Therapy on Balance in Patients with Stroke: A Single-Blind Randomized Controlled Trial. Am J Phys Med Rehabil 2017;96:184–90. https://doi.org/10.1097/PHM.0000000000000567.
2017 De	De S, Chopra C, Mehta DD, Mehndiratta MM. Comparison between Mirror Therapy and Mental Imagery in Improving Ankle Motor Recovery in Sub Acute Stroke Patients. Indian J Physiother Occup Ther - Int J 2017;11:169. https://doi.org/10.5958/0973-5674.2017.00097.1.
2017 de Lima Gomes	de Lima Gomes W, de Nadai Dias LI, Guimarães RP, Stival CM, Faria G da R, Bovi ACN, et al. Effects of Treadmill Training in chronic hemiparetic: a randomized, double-blind clinical trial. Man Ther Posturology Rehabil J 2017;15:1–5. https://doi.org/10.17784/mtprehabjournal.2017.15.530.
2017 Dujović	Dujović SD, Malešević J, Malešević N, Vidaković AS, Bijelić G, Keller T, et al. Novel multi-pad functional electrical stimulation in stroke patients: A single-blind randomized study. NeuroRehabilitation 2017;41:791–800. https://doi.org/10.3233/NRE-172153.
2017 In	In T, Jin Y, Jung K, Cho H. Treadmill training with Thera-Band improves motor function, gait and balance in stroke patients. NeuroRehabilitation 2017;40:109–14. https://doi.org/10.3233/NRE-161395.
2017 Ivey FM	Ivey FM, Prior SJ, Hafer-Macko CE, Katzel LI, Macko RF, Ryan AS. Strength Training for Skeletal Muscle Endurance after Stroke. J Stroke Cerebrovasc Dis 2017;26:787–94. https://doi.org/10.1016/j.jstrokecerebrovasdis.2016.10.018.
2017 Jung	Jung K-S, In T-S, Cho H. Effects of sit-to-stand training combined with transcutaneous electrical stimulation on spasticity, muscle strength and balance ability in patients with stroke: A randomized controlled study. Gait Posture 2017;54:183–7. https://doi.org/10.1016/j.gaitpost.2017.03.007.
2017 Lee	Lee HJ, Kim YM, Lee DK. The effects of action observation training and mirror therapy on gait and balance in stroke patients. J Phys Ther Sci 2017;29:523–6. https://doi.org/10.1589/jpts.29.523.
2017 Nikamp	Nikamp CDM, Hobbelink MSH, van der Palen J, Hermens HJ, Rietman JS, Buurke JH. A randomized controlled trial on providing ankle-foot orthoses in patients with (sub-)acute stroke: Short-term kinematic and spatiotemporal effects and effects of timing. Gait Posture 2017;55:15–22. https://doi.org/10.1016/j.gaitpost.2017.03.028.

2017 Rasheeda	Vadnagarwala R, Ramachandran S. The Effect of Swiss Ball Therapy on Sit-to-Stand Function, Paretic Limb Weight Bearing and Lower Limb Motor Score in Patients with Hemiplegia. Int J Physiother 2017;4. https://doi.org/10.15621/ijphy/2017/v4i6/163917.
2017 Wang	Wang H, Zhao Z, Jiang P, Li X, Lin Q, Wu Q. Effect and mechanism of mirror therapy on rehabilitation of lower limb motor function in patients with stroke hemiplegia. Biomed Res India 2017;28:10165–70.
2017 Xu	Xu Q, Guo F, Salem HMA, Chen H, Huang X. Effects of mirror therapy combined with neuromuscular electrical stimulation on motor recovery of lower limbs and walking ability of patients with stroke: a randomized controlled study. Clin Rehabil 2017;31:1583–91. https://doi.org/10.1177/0269215517705689.
2018 Dalal	Dalal KK, Joshua AM, Nayak A, Mithra P, Misri Z, Unnikrishnan B. Effectiveness of prowling with proprioceptive training on knee hyperextension among stroke subjects using videographic observation- a randomised controlled trial. Gait Posture 2018;61:232–7. https://doi.org/10.1016/j.gaitpost.2018.01.018.
2018 Du	Du J, Yang F, Hu J, Hu J, Xu Q, Cong N, et al. Effects of high- and low-frequency repetitive transcranial magnetic stimulation on motor recovery in early stroke patients: Evidence from a randomized controlled trial with clinical, neurophysiological and functional imaging assessments. NeuroImage Clin 2019;21:101620. https://doi.org/10.1016/j.nicl.2018.101620.
2018 Dubey	Dubey L, Karthikbabu S, Mohan D. Effects of Pelvic Stability Training on Movement Control, Hip Muscles Strength, Walking Speed and Daily Activities after Stroke: A Randomized Controlled Trial. Ann Neurosci 2018;25:80–9. https://doi.org/10.1159/000486273.
2018 Elhakk	Elhakk SMA, Ragab WM, Zakaria HM, Faggal MS, Taha SI, Badawy WM, et al. Task oriented approach via virtual reality for improving postural control in stroke patients. Biosci Res 2018;15:3926–33.
2018 Erbil	Erbil D, Tugba G, Murat TH, Melike A, Merve A, Cagla K, et al. Effects of robot-assisted gait training in chronic stroke patients treated by botulinum toxin-a: A pivotal study. Physiother Res Int 2018;23:e1718. https://doi.org/10.1002/pri.1718.
2018 Ferreira	Ferreira LAB, Cimolin V, Neto HP, Grecco LAC, Lazzari RD, Dumont AJL, et al. Effect of postural insoles on gait pattern in individuals with hemiparesis: A randomized controlled clinical trial. J Bodyw Mov Ther 2018;22:792–7. https://doi.org/10.1016/j.jbmt.2017.08.004.
2018 Ganesh	Ganesh GS, Kumari R, Pattnaik M, Mohanty P, Mishra C, Kaur P, et al. Effectiveness of Faradic and Russian currents on plantar flexor muscle spasticity, ankle motor recovery, and functional gait in stroke patients. Physiother Res Int 2018;23:e1705. https://doi.org/10.1002/pri.1705.
2018 Ghasemi (a)	Ghasemi E, Khademi-Kalantari K, Khalkhali-Zavieh M, Rezasoltani A, Ghasemi M, Baghban AA, et al. The effect of functional stretching exercises on functional outcomes in spastic stroke patients: A randomized controlled clinical trial. J Bodyw Mov Ther 2018;22:1004–12. https://doi.org/10.1016/j.jbmt.2017.09.021.
2018 Ghasemi (b)	Ghasemi E, Khademi-Kalantari K, Khalkhali-Zavieh M, Rezasoltani A, Ghasemi M, Akbarzadeh Baghban A, et al. The Effect of Functional Stretching Exercises on Neural and Mechanical Properties of the Spastic Medial Gastrocnemius Muscle in Patients with Chronic Stroke: A Randomized Controlled Trial. J Stroke Cerebrovasc Dis Off J Natl Stroke Assoc 2018;27:1733–42. https://doi.org/10.1016/j.jstrokecerebrovasdis.2018.01.024.
2018 Huang	Huang Y-Z, Lin L-F, Chang K-H, Hu C-J, Liou T-H, Lin Y-N. Priming With 1-Hz Repetitive Transcranial Magnetic Stimulation Over Contralesional Leg Motor Cortex Does Not Increase the Rate of Regaining Ambulation Within 3 Months of Stroke: A Randomized Controlled Trial. Am J Phys Med Rehabil 2018;97:339–45. https://doi.org/10.1097/PHM.0000000000000850.
2018 Kim	Kim M-K, Choe Y-W, Shin Y-J, Peng C, Choi E-H. Effect of mirror use on lower extremity muscle strength of patients with chronic stroke. J Phys Ther Sci 2018;30:213–5. https://doi.org/10.1589/jpts.30.213.
2018 Klomjai	Klomjai W, Aneksan B, Pheungphrarattanatrai A, Chantanachai T, Choowong N, Bunleukhet S, et al. Effect of single-session dual-tDCS before physical therapy on lower-limb performance in sub-acute stroke patients: A randomized sham-controlled crossover study. Ann Phys Rehabil Med 2018;61:286–91. https://doi.org/10.1016/j.rehab.2018.04.005.

2018 Lee	Lee S, Lee K, Song C. Gait Training with Bilateral Rhythmic Auditory Stimulation in Stroke Patients: A Randomized Controlled Trial. Brain Sci 2018;8:164. https://doi.org/10.3390/brainsci8090164.
2018 Maciaszek	Maciaszek J. Effects of Posturographic Platform Biofeedback Training on the Static and Dynamic Balance of Older Stroke Patients. J Stroke Cerebrovasc Dis 2018;27:1969–74. https://doi.org/10.1016/j.jstrokecerebrovasdis.2018.02.055.
2018 Mainka	Mainka S, Wissel J, Völler H, Evers S. The Use of Rhythmic Auditory Stimulation to Optimize Treadmill Training for Stroke Patients: A Randomized Controlled Trial. Front Neurol 2018;9:755. https://doi.org/10.3389/fneur.2018.00755.
2018 Nikamp	Nikamp CDM, van der Palen J, Hermens HJ, Rietman JS, Buurke JH. The influence of early or delayed provision of ankle-foot orthoses on pelvis, hip and knee kinematics in patients with sub-acute stroke: A randomized controlled trial. Gait Posture 2018;63:260–7. https://doi.org/10.1016/j.gaitpost.2018.05.012.
2018 Pan	Pan X-L. Efficacy of early rehabilitation therapy on movement ability of hemiplegic lower extremity in patients with acute cerebrovascular accident. Medicine (Baltimore) 2018;97:e9544. https://doi.org/10.1097/MD.0000000000009544.
2018 Park	Park D, Lee J-H, Kang T-W, Cynn H-S. Four-week training involving ankle mobilization with movement versus static muscle stretching in patients with chronic stroke: a randomized controlled trial. Top Stroke Rehabil 2019;26:81–6. https://doi.org/10.1080/10749357.2018.1550614.
2018 Son	Son Y-L, Kim J-W. The effects of mirror neuron system-based self-observation training on lower limb muscle activity and dynamic balance in patients with chronic stroke. J Phys Ther Sci 2018;30:1241–4. https://doi.org/10.1589/jpts.30.1241.
2018 Tsaih	Tsaih P-L, Chiu M-J, Luh J-J, Yang Y-R, Lin J-J, Hu M-H. Practice Variability Combined with Task-Oriented Electromyographic Biofeedback Enhances Strength and Balance in People with Chronic Stroke. Behav Neurol 2018;2018:1–9. https://doi.org/10.1155/2018/7080218.
2018 Zhang	Zhang Y, Qin L, Shi Y, Zhong C. Effects of high-quality nursing services on the neurological functions and abilities of daily living of stroke patients. NeuroQuantology 2018;16:7–12. https://doi.org/10.14704/nq.2018.16.5.1327.
2018 Zhao	Zhao N, Zhang J, Qiu M, Wang C, Xiang Y, Wang H, et al. Scalp acupuncture plus low-frequency rTMS promotes repair of brain white matter tracts in stroke patients: A DTI study. J Integr Neurosci 2018;17:125–39. https://doi.org/10.3233/JIN-170043.
2018 Zheng	Zheng X, Chen D, Yan T, Jin D, Zhuang Z, Tan Z, et al. A Randomized Clinical Trial of a Functional Electrical Stimulation Mimic to Gait Promotes Motor Recovery and Brain Remodeling in Acute Stroke. Behav Neurol 2018;2018:1–10. https://doi.org/10.1155/2018/8923520.
2019 Ancona	Ancona E, Quarenghi A, Simonini M, Saggini R, Mazzoleni S, De Tanti A, et al. Effect of verticalization with Erigo® in the acute rehabilitation of severe acquired brain injury. Neurol Sci Off J Ital Neurol Soc Ital Soc Clin Neurophysiol 2019;40:2073–80. https://doi.org/10.1007/s10072-019-03917-0.
2019 Chung	Chung SH, Kim JH, Yong SY, Lee YH, Park JM, Kim SH, et al. Effect of Task-Specific Lower Extremity Training on Cognitive and Gait Function in Stroke Patients: A Prospective Randomized Controlled Trial. Ann Rehabil Med 2019;43:1–10. https://doi.org/10.5535/arm.2019.43.1.1.
2019 de Sousa	de Sousa DG, Harvey LA, Dorsch S, Varettas B, Jamieson S, Murphy A, et al. Two weeks of intensive sit-to-stand training in addition to usual care improves sit-to-stand ability in people who are unable to stand up independently after stroke: a randomised trial. J Physiother 2019;65:152–8. https://doi.org/10.1016/j.jphys.2019.05.007.
2019 dos Santos	dos Santos RBC, Galvão SCB, Frederico LMP, Amaral NSL, Carneiro MIS, de Moura Filho AG, et al. Cortical and spinal excitability changes after repetitive transcranial magnetic stimulation combined to physiotherapy in stroke spastic patients. Neurol Sci 2019;40:1199–207. https://doi.org/10.1007/s10072-019-03765-y.
2019 Gamez	Gámez AB, Hernandez Morante JJ, Martínez Gil JL, Esparza F, Martínez CM. The effect of surface electromyography biofeedback on the activity of extensor and dorsiflexor muscles in elderly adults: a randomized trial. Sci Rep 2019;9:13153. https://doi.org/10.1038/s41598-019-49720-x.

2019 Guan	Guan Y, Guo N, Gao H, Sun J, Sun L, Liu X, et al. Study on application of continuous nursing in rehabilitation period of stroke patients. Acta Medica Mediterr 2019;35:539–43. https://doi.org/10.19193/0393-6384_2019_1s_83.
2019 Handelzalts	Handelzalts S, Kenner-Furman M, Gray G, Soroker N, Shani G, Melzer I. Effects of Perturbation-Based Balance Training in Subacute Persons With Stroke: A Randomized Controlled Trial. Neurorehabil Neural Repair 2019;33:213–24. https://doi.org/10.1177/1545968319829453.
2019 Hsieh	Hsieh H-C. Training by Using an Adaptive Foot Switch and Video Games to Improve Balance and Mobility Following Stroke: A Randomised Controlled Trial. Brain Impair 2019;20:16–23. https://doi.org/10.1017/BrImp.2018.15.
2019 Kim HY	Kim HY, Shin J-H, Yang SP, Shin MA, Lee SH. Robot-assisted gait training for balance and lower extremity function in patients with infratentorial stroke: a single-blinded randomized controlled trial. J NeuroEngineering Rehabil 2019;16:99. https://doi.org/10.1186/s12984-019-0553-5.
2019 Kim JY	Kim J, Kim DY, Chun MH, Kim SW, Jeon HR, Hwang CH, et al. Effects of robot-(Morning Walk®) assisted gait training for patients after stroke: a randomized controlled trial. Clin Rehabil 2019;33:516–23. https://doi.org/10.1177/0269215518806563.
2019 Lee	Lee K. Speed-Interactive Pedaling Training Using Smartphone Virtual Reality Application for Stroke Patients: Single-Blinded, Randomized Clinical Trial. Brain Sci 2019;9:295. https://doi.org/10.3390/brainsci9110295.
2019 Nam	Nam Y-G, Lee JW, Park JW, Lee HJ, Nam KY, Park JH, et al. Effects of Electromechanical Exoskeleton-Assisted Gait Training on Walking Ability of Stroke Patients: A Randomized Controlled Trial. Arch Phys Med Rehabil 2019;100:26–31. https://doi.org/10.1016/j.apmr.2018.06.020.
2019 Nordin	Nordin NAM, Aziz NA, Sulong S, Aljunid SM. Effectiveness of home-based carer-assisted in comparison to hospital-based therapist-delivered therapy for people with stroke: A randomised controlled trial. NeuroRehabilitation 2019;45:87–97. https://doi.org/10.3233/NRE-192758.
2019 Park D	Park D, Cynn H-S, Yi C, Choi WJ, Shim J-H, Oh D-W. Four-week training involving self-ankle mobilization with movement versus calf muscle stretching in patients with chronic stroke: a randomized controlled study. Top Stroke Rehabil 2020;27:296–304. https://doi.org/10.1080/10749357.2019.1690831.
2019 Park J-S	Park J-S, Choi J-B, Lee G, Lee S-H, Jung Y. EFFECT OF MOTOR IMAGERY TRAINING IN COMBINATION WITH ELECTROMYOGRAPHY-TRIGGERED ELECTRICAL STIMULATION IN STROKE WITH HEMIPLEGIA PATIENTS: A RANDOMIZED CONTROLLED STUDY. J Mech Med Biol 2019;19:1940062. https://doi.org/10.1142/S0219519419400621.
2019 Pradines	Pradines M, Ghedira M, Portero R, Masson I, Marciniak C, Hicklin D, et al. Ultrasound Structural Changes in Triceps Surae After a 1-Year Daily Self-stretch Program: A Prospective Randomized Controlled Trial in Chronic Hemiparesis. Neurorehabil Neural Repair 2019;33:245–59. https://doi.org/10.1177/1545968319829455.
2019 Saadatnia	Saadatnia M, Shahnazi H, Khorvash F, Esteki-Ghashghaei F. The Impact of Home-Based Exercise Rehabilitation on Functional Capacity in Patients With Acute Ischemic Stroke: A Randomized Controlled Trial. Home Health Care Manag Pract 2020;32:141–7. https://doi.org/10.1177/1084822319895982.
2019 Sczesny-Kaiser	Sczesny-Kaiser M, Trost R, Aach M, Schildhauer TA, Schwenkreis P, Tegenthoff M. A Randomized and Controlled Crossover Study Investigating the Improvement of Walking and Posture Functions in Chronic Stroke Patients Using HAL Exoskeleton – The HALESTRO Study (HAL-Exoskeleton STROke Study). Front Neurosci 2019;13:259. https://doi.org/10.3389/fnins.2019.00259.
2019 Sheng	Sheng Y, Kan S, Wen Z, Chen W, Qi Q, Qu Q, et al. Effect of Kinesio Taping on the Walking Ability of Patients with Foot Drop after Stroke. Evid Based Complement Alternat Med 2019;2019:1–7. https://doi.org/10.1155/2019/2459852.
2019 Stolz	Stolz R, Nayyar R, Louie J, Bower KJ, Paul SK, Ng L. The effectiveness of a novel cable-driven gait trainer (Robowalk) combined with conventional physiotherapy compared to conventional physiotherapy alone following stroke: a randomised controlled trial. Int J Rehabil Res 2019;42:377–84. https://doi.org/10.1097/MRR.0000000000000375.

2019 Wang	Wang M, Liu S, Peng Z, Zhu Y, Feng X, Gu Y, et al. Tibetan Medicated Bathing Therapy for Patients With Post-stroke Limb Spasticity: A Randomized Controlled Clinical Trial. J Am Med Dir Assoc 2020;21:374-380.e4. https://doi.org/10.1016/j.jamda.2019.10.018.
2019 Zhang	Zhang X, Sheng Y, Gao S. Effects of early rehabilitation nursing on improvement of conditions and quality of life in patients after ischemic strokes. Int J Clin Exp Med 2019;12:11412–9.
2020 Ambrosini	Ambrosini E, Peri E, Nava C, Longoni L, Monticone M, Pedrocchi A, et al. A multimodal training with visual biofeedback in subacute stroke survivors: a randomized controlled trial. Eur J Phys Rehabil Med 2020;56. https://doi.org/10.23736/S1973-9087.19.05847-7.
2020 Chen	Chen X, Gan Z, Tian W, Lv Y. Effects of rehabilitation training of core muscle stability on stroke patients with hemiplegia. Pak J Med Sci 2020;36. https://doi.org/10.12669/pjms.36.3.1466.
2020 Kumar	Kumar S, Yadav R, Aafreen. Comparison between Erigo tilt-table exercise and conventional physiotherapy exercises in acute stroke patients: a randomized trial. Arch Physiother 2020;10:3. https://doi.org/10.1186/s40945-020-0075-2.
2020 Lee	Lee K. Balance Training with Electromyogram-Triggered Functional Electrical Stimulation in the Rehabilitation of Stroke Patients. Brain Sci 2020;10:80. https://doi.org/10.3390/brainsci10020080.
2020 Liu	Liu J, Feng W, Zhou J, Huang F, Long L, Wang Y, et al. Effects of sling exercise therapy on balance, mobility, activities of daily living, quality of life and shoulder pain in stroke patients: a randomized controlled trial. Eur J Integr Med 2020;35:101077. https://doi.org/10.1016/j.eujim.2020.101077.
2020 Mustafaouglu	Mustafaoglu R, Erhan B, Yeldan I, Gunduz B, Tarakci E. Does robot-assisted gait training improve mobility, activities of daily living and quality of life in stroke? A single-blinded, randomized controlled trial. Acta Neurol Belg 2020;120:335–44. https://doi.org/10.1007/s13760-020-01276-8.
2020 Sharma	Sharma H, Vishnu VY, Kumar N, Sreenivas V, Rajeswari MR, Bhatia R, et al. Efficacy of Low-Frequency Repetitive Transcranial Magnetic Stimulation in Ischemic Stroke: A Double-Blind Randomized Controlled Trial. Arch Rehabil Res Clin Transl 2020;2:100039. https://doi.org/10.1016/j.arrct.2020.100039.
2020 Sürücü	Dost Sürücü G, Tezen Ö. The effect of EMG biofeedback on lower extremity functions in hemiplegic patients. Acta Neurol Belg 2020. https://doi.org/10.1007/s13760-019-01261-w.
2020 Swank	Swank C, Trammell M, Callender L, Bennett M, Patterson K, Gillespie J, et al. The impact of a patient-directed activity program on functional outcomes and activity participation after stroke during inpatient rehabilitation—a randomized controlled trial. Clin Rehabil 2020;34:504–14. https://doi.org/10.1177/0269215519901153.
2020 Wang	Wang H-Q, Hou M, Li H, Bao C-L, Min L, Dong G-R, et al. Effects of acupuncture treatment on motor function in patients with subacute hemorrhagic stroke: A randomized controlled study. Complement Ther Med 2020;49. https://doi.org/10.1016/j.ctim.2019.102296.
2020 Yadav	Yadav R, Kumar S, Afrin A. EFFICACY OF ERIGO TILT-TABLE AND CONVENTIONAL PHYSIOTHERAPY ON REHABILITATION OUTCOME WITHIN HAEMORRHAGIC AND ISCHEMIC STROKE PATIENTS. J Musculoskelet Res 2019;22:1950003. https://doi.org/10.1142/S0218957719500039.