APPENDIX

Review

Sequelae of COVID-19 among previously hospitalized patients up to 1 year after discharge: a systematic review and meta-analysis

Tianqi Yang[a*], Michael Zhipeng Yan[b*], Xingyi Li[a], Eric H Y Lau[a,c†]

Affiliation:

[a]School of Public Health, The University of Hong Kong, Hong Kong, China

[b]Department of Medicine, Li Ka Shing Faculty of Medicine, The University of Hong Kong, Hong Kong, China

[c]Laboratory of Data Discovery for Health Limited, Hong Kong Science and Technology Park, Hong Kong, China

School of Public Health, The University of Hong Kong, Hong Kong, China

Email addresses: ehylau@hku.hk (Eric HY Lau)

†Corresponding author:

Dr Eric Lau, School of Public Health, Li Ka Shing Faculty of Medicine, The University of Hong Kong, 7 Sassoon Road, Pokfulam, Hong Kong.

Tel: +852 3917 6719; Fax: +852 3520 1945; email: ehylau@hku.hk

Appendix 1 The search queries of Web of Science and PubMed

Database	**Search query**
PubMed	(COVID-19[TIAB] OR SARS-CoV-2[TIAB] OR 2019-nCoV[TIAB]) AND (((persistent[TIAB] OR post[TIAB] OR long term[TIAB]) AND (effect[TIAB] OR consequence[TIAB] OR symptom[TIAB] OR manifestations[TIAB] OR syndrome[TIAB])) OR sequelae[TIAB]) AND (study[TIAB] OR trial[TIAB])
Web of Science	ALL=((COVID-19 OR SARS-CoV-2 OR 2019-nCoV) AND (((persistent OR post OR long term) AND (effect OR consequence OR symptom OR manifestations OR syndrome)) OR sequelae) AND (study OR trial))

Appendix 2 Overall star rating for studies by NOS

Study	Selection	Comparability	Outcome	Overall star rating
Xiong et al.	++++	++	+	7
Fernández-de-las-Peñas et al.	+++		+++	6
Bellan et al.	+++		++	5
Romero-Duarte et al.	+++		+++	6
Simani et al.	+++		++	5
Mei et al.	++		++	4
Qu et al.	++		++	4
Shang et al.	++		++	4
Suárez-Robles et al.	++		++	4
Huang et al.	+++		++	5
Tarsitani et al.	+++		++	5
Froidure et al.	+++		++	5
Morin et al.	+++		++	5
Ayoubkhani et al.	+++	++	++	7
L. Sykes et al.	+++		++	5
Arnold et al.	+++		+++	6
Garrigues et al.	+++		++	5
Caruso et al.	+++		++	5
Castro et al.	+++	++	++	7
Darcis et al.	+++		+	4
González-Hermosillo et al.	+++		++	5
Gramaglia et al.	+++		++	5
Bozzetti et al.	+++		++	5
Maestre-Muñiz et al.	+++		++	5
Vincent et al.	+++		++	5
Sun et al.	+++		++	5
Vlake et al.	+++	++	++	7
Huang et al.	+++	++	++	7
Li et al.	+++		++	5
Sibila et al.	+++		++	5
Liu et al.	+++		++	5
Lombardo et al.	+++		++	5
Maestrini et al.	+++		+	4
Sigfrid et al.	++		++	4
Ahmed et al.	++		++	4
Anjana et al.	+++		++	5

Bai et al.	++		++	4
Chen et al.	++		++	4
Damiano et al.	++		++	4
Eloy et al.	++		++	4
Evans et al.	++		++	4
Gamberini et al.	++		++	4
Garcia-Abellan et al.	++		++	4
Ghosn et al.	++		++	4
Hodgson et al.	++		++	4
Horwitz et al.	++		+++	5
Huang et al.	++		++	4
Huarcaya-Victoria et al.	++		++	4
Karaarslan et al.	++		+++	5
Boglione et al.	++		+++	5
Chand et al.	+++		++	5
Kumar et al.	+++		+++	6
Mei et al.	++		+++	5
Nesan et al.	+++		++	5
Ong et al.	+++		++	5
Rass et al.	++		++	4
Righi et al.	++		+++	5
Rivera-Izquierdo et al.	++	++	++	6
Sibila et al.	+++		++	5
Staudt et al.	++		++	4
Tessitore et al.	+++		++	5
Tleyjeh et al.	++		++	4
Torres-Ruiz et al.	+++		++	5
Vejen et al.	++		++	4
Wang et al.	++		++	4
Wong-Chew et al.	++		++	4
Zhang et al.	+++		++	5
Zhou et al.	++		++	4
Zuschlag et al.	++		++	4

Appendix 3 Pooled prevalence of sequelae at 3 - 4, 5 - 8 and ≥ 9 months (9 - 13 months) follow-up

	3 - 4 months	5 - 8 months	≥ 9 months
Fatigue			
Prevalence %	26.4	29.4	26.2
95% CI %	[19.6 - 34.5]	[19.8 - 41.3]	[17.3 - 37.6]
Range %	(1.8 - 55.0)	(1.5 - 84.9)	(3.7 - 68.5)
I^2	98.3%	99.0%	98.9%
P_H	< 0.001	< 0.001	< 0.001
The number of studies	19	20	14
Time Days	107	169	349
P-value of meta regression		0.743	0.985
Somnipathy			
Prevalence %	19.2	27.1	15.1
95% CI %	[11.4 - 30.5]	[15.3 - 43.4]	[9.3 - 23.6]
Range %	(3.6 - 53.6)	(2.1 - 64.8)	(1.2 - 47.1)
I^2	98.6%	99.2%	97.3%
P_H	< 0.001	< 0.001	< 0.001
The number of studies	11	10	9
Time Days	109	162	367
P-value of meta regression		0.414	0.522
Anxiety			
Prevalence %	20.9	17.9	12.9
95% CI %	[13.4 - 31.2]	[11.3 - 27.2]	[7.7 - 20.9]
Range %	(6.1 - 47.8)	(6.8 - 35.4)	(0.6 - 41.7)
I^2	98.4%	96.8%	96.4%
P_H	< 0.001	< 0.001	< 0.001
The number of studies	11	8	5
Time Days	110	183	373
P-value of meta regression		0.605	0.227
Dyspnea			
Prevalence %	14.6	18.5	13.1
95% CI %	[7.9 - 25.3]	[11.3 - 28.6]	[6.5 - 24.8]
Range %	(1.0 - 55.0)	(0.8 - 47.7)	(2.7 - 58.4)
I^2	98.1%	98.4%	98.6%
P_H	< 0.001	< 0.001	< 0.001
The number of studies	14	12	9
Time Days	94	169	350
P-value of meta regression		0.645	0.845
PTSD			
Prevalence %	15.2	16.0	1.2
95% CI %	[11.0 - 20.7]	[10.9 - 22.8]	[0.2 - 4.4]
Range %	(7.4 - 28.6)	(5.8 - 32.0)	-
I^2	84.5%	94.4%	-
P_H	< 0.001	< 0.001	-

The number of studies	8	9	1
Time Days	100	186	360
P-value of meta regression		0.854	< 0.001
Hypomnesia			
Prevalence %	9.9	21.8	4.9
95% CI %	[3.5 - 25.2]	[14.0 - 32.3]	[0.7 - 27.6]
Range %	(1.7 - 37.3)	(0.8 - 53.8)	(0.6 - 22.4)
I^2	98.8%	97.9%	96.0%
P_H	< 0.001	< 0.001	< 0.001
The number of studies	7	10	3
Time Days	115	208	358
P-value of meta regression		0.176	0.527
Arthralgia			
Prevalence %	7.7	18.0	11.5
95% CI %	[4.3 - 13.2]	[9.4 - 31.7]	[3.8 - 29.8]
Range %	(2.6 - 24.3)	(0.0 - 47.8)	(2.9 - 32.5)
I^2	94.2%	99.0%	97.2%
P_H	< 0.001	< 0.001	< 0.001
The number of studies	7	11	3
Time Days	92	163	350
P-value of meta regression		0.064	0.605
Depression			
Prevalence %	14.9	10.9	12.6
95% CI %	[9.0 - 23.7]	[5.8 - 19.6]	[5.8 - 25.1]
Range %	(4.3 - 30.2)	(2.6 - 34.6)	(0.6 - 37.5)
I^2	94.7%	97.1%	96.4%
P_H	< 0.001	< 0.001	< 0.001
The number of studies	8	9	5
Time Days	103	184	381
P-value of meta regression		0.411	0.635
Alopecia			
Prevalence %	27.9	7.0	10.5
95% CI %	[21.9 - 34.9]	[2.6 - 17.5]	[4.3 - 24.4]
Range %	(20.0 - 35.5)	(0.5 - 47.0)	(0.0 - 40.6)
I^2	77.8%	99.0%	97.5%
P_H	0.004	< 0.001	< 0.001
The number of studies	4	8	5
Time Days	94	157	352
P-value of meta regression		0.007	0.126
Myalgia			
Prevalence %	10.8	15.2	6.0
95% CI %	[5.5 - 20.1]	[8.3 - 26.2]	[3.6 - 9.6]
Range %	(3.2 - 51.5)	(0.2 - 57.2)	(1.0 - 15.5)
I^2	96.8%	98.7%	92.3%
P_H	< 0.001	< 0.001	< 0.001
The number of studies	9	12	6

Time Days	92	163	355
P-value of meta regression		0.583	0.182
Cough			
Prevalence %	11.1	12.6	4.6
95% CI %	[7.8 - 15.5]	[9.2 - 17.1]	[2.9 - 7.3]
Range %	(0.0 - 35.1)	(1.7 - 35.2)	(1.9 - 11.0)
I^2	92.3%	96.1%	90.2%
P_H	< 0.001	< 0.001	< 0.001
The number of studies	15	17	8
Time Days	95	174	350
P-value of meta regression		0.580	0.008
Headache			
Prevalence %	5.9	12.8	4.1
95% CI %	[3.6 - 9.6]	[7.9 - 20.2]	[2.5 - 6.7]
Range %	(1.6 - 24.6)	(0.0 - 39.4)	(0.6 - 10.3)
I^2	96.1%	97.7%	90.0%
P_H	< 0.001	< 0.001	< 0.001
The number of studies	12	14	7
Time Days	108	177	359
P-value of meta regression		0.050	0.353
Chest pain			
Prevalence %	7.3	10.0	3.0
95% CI %	[4.7 - 11.0]	[6.5 - 15.0]	[1.8 - 4.9]
Range %	(0.6 - 17.9)	(0.0 - 27.7)	(0.0 - 7.2)
I^2	90.3%	97.0%	87.2%
P_H	< 0.001	< 0.001	< 0.001
The number of studies	10	12	7
Time Days	95	163	347
P-value of meta regression		0.320	0.030
Dizziness			
Prevalence %	3.8	13.1	3.2
95% CI %	[1.9 - 7.5]	[6.6 - 24.4]	[2.1 - 5.0]
Range %	(1.8 - 6.8)	(1.9 - 35.8)	(0.6 - 5.1)
I^2	88.6%	98.3%	79.2%
P_H	< 0.001	< 0.001	0.002
The number of studies	4	9	4
Time Days	92	173	359
P-value of meta regression		0.018	0.626
Palpitation			
Prevalence %	7.0	7.7	5.0
95% CI %	[1.4 - 28.3]	[4.3 - 13.5]	[3.2 - 7.8]
Range %	(0.5 - 21.6)	(1.7 - 23.2)	(1.6 - 9.2)
I^2	98.0%	98.1%	89.8%
P_H	< 0.001	< 0.001	< 0.001
The number of studies	3	9	6
Time Days	90	181	359

P-value of meta regression		0.918	0.804
Ageusia			
Prevalence %	4.2	9.4	3.5
95% CI %	[2.2 - 7.8]	[5.1 - 16.9]	[1.3 - 9.1]
Range %	(0.0 - 21.6)	(2.7 - 49.9)	(0.0 - 20.8)
I^2	93.4%	98.2%	94.5%
P_H	< 0.001	< 0.001	< 0.001
The number of studies	11	13	7
Time Days	92	197	355
P-value of meta regression		0.041	0.789
Anosmia			
Prevalence %	4.0	10.9	3.2
95% CI %	[2.3 - 6.7]	[6.3 - 18.2]	[1.6 - 6.4]
Range %	(0.0 - 26.1)	(4.1 - 53.8)	(0.0 - 9.5)
I^2	94.8%	98.0%	92.3%
P_H	< 0.001	< 0.001	< 0.001
The number of studies	15	14	7
Time Days	107	197	354
P-value of meta regression		0.003	0.598
Sore throat			
Prevalence %	3.6	4.9	1.7
95% CI %	[1.2 - 9.8]	[2.9 - 8.2]	[0.7 - 4.0]
Range %	(0.7 - 12.7)	(0.0 - 13.3)	(0.0 - 3.5)
I^2	95.5%	92.6%	88.6%
P_H	< 0.001	< 0.001	< 0.001
The number of studies	5	6	4
Time Days	92	201	359
P-value of meta regression		0.619	0.333
Diarrhea			
Prevalence %	3.1	5.0	0.9
95% CI %	[1.6 - 6.0]	[2.6 - 9.1]	[0.4 - 2.1]
Range %	(0.6 - 6.7)	(0.0 - 17.7)	(0.0 - 3.3)
I^2	76.1%	96.9%	63.7%
P_H	0.002	< 0.001	0.027
The number of studies	5	11	5
Time Days	90	183	355
P-value of meta regression		0.403	0.040
Fever			
Prevalence %	1.1	1.9	0.2
95% CI %	[0.2 - 4.8]	[0.9 - 4.3]	[0.0 - 0.8]
Range %	(0.0 - 10.4)	(0.0 - 11.0)	(0.0 - 0.6)
I^2	88.5%	91.0%	22.0%
P_H	< 0.001	< 0.001	0.279
The number of studies	6	8	4
Time Days	95	173	363

P-value of meta regression	0.702	0.079

Appendix 4 Pooled prevalence of symptoms in Asia, Europe and other regions

	Asia	Europe	Others
Fatigue			
Prevalence %	17.5	33.9	33.9
95% CI %	[11.7 - 25.5]	[27.2 - 41.2]	[18.0 - 54.4]
Range %	(1.5 - 62.7)	(5.9 - 82.8)	(10.9 - 84.9)
I^2	99.1%	97.7%	98.9%
P_H	< 0.001	< 0.001	< 0.001
The number of studies	18	29	6
P-value of meta regression		0.018	0.153
Somnipathy			
Prevalence %	12.3	24.1	30.3
95% CI %	[7.0 - 20.8]	[16.8 - 33.3]	[11.7 - 58.8]
Range %	(2.1 - 43.3)	(1.2 - 53.6)	(6.8 - 64.8)
I^2	99.0%	97.4%	99.4%
P_H	< 0.001	< 0.001	< 0.001
The number of studies	10	15	5
P-value of meta regression		0.118	0.104
Anxiety			
Prevalence %	24.1	17.8	16.5
95% CI %	[10.9 - 45.0]	[13.0 - 23.9]	[9.2 - 27.8]
Range %	(10.8 - 41.7)	(0.6 - 47.8)	(6.1 - 35.4)
I^2	99.0%	96.1%	97.9%
P_H	< 0.001	< 0.001	< 0.001
The number of studies	3	14	7
P-value of meta regression		0.467	0.441
Dyspnea			
Prevalence %	5.6	27.0	8.4
95% CI %	[2.2 - 13.4]	[22.5 - 32.1]	[4.8 - 14.1]
Range %	(0.8 - 55.0)	(4.3 - 58.4)	(6.7 - 11.7)
I^2	99.0%	94.9%	70.1%
P_H	< 0.001	< 0.001	0.068
The number of studies	12	21	2
P-value of meta regression		<0.001	0.428
PTSD			
Prevalence %	10.0	12.7	23.5
95% CI %	[6.8 - 14.4]	[8.5 - 18.7]	[16.5 - 32.1]
Range %	(5.8 - 16.0)	(1.2 - 30.8)	(13.6 - 32.0)
I^2	69.5%	92.7%	88.5%
P_H	0.020	< 0.001	< 0.001
The number of studies	4	9	5
P-value of meta regression		0.458	0.001
Hypomnesia			
Prevalence %	4.5	21.2	12.8
95% CI %	[2.1 - 9.4]	[15.1 - 28.9]	[2.7 - 44.0]

Range %	(0.6 - 12.1)	(3.0 - 42.0)	(1.7 - 53.8)
I^2	91.5%	96.1%	99.3%
P_H	< 0.001	< 0.001	< 0.001
The number of studies	4	11	5
P-value of meta regression		0.017	0.193
Arthralgia			
Prevalence %	8.8	16.2	20.0
95% CI %	[4.5 - 16.6]	[8.9 - 27.9]	[8.0 - 42.0]
Range %	(0.0 - 40.0)	(4.5 - 47.8)	(6.8 - 43.8)
I^2	98.5%	97.9%	97.2%
P_H	< 0.001	< 0.001	< 0.001
The number of studies	10	7	4
P-value of meta regression		0.248	0.163
Depression			
Prevalence %	14.1	11.9	14.6
95% CI %	[1.3 - 67.7]	[8.2 - 16.9]	[7.2 - 27.4]
Range %	(4.3 - 37.5)	(0.6 - 29.5)	(2.6 - 34.6)
I^2	98.8%	96.0%	95.2%
P_H	< 0.001	< 0.001	< 0.001
The number of studies	2	14	6
P-value of meta regression		0.871	0.978
Alopecia			
Prevalence %	7.5	12.4	17.3
95% CI %	[2.8 - 18.3]	[5.8 - 24.4]	[7.4 - 35.5]
Range %	(0.0 - 47.0)	(2.2 - 40.6)	(5.2 - 32.2)
I^2	99.2%	97.1%	93.3%
P_H	< 0.001	< 0.001	< 0.001
The number of studies	7	7	3
P-value of meta regression		0.452	0.302
Myalgia			
Prevalence %	3.6	19.4	21.4
95% CI %	[1.6 - 7.6]	[13.0 - 28.1]	[8.8 - 43.5]
Range %	(0.2 - 41.4)	(4.7 - 57.2)	(9.1 - 38.1)
I^2	98.4%	97.1%	97.2%
P_H	< 0.001	< 0.001	< 0.001
The number of studies	10	13	4
P-value of meta regression		< 0.001	0.002
Cough			
Prevalence %	6.7	12.1	10.9
95% CI %	[4.5 - 9.8]	[9.1 - 15.9]	[4.9 - 22.3]
Range %	(0.6 - 15.6)	(1.9 - 35.2)	(0.0 - 23.3)
I^2	95.9%	94.2%	90.3%
P_H	< 0.001	< 0.001	< 0.001
The number of studies	13	23	4

P-value of meta regression		0.026	0.410
Headache			
Prevalence %	3.0	8.6	16.8
95% CI %	[1.4 - 6.5]	[5.6 - 13.1]	[8.7 - 30.1]
Range %	(0.0 - 24.9)	(1.7 - 39.4)	(6.1 - 36.9)
I^2	97.3%	96.6%	98.1%
P_H	< 0.001	< 0.001	< 0.001
The number of studies	9	18	6
P-value of meta regression		0.017	< 0.001
Chest pain			
Prevalence %	4.7	8.1	9.3
95% CI %	[3.2 - 7.0]	[5.7 - 11.5]	[3.6 - 21.7]
Range %	(0.0 - 12.3)	(0.0 - 21.9)	(3.9 - 27.7)
I^2	94.4%	93.9%	94.3%
P_H	< 0.001	< 0.001	< 0.001
The number of studies	9	16	4
P-value of meta regression		0.117	0.153
Dizziness			
Prevalence %	4.3	6.9	19.8
95% CI %	[2.5 - 7.4]	[2.6 - 17.1]	[5.8 - 49.7]
Range %	(1.8 - 19.3)	(0.6 - 35.8)	(6.5 - 35.4)
I^2	95.9%	97.9%	98.1%
P_H	< 0.001	< 0.001	< 0.001
The number of studies	7	7	3
P-value of meta regression		0.495	0.019
Palpitation			
Prevalence %	4.1	9.5	11.3
95% CI %	[2.1 - 7.6]	[5.4 - 16.0]	[4.6 - 25.0]
Range %	(0.5 - 20.4)	(3.1 - 23.2)	(7.0 - 16.7)
I^2	98.2%	96.5%	80.2%
P_H	< 0.001	< 0.001	0.025
The number of studies	8	8	2
P-value of meta regression		0.087	0.089
Ageusia			
Prevalence %	2.4	8.2	5.8
95% CI %	[1.2 - 4.8]	[4.7 - 13.8]	[2.4 - 13.0]
Range %	(0.0 - 8.4)	(1.0 - 49.9)	(0.0 - 12.7)
I^2	95.6%	97.4%	89.3%
P_H	< 0.001	< 0.001	< 0.001
The number of studies	7	19	5
P-value of meta regression		0.004	0.109
Anosmia			
Prevalence %	2.8	8.2	5.7
95% CI %	[1.5 - 5.4]	[4.7 - 13.9]	[2.8 - 11.6]
Range %	(0.4 - 13.0)	(0.0 - 53.8)	(0.0 - 15.1)

I^2	96.6%	97.6%	94.5%
P_H	< 0.001	< 0.001	< 0.001
The number of studies	10	19	7
P-value of meta regression		0.007	0.141
Sore throat			
Prevalence %	2.6	3.9	10.3
95% CI %	[1.2 - 5.3]	[2.2 - 6.8]	[8.5 - 12.5]
Range %	(0.0 - 13.3)	(0.0 - 12.7)	-
I^2	95.6%	87.2%	-
P_H	< 0.001	< 0.001	-
The number of studies	7	7	1
P-value of meta regression		0.490	0.001
Diarrhea			
Prevalence %	1.6	3.5	7.3
95% CI %	[0.6 - 4.1]	[1.7 - 6.9]	[2.6 - 18.5]
Range %	(0.0 - 6.7)	(0.0 - 17.7)	(3.2 - 14.6)
I^2	94.9%	96.0%	93.1%
P_H	< 0.001	< 0.001	< 0.001
The number of studies	8	10	3
P-value of meta regression		0.335	0.031
Fever			
Prevalence %	0.3	2.1	0.9
95% CI %	[0.0 - 2.8]	[1.1 - 4.1]	[0.5 - 1.8]
Range %	(0.0 - 2.8)	(0.0 - 11.0)	(0.0 - 1.0)
I^2	88.2%	84.6%	0.0%
P_H	< 0.001	< 0.001	0.628
The number of studies	4	12	2
P-value of meta regression		0.163	0.292

Appendix 5 Pooled prevalence of symptoms in < 60 years old and ≥ 60 years old

	< 60 years old		**≥ 60 years old**
Fatigue			
Prevalence %	28.1		26.9
95% CI %	[18.7 - 39.9]		[20.6 - 34.4]
Range %	(1.5 - 82.8)		(3.7 - 84.9)
I^2	99.0%		98.8%
P_H	< 0.001		< 0.001
The number of studies	21		29
P-value of meta regression		0.910	
Somnipathy			
Prevalence %	27.7		15.3
95% CI %	[18.6 - 39.2]		[9.2 - 24.3]
Range %	(2.1 - 64.8)		(1.2 -53.6)
I^2	98.8%		98.8%
P_H	< 0.001		< 0.001
The number of studies	15		14
P-value of meta regression		0.088	
Anxiety			
Prevalence %	23.2		12.7
95% CI %	[16.6 - 31.4]		[8.8 - 18.1]
Range %	(6.1 - 47.8)		(0.6 - 33.1)
I^2	95.9%		97.4%
P_H	< 0.001		< 0.001
The number of studies	12		11
P-value of meta regression		0.029	
Dyspnea			
Prevalence %	11.1		20.2
95% CI %	[4.9 - 23.4]		[14.9 - 26.8]
Range %	(0.8 - 46.0)		(2.7 - 58.4)
I^2	98.7%		97.9%
P_H	< 0.001		< 0.001
The number of studies	11		21
P-value of meta regression		0.178	
PTSD			
Prevalence %	14.5		13.5
95% CI %	[10.9 - 19.0]		[6.0 - 27.5]
Range %	(5.8 - 32.0)		(1.2 - 30.8)
I^2	91.4%		91.9%
P_H	< 0.001		< 0.001
The number of studies	13		4
P-value of meta regression		0.777	
Hypomnesia			
Prevalence %	16.6		12.0
95% CI %	[7.5 - 32.7]		[6.6 - 20.8]
Range %	(0.8 - 53.8)		(0.6 - 41.3)
I^2	98.3%		98.6%
P_H	< 0.001		< 0.001

The number of studies	7		13
P-value of meta regression		0.650	
Arthralgia			
Prevalence %	13.3		11.6
95% CI %	[7.7 - 22.2]		[6.6 - 19.6]
Range %	(0.0 - 47.8)		(2.9 - 33.3)
I^2	98.8%		94.5%
P_H	< 0.001		< 0.001
The number of studies	15		6
P-value of meta regression		0.766	
Depression			
Prevalence %	15.1		9.9
95% CI %	[10.2 - 21.7]		[6.1 - 15.8]
Range %	(2.6 - 37.5)		(0.6 - 29.5)
I^2	95.4%		95.1%
P_H	< 0.001		< 0.001
The number of studies	13		9
P-value of meta regression		0.241	
Alopecia			
Prevalence %	19.0		6.8
95% CI %	[9.4 - 34.5]		[3.2 - 14.1]
Range %	(1.8 - 47.0)		(0.0 - 40.6)
I^2	99.2%		96.9%
P_H	< 0.001		< 0.001
The number of studies	7		10
P-value of meta regression		0.092	
Myalgia			
Prevalence %	10.4		12.3
95% CI %	[4.9 - 21.0]		[6.9 - 21.1]
Range %	(0.2 - 57.2)		(1.0 - 38.1)
I^2	99.0%		98.0%
P_H	< 0.001		< 0.001
The number of studies	15		11
P-value of meta regression		0.744	
Cough			
Prevalence %	9.7		9.6
95% CI %	[6.3 - 14.6]		[7.1 - 12.7]
Range %	(0.0 - 35.1)		(1.9 - 35.2)
I^2	96.3%		95.3%
P_H	< 0.001		< 0.001
The number of studies	16		23
P-value of meta regression		0.995	
Headache			
Prevalence %	10.1		6.6
95% CI %	[5.8 - 17.0]		[4.7 - 9.1]
Range %	(0.0 - 39.4)		(0.6 - 26.2)
I^2	98.0%		94.4%
P_H	< 0.001		< 0.001
The number of studies	15		17

P-value of meta regression		0.248	
Chest pain			
Prevalence %	9.5		6.0
95% CI %	[6.5 - 13.8]		[4.2 - 8.5]
Range %	(0.0 - 27.7)		(0.0 - 20.5)
I^2	96.4%		91.2%
P_H	< 0.001		< 0.001
The number of studies	13		15
P-value of meta regression		0.083	
Dizziness			
Prevalence %	10.2		3.7
95% CI %	[5.1 - 19.4]		[1.5 - 9.1]
Range %	(1.8 - 35.8)		(0.6 - 28.6)
I^2	98.6%		96.0%
P_H	< 0.001		< 0.001
The number of studies	10		7
P-value of meta regression		0.081	
Palpitation			
Prevalence %	12.1		5.3
95% CI %	[6.5 - 21.4]		[3.8 - 7.2]
Range %	(1.7 - 23.2)		(1.6 - 16.7)
I^2	98.5%		86.9%
P_H	< 0.001		< 0.001
The number of studies	7		10
P-value of meta regression		0.036	
Ageusia			
Prevalence %	5.9		6.0
95% CI %	[4.0 - 8.7]		[2.8 - 12.2]
Range %	(0.0 - 21.6)		(0.0 - 49.9)
I^2	92.1%		98.1%
P_H	< 0.001		< 0.001
The number of studies	13		17
P-value of meta regression		0.813	
Anosmia			
Prevalence %	6.4		6.0
95% CI %	[4.5 - 9.1]		[2.9 - 11.8]
Range %	(0.0 - 26.1)		(0.0 - 53.8)
I^2	92.8%		98.5%
P_H	< 0.001		< 0.001
The number of studies	16		18
P-value of meta regression		0.986	
Sore throat			
Prevalence %	6.1		2.4
95% CI %	[3.6 - 10.1]		[1.3 - 4.4]
Range %	(0.0 - 13.3)		(0.0 - 5.9)
I^2	93.8%		89.8%
P_H	< 0.001		< 0.001
The number of studies	7		7
P-value of meta regression		0.030	

Diarrhea			
Prevalence %	4.3		1.8
95% CI %	[2.3 - 8.2]		[0.8 - 4.1]
Range %	(0.0 - 17.7)		(0.0 - 10.3)
I^2	96.3%		94.7%
P_H	< 0.001		< 0.001
The number of studies	11		10
P-value of meta regression		0.190	
Fever			
Prevalence %	1.3		0.8
95% CI %	[0.4 - 3.8]		[0.3 - 2.3]
Range %	(0.0 - 11.0)		(0.0 - 7.0)
I^2	92.4%		85.5%
P_H	< 0.001		< 0.001
The number of studies	9		9
P-value of meta regression		0.518	

Appendix 6 Pooled prevalence of symptoms in ICU admission < 20% and ≥ 20%

	ICU admission < 20%		ICU admission ≥ 20%
Fatigue			
Prevalence %	29.7		35.1
95% CI %	[21.1 - 40.0]		[25.2 - 46.4]
Range %	(1.7 - 68.5)		(8.2 - 84.9)
I^2	99.4%		97.5%
P_H	< 0.001		< 0.001
The number of studies	19		14
P-value of meta regression		0.527	
Somnipathy			
Prevalence %	15.0		29.4
95% CI %	[9.3 - 23.2]		[19.4 - 42.0]
Range %	(1.2 - 53.6)		(3.8 - 53.6)
I^2	99.1%		96.4%
P_H	< 0.001		< 0.001
The number of studies	13		8
P-value of meta regression		0.083	
Anxiety			
Prevalence %	13.3		18.9
95% CI %	[9.0 - 19.2]		[9.1 - 35.0]
Range %	(0.6 - 33.1)		(7.3 - 47.8)
I^2	98.3%		96.5%
P_H	< 0.001		< 0.001
The number of studies	12		6
P-value of meta regression		0.348	
Dyspnea			
Prevalence %	11.5		30.5
95% CI %	[6.9 - 18.5]		[22.5 - 39.9]
Range %	(1.0 - 40.3)		(11.0 - 58.4)
I^2	98.6%		95.6%
P_H	< 0.001		< 0.001
The number of studies	13		10
P-value of meta regression		0.005	
PTSD			
Prevalence %	9.2		16.7
95% CI %	[3.5 - 21.9]		[11.1 - 24.4]
Range %	(1.2 - 30.8)		(9.7 - 32.0)
I^2	94.8%		83.1%
P_H	< 0.001		< 0.001
The number of studies	5		5
P-value of meta regression		0.247	
Hypomnesia			
Prevalence %	5.1		17.1
95% CI %	[1.7 - 14.6]		[9.1 - 29.7]
Range %	(0.6 - 35.6)		(3.0 - 41.3)
I^2	99.3%		96.2%
P_H	< 0.001		< 0.001

The number of studies	6		7
P-value of meta regression		0.115	
Arthralgia			
Prevalence %	9.5		17.6
95% CI %	[6.7 - 13.2]		[9.8 - 29.5]
Range %	(0.0 - 17.6)		(4.7 - 37.0)
I^2	91.0%		96.1%
P_H	< 0.001		< 0.001
The number of studies	6		5
P-value of meta regression		0.751	
Depression			
Prevalence %	11.3		9.2
95% CI %	[7.0 - 17.7]		[4.6 - 17.7]
Range %	(0.6 - 29.5)		(2.6 - 20.6)
I^2	95.2%		92.2%
P_H	< 0.001		< 0.001
The number of studies	9		6
P-value of meta regression		0.721	
Alopecia			
Prevalence %	14.9		4.1
95% CI %	[9.2 - 23.2]		[0.5 - 27.4]
Range %	(0.0 -40.6)		(0.5 - 20.0)
I^2	98.0%		96.3%
P_H	< 0.001		< 0.001
The number of studies	8		3
P-value of meta regression		0.342	
Myalgia			
Prevalence %	5.6		29.4
95% CI %	[3.4 - 8.9]		[21.3 - 39.0]
Range %	(1.0 -23.1)		(7.9 - 51.5)
I^2	96.3%		92.0%
P_H	< 0.001		< 0.001
The number of studies	9		6
P-value of meta regression		< 0.001	
Cough			
Prevalence %	7.9		14.2
95% CI %	[5.0 - 12.2]		[11.2 - 17.9]
Range %	(1.7 - 26.9)		(5.0 - 35.1)
I^2	96.9%		86.5%
P_H	< 0.001		< 0.001
The number of studies	13		13
P-value of meta regression		0.033	
Headache			
Prevalence %	5.5		8.5
95% CI %	[3.5 - 8.5]		[4.5 - 15.7]
Range %	(0.0 - 24.6)		(1.7 - 39.4)
I^2	97.2%		96.5%
P_H	< 0.001		< 0.001
The number of studies	14		10

P-value of meta regression		0.263	
Chest pain			
Prevalence %	4.3		8.1
95% CI %	[2.7 - 6.9]		[5.8 - 11.0]
Range %	(0.0 - 20.5)		(1.1 - 17.9)
I^2	95.6%		82.6%
P_H	< 0.001		< 0.001
The number of studies	10		11
P-value of meta regression		0.097	
Dizziness			
Prevalence %	3.7		21.7
95% CI %	[2.7 - 5.1]		[10.7 - 39.1]
Range %	(0.6 - 6.5)		(6.8 - 35.8)
I^2	86.2%		92.1%
P_H	< 0.001		< 0.001
The number of studies	8		3
P-value of meta regression		0.002	
Palpitation			
Prevalence %	4.9		10.3
95% CI %	[3.2 - 7.3]		[5.2 - 19.2]
Range %	(0.5 - 21.6)		(4.2 - 23.2)
I^2	95.6%		89.9%
P_H	< 0.001		< 0.001
The number of studies	9		5
P-value of meta regression		0.095	
Ageusia			
Prevalence %	5.8		8.3
95% CI %	[2.3 - 14.1]		[5.8 - 11.7]
Range %	(0.0 - 49.9)		(1.0 - 12.7)
I^2	98.9%		76.7%
P_H	< 0.001		< 0.001
The number of studies	11		9
P-value of meta regression		0.446	
Anosmia			
Prevalence %	5.3		8.4
95% CI %	[2.3 - 11.8]		[5.9 - 11.8]
Range %	(0.4 - 53.8)		(1.9 - 15.1)
I^2	99.1%		81.7%
P_H	< 0.001		< 0.001
The number of studies	12		10
P-value of meta regression		0.300	
Sore throat			
Prevalence %	2.5		5.8
95% CI %	[1.3 - 4.6]		[2.3 - 13.9]
Range %	(0.0 - 10.3)		(1.0 - 12.7)
I^2	95.7%		86.9%
P_H	< 0.001		< 0.001
The number of studies	9		3
P-value of meta regression		0.274	

Diarrhea			
Prevalence %	1.9		3.1
95% CI %	[0.7 - 5.0]		[1.0 - 9.8]
Range %	(0.0 - 10.3)		(0.6 - 17.7)
I^2	96.7%		93.1%
P_H	< 0.001		< 0.001
The number of studies	6		6
P-value of meta regression		0.450	
Fever			
Prevalence %	0.7		9.4
95% CI %	[0.2 - 2.0]		[5.2 - 16.3]
Range %	(0.0 - 7.0)		(0.0 - 11.0)
I^2	91.7%		60.7%
P_H	< 0.001		0.078
The number of studies	9		3
P-value of meta regression		< 0.001	

Appendix 7 Figure for pooled prevalence of symptoms in ICU admission < 20% and ≥ 20%

