Table 2. Maternal presenting symptoms and outcomes (of women that tested positive for).

Author	Gestational age at presentation	Gestational age at delivery	Presenting symptoms	Complications	Shah's Classification System for Maternal-Fetal-Neonatal SARS-CoV-2 Intrauterine Infections[11]
Abasse et al.[12]	33 weeks	33 weeks	• Cough • Dyspnoea	• Preterm labour	Unclassified – Possible (1)*
Algarroba et al.[13]	28+4 weeks	28+4 weeks	• Cough • Dyspnoea • Hypoxia	• Sepsis pneumonia • Metabolic acidosis • Preterm labour	Possible (1)
Alzamora et al.[14]	33 weeks	33 weeks	• General malaise • Fatigue • Low-grade fever • Dyspnoea	• Metabolic acidosis • Preterm labour	Unclassified – Possible (1)*
Blauvelt et al.[15]	28 weeks	28 weeks	• Malaise • Fever • Myalgias • Cough • Dyspnoea	• ICU admission requiring mechanical ventilation • Preterm labour	Unlikely (1)
Cao et al.[16]	33+6 to 40+5 weeks	3 late preterm	• Fever (2/10) • Cough (1/10) • Chest tightness (1/10)	• Preterm labour (3/10)	Unlikely (11)
Chen et al.[17]	36 weeks to 39+4 weeks	4 late preterm (all beyond 36 weeks)	• Fever (7/9) • Cough (4/9) • Myalgia (3/9) • Sore throat (2/9) • Malaise (2/9) • Dyspnoea (1/9)	• Preterm labour (4/9)	Not infected (9)
Demirjian et al.[18]	38+5 weeks	39+5 weeks	• Fever • Dyspnoea	• ICU admission requiring mechanical ventilation	Unlikely (1)
Dong et al.[19]	34+2 weeks	38 weeks	• Fever • Nasal congestion • Dyspnoea	• Preterm labour	Possible (1)
Fan et al.[20]	36 weeks and 37 weeks	37 weeks and 36 weeks	• Fever (1/2) • Chills (1/2) • Nasal congestion (2/2) • Sore throat (1/2)	• Preterm labour (½)	Not infected (2)
Fenizia et al.[21]	33+6 weeks to 38 weeks	Not reported	• Not reported	• ICU admission required for mechanical ventilation (1/31) • Interstitial pneumonia (14/31) • Preterm delivery (1/31) • C-section due to severe COVID-19 illness (3/31) • Induction of delivery due to COVID-19 (6/31)	Confirmed (1) Probable (1) Not infected (29)
Gidlöf et al.[22]	36+2 weeks	36+2 weeks	• Headache • Hoarseness • Malaise	• Preeclampsia • Preterm labour	Unlikely (2)
Govind et al.[23]	39 weeks+	39 weeks	• Anosmia (7/9) • Cough (8/9) • Myalgia (5/9) • Fever (4/9)		Possible (1) Unlikely (8)

(continued)

Table 2. Continued.

Author	Gestational age at presentation	Gestational age at delivery	Presenting symptoms	Complications	Shah's Classification System for Maternal-Fetal-Neonatal SARS-CoV-2 Intrauterine Infections[11]
Hu et al.[24]	Not reported	Not reported	• Sore throat (4/9) • Lethargy (6/9) • Dyspnoea (4/9) • Low grade fever (4/7) • Cough (2/7) • Diarrhoea (1/7)		Unclassified – Possible (1)* Unlikely (6)
Kalafat et al.[25]	35+3 weeks	36 weeks	• Cough • Dyspnoea	• ICU admission required for mechanical ventilation • Preterm labour	Not infected (1)
Khan et al.[26]	34+6 to 38+2 weeks	34+6 weeks to 39 +1 weeks	• Fever (2/3) • Cough (3/3) • Chest tightness (1/3)	• Preterm labour	Not infected (3)
Khan et al.[27]	35 weeks to 41 weeks	35+5 to 41 (average=38.1 weeks)	• Fever (3/17) • Cough (6/17) • Diarrhoea (3/17) • Dyspnoea (2/17) • Nasal congestion (2/17) • Sputum (1/17)	• Preterm labour (5/17)	Unlikely (17)
Kirtsman et al.[28]	35+5 weeks	35+5 weeks	• Myalgia • Decreased appetite • Cough • Fever	• Preterm labour	Probable (1)
Knight et al.[29]	Average=34 weeks	Median=38 weeks	• Fever (275/427) • Cough (230/427) • Dyspnoea (155/427) • Lethargy (60/427) • Headache (55/427) • joint pain (50/427) • sore throat (45/427) • Diarrhoea (20/427)	• ICU admission required for mechanical ventilation (10/427) • Interstitial pneumonia (24/427) • Maternal death (1)	Unclassified – Probable (6)* Possible (6) Unlikely (253)
Lang et al.[30]	35+2 weeks	35+2 weeks	• Cough	• Preterm labour	Not infected (1)
Li et al.[31]	35 weeks	35 + 4 weeks	• Cough	• Preterm labour	Not infected (1)
Liao et al.[32]	31+1 weeks to 42 weeks	31+1 weeks to 42 weeks	• Fever (5/10) • Cough (3/10) • Sore throat (1/17) • Chest pain (1/10)	• Preterm labour	Unlikely (10)
Liu et al.[33]	35+2 weeks to 41+2 weeks	38.6±1.5 weeks	• Fever (11/19) • Cough (5/19) • Dyspnoea (5/19) • Diarrhoea (2/19)	• Preterm labour (2/19)	Unlikely (19)
Lowe et al.[34]	40 weeks	40+2 weeks	• Asymptomatic		Unlikely (1)
Lu et al.[35]	38 weeks	38 +2 weeks	• Asymptomatic		Unlikely (1)
Marzollo et al.[36]	38 weeks	38 weeks	• Fever	• Severe thrombocytopenia	Unclassified – Possible (1)*
Nie et al.[37]	28+weeks to 40 weeks	Not reported	• Fever (21/33) • Cough (13/33) • Fatigue (7/33) • Dyspnoea (7/33)	• Preterm labour (10/33)	Unlikely (1) Not infected (25)
Oncel et al.[38]	Not reported	26–38 weeks	• Not reported	• ICU admission requiring mechanical ventilation (8/125)	Unlikely (125)

(continued)

Table 2. Continued.

Author	Gestational age at presentation	Gestational age at delivery	Presenting symptoms	Complications	Shah's Classification System for Maternal-Fetal-Neonatal SARS-CoV-2 Intrauterine Infections[11]
				• Preterm labour (33/125) • Maternal death (6/125)	
Patanè et al.[39]	35+1 to 37+6 weeks	35+1 to 37+6 weeks	• Fever (2) • Cough (2)	• Preterm labour (½)	Probable (2) Unlikely (20)
Patil et al.[40]	30+5 weeks to 41 weeks	30+5 weeks to 41 weeks	• Fever (15/45) • Cough (16/45) • Sore throat (4/45) • Shortness of breath (6/45)	• ICU admission requiring mechanical ventilation (2/45) • Preterm labour (4/45)	Possible (2) Unlikely (43)
Penfield et al.[41]	26+5 weeks to 41+3 weeks	26+5 weeks to 41+3 weeks	• Not reported	• Preterm labour (3/9)	Possible (3) Not infected (8)
Peng et al.[42]	35+3 weeks	36 weeks	• Fever • Fatigue • Dyspnoea	Preterm labour	Not infected (1)
Pereira et al.[43]	Median=32 weeks	Median=32 weeks	• Asymptomatic (15/60) • Dyspnoea (17/60) • Fever (34/60) • Cough (34/60)	• ICU admission required for mechanical ventilation (1/60) • Preterm labour (2/60)	Unlikely (17) Not infected (6)
Pierce-Williams et al.[44]	Mean=30 + 6 weeks	Mean=30+6 weeks	• Not reported	• Maternal cardiac arrest (1/17) • Preterm labour (15/17)	Unlikely (33)
Sisman et al.[45]	34 weeks	34 weeks	• Fever • Diarrhea • Back pain	• Preterm labour	Confirmed (1)
Vivanti et al.[46]	35+2 weeks	35+5 weeks	• Fever • Cough	• Preterm labour	Confirmed (1)
Wang et al.[47]	40 weeks	40 weeks	• Fever • Abdominal pain • Vaginal bleeding		Unlikely (1)
Wu et al.[48]	35 weeks to 40 weeks	35 weeks to 40 weeks	• Fever (8/29) • Cough (9/29) • Diarrhoea (2/29) • Dyspnoea (3/29) • Asymptomatic (15/29)	• Preterm labour (3/29)	Unclassified – Possible (4)* Unlikely (1) Not infected (25)
Xiong et al.[49]	33+1 weeks	38+4 weeks	• Fever • Cough • Chills		Not infected (1)
Yan et al.[50]	Median = 38 weeks	Median = 38 weeks	• Fever (59/116) • Cough (33/116) • Asymptomatic (27/116)	• ICU admission requiring mechanical ventilation (2/116) • Preterm labour (6/116)	Unlikely (76) Not infected (10)
Yang et al.[51]	36 weeks to 37 weeks	36 weeks to 37 weeks	• Fever (6/7) • Cough (6/7) • Abdominal pain (6/7) • Diarrhoea (6/7)	• Preeclampsia (2/7) • Preterm labour (4/7)	Unlikely (7)
Yang et al.[52]	30 weeks to 40 weeks	30 weeks to 40 weeks	• Fever (13/27) • Cough (9/27) • Vomiting (1/27)	• Preterm labour (1/27)	Unlikely (24)

(continued)

Table 2. Continued.

Author	Gestational age at presentation	Gestational age at delivery	Presenting symptoms	Complications	Shah's Classification System for Maternal-Fetal-Neonatal SARS-CoV-2 Intrauterine Infections[11]
Yin et al.[53]	Mean=35 weeks	Mean=35 weeks	• Fever (17/31) • Cough (15/31) • Fatigue (6/31) • Dyspnoea (8/31) • Diarrhoea (2/31) • Myalgia (3/ 31)	• Preterm labour (5/31)	Not infected (31)
Yu et al.[54]	37 weeks to 41+2 weeks	Mean=39+2 days	• Fever (6/7) • Cough (1/7) • Dyspnoea (1/7) • Diarrhoea (6/7)		Unlikely (3)
Zamaniyan et al.[55]	32 weeks	33 weeks	• Dyspnoea • Myalgia • Anorexia • Nausea • Cough • Fever	• ICU admission requiring mechanical ventilation • Premature labour • Maternal death	Probable (1)
Zeng et al.[56]	31 weeks to 40 weeks	31 weeks to 40 weeks	• Fever (8/33) • Dyspnoea (4/33) • Cough (10/33)	• Preterm labour (3/33)	Unclassified – Possible (3)* Unlikely (30)
Zeng et al.[57]	>30 weeks	Not reported	Not reported		Possible (3) Unlikely (3)
Zhu et al.[58]	>34 weeks	>34 weeks	• Fever (8/9) • Cough (4/9) • Cholecystitis (1/9) • Diarrhoea (1/9)	• Preterm labour (6/9)	Unlikely (10)

Table 3. Infected neonatal symptoms and outcomes.

Author	Shah's Classification System for Maternal-Fetal-Neonatal SARS-CoV-2 Intrauterine Infections[11]	Details of mother of affected neonate	Details of affected neonate	Neonatal complications	1 min Apgar score	5 min Apgar score
Abasse et al.[12]	Unclassified – Possible (1)*			• Fever • Respiratory difficulties • NICU admission for monitoring	5	7
Alzamora et al.[14]	Unclassified – Possible(1)*	Mother possible 1: admitted at 33 weeks due to respiratory insufficiency. Was admitted to ICU for mechanical ventilation. Caesarean delivery due to decreased respiratory compromise.	Neonate possible: 33 weeks gestation, delivered by Caesarean section, required resuscitation and admitted to NICU for mechanical ventilation	• Cough • Respiratory difficulties • Lethargy • NICU admission: mechanical ventilation	6	8
Demirjian et al.[18]	Unlikely (1)			• Fever • Respiratory difficulties	5	9
Fenizia et al.[21]	Confirmed (1), Possible (1)	Mother confirmed 1: admitted at 33+6 with fever and dyspnoea, delivered spontaneously Mother confirmed 2: No reported symptoms, delivered spontaneously	Neonate confirmed 1: 34+4 weeks gestation, delivered spontaneous vaginal, asymptomatic, UA pH 7.12. Neonate possible 1: 39+2 weeks gestation, delivered spontaneous vaginal, asymptomatic, UA pH 7.14	• NICU admission for observation (2)	9 (2)	10 (2)
Govind et al.[23]	Possible (1)	Mother possible 1: admitted at 38 weeks with maternal pneumonia and respiratory distress.	Neonate possible 1: 38 weeks gestation, delivered by Caesarean section, required resuscitation and ECMO. Admitted to NICU for mechanical ventilation.	• Fever • Cough • Lethargy • Foetal distress • Respiratory difficulties • NICU admission: mechanical ventilation	5	9
Hu et al.[24]	Unclassified – Possible (1)*			• Asymptomatic	8	9
Kirtsman et al.[28]	Probable (1)			• Vomiting/feeding difficulties • Intermittent hypoglycaemic episodes • NICU admission for observation	9	9
Knight et al.[29]	Unclassified – Probable (6)*, Possible (6)	• Maternal symptoms not reported.	• Six neonates tested positive within the first 12 h after birth. • Two of the six infants with early onset SARS-CoV-2 infection were from unassisted vaginal births; four were born by caesarean, three of which were pre-labour. • The six infants who	NICU admission (6/12)	NR	NR

(continued)

Table 3. Continued.

Author	Shah's Classification System for Maternal-Fetal-Neonatal SARS-CoV-2 Intrauterine Infections[11]	Details of mother of affected neonate	Details of affected neonate	Neonatal complications	1 min Apgar score	5 min Apgar score
			developed later infection were born by pre-labour caesarean (*n*=4) and vaginal birth (*n*=2). Only one of the infants with an early positive test for SARS-CoV-2 RNA was admitted to a neonatal unit for observation, compared with five infants with a later positive test. • Symptoms were not reported			
Marzollo et al.[36]	Unclassified – Possible (1)*			• Respiratory difficulties • Vomiting/feeding difficulties • NICU admission: mechanical ventilation	9	10
Nie et al.[37]	Unlikely (1)			Asymptomatic	9	9
Oncel et al.[38]	Unlikely (4)		• Prematurity 2/4	• Fever (1/4) • Cough (1/4) • Respiratory difficulties (3/4) • Vomiting/feeding difficulties (1/4) • NICU admission: mechanical ventilation (1/4)	2 7 8 (2)	6 8 9 (2)
Patanè et al.[39]	Probable (2)			• Vomiting/feeding difficulties (2/2)	9 (2)	10 (2)
Patil et al.[40]	Possible (2)			Asymptomatic (2)	9	9
Sisman et al.[45]	Confirmed (1)			• Fever • Respiratory difficulties	7	9
Vivanti et al.[46]	Confirmed (1)			• Foetal distress • Neurological symptoms (hypertonia/ irritability) • Resuscitation required • NICU admission: mechanical ventilation	4	1
Wang et al.[47]	Unlikely (1)			• Vomiting/feeding difficulties	8	9
Wu et al.[48]	Unclassified – Possible (2)*, Unlikely (3)	Mother of neonate possible 1: presented with cough, had negative throat swab, clinically suspected with COVID-19. Mother of	Neonate Possible 1 (5): 40+4 weeks gestation, delivered by caesarean delivery, asymptomatic, admitted to NICU for observation Neonate Possible 2 (9): 39	• Fever (1/5) • NICU admission for monitoring (2/5)	9 (2) 10 (3)	10 (5)

(continued)

Table 3. Continued.

Author	Shah's Classification System for Maternal-Fetal-Neonatal SARS-CoV-2 Intrauterine Infections[11]	Details of mother of affected neonate	Details of affected neonate	Neonatal complications	1 min Apgar score	5 min Apgar score
		neonate possible 2: asymptomatic, noted to have CT findings of bilateral focal GGO, negative throat swab, clinically suspected with COVID-19 Mother of neonate unlikely 1: asymptomatic, CT findings positive for left lung infectious lesions, bilateral pleural thickening and pleural effusion. Throat swab positive for Mother of Neonate Unlikely 2: Presented with fever, cough, vomiting with positive throat swab for Mother of Neonate Unlikely 3: Presented with fever, stuffy nose, shortness of breath, CT positive for bilateral scattered GGO, throat swab positive for	+1 weeks gestation, delivered by caesarean delivery, developed fever and admitted to NICU for observation Neonate Unlikely 1: 38+3 weeks gestation, delivered by cesarean delivery, asymptomatic, throat swab negative for Neonate Unlikely 2: 37+1 weeks gestation, delivered by caesarean delivery, asymptomatic, throat swab negative for Neonate Unlikely 3: 37+6 weeks gestation, delivered by caesarean delivery, asymptomatic, throat swab negative for			
Yu et al.[54]	Unlikely (3)		One neonate was found to have a mild pulmonary infection. The shortness of breath relieved quickly under neonatal care and was admitted to the NICU for monitoring	• Respiratory difficulties (1/3) • NICU admission for monitoring (1/3)	9	10
Zamaniyan et al.[55]	Probable (1)			• Fever	8	9
Zeng et al.[56]	Unclassified – Possible (3)*		Neonate 1: 40 weeks gestation by caesarean delivery, developed lethargy and fever and required NICU admission. Neonate 2: 40+4 weeks gestation by caesarean delivery presenting with lethargy, vomiting, and fever. Neonate 3: 31+2 weeks gestation by caesarean delivery. Resuscitation was required. Apgar scores were 3, 4, and 5 at 1, 5, and 10 minutes after birth. Required NICU admission for mechanical ventilation with neonatal respiratory distress syndrome and pneumonia.	• Fever (2/3) • Lethargy (2/3) • Foetal distress (1/3) • Respiratory difficulties (1/3) • Vomiting/feeding difficulties (1/3) • NICU admission for monitoring (2): mechanical ventilation (1)	3 (1), NR (2)	4 (1), NR (2)