

# Case PnYmzBrDxbjITyQ14316 — Questions

## Case Details

**Demographics** 62-year-old Hispanic female; teaching assistant

**Chief complaint** blurred vision

**History of present illness**

**Secondary complaints/symptoms** none

**Patient ocular history** last eye exam 1 month ago at another office, wears single vision reading glasses only, new glasses are 2 weeks old; history of refractive surgery (unsure of procedure)

**Family ocular history** father: macular degeneration

**Patient medical history** rheumatoid arthritis

**Medications taken by patient** Plaquenil® 200 mg b.i.d.

**Patient allergy history** Demerol®

**Family medical history** mother: hyperthyroidism, father: tuberculosis

**Review of systems**

**Mental status**

**Clinical findings**

**Uncorrected visual acuity**

**New spectacle Rx**

**Pupils:** PERRL, negative APD

**EOMs:** full, no restrictions OU

**Cover test:** distance: orthophoria, near: 4 exophoria

**Confrontation fields:** full to finger counting OD, OS

**Subjective refraction**

**Slit lamp**

**IOPs:** OD: 16 mmHg, OS: 18 mmHg @ 10:05 am by Goldmann applanation tonometry

**Fundus OD**


**Fundus OS** deep and quiet OD, OS

**Blood pressure:** 112/82 mmHg, right arm, sitting

**Pulse:** 74 bpm, regular

- Character/signs/symptoms: blurry vision at near with new glasses
- Location: OD, OS
- Severity: moderate
- Nature of onset: noticed after picking up her new glasses
- Duration: 2 weeks
- Frequency: constant
- Exacerbations/remissions: vision improves if she brings reading material closer
- Relationship to activity or function: near vision tasks only
- Accompanying signs/symptoms: eye strain and fatigue
- Constitutional/general health: denies
- Ear/nose/throat: denies
- Cardiovascular: denies
- Pulmonary: denies
- Dermatological: denies
- Gastrointestinal: denies
- Genitourinary: denies
- Musculoskeletal: joint pain and stiffness
- Neuropsychiatric: denies
- Endocrine: denies
- Hematologic: denies
- Immunologic: denies
- Orientation: oriented to time, place, and person
- Mood: appropriate
- Affect: appropriate
- OD: VA distance: 20/20
- OS: VA distance: 20/30
- OD: +3.50 -0.25 x 133; VA near: 20/20 @ 33 cm
- OS: +4.25 -0.50 x 062; VA near: 20/20 @ 33 cm
- OD: +0.25 DS add: +2.50; VA distance: 20/20, VA near: 20/20 @ 40 cm
- OS: +1.00 -0.50 x 060 add: +2.50; VA distance: 20/20, VA near: 20/20 @ 40 cm
- lids/lashes/adnexa: dermatochalasis OD, OS
- conjunctiva: normal OD, OS
- cornea: OD similar to OS, OS 8 small circular scars noted circumferentially in peripheral cornea (see images 1 & 2)

- anterior chamber: deep and quiet OD, OS
- iris: normal OD, OS
- lens: trace nuclear sclerosis OD, OS
- vitreous: posterior vitreous detachment OD, OS
- C/D: 0.40 H/0.40 V
- macula: normal
- posterior pole: normal
- periphery: see images 3 & 4
- C/D: 0.40 H/0.40 V
- macula: normal
- posterior pole: normal
- periphery: similar to images 3 & 4



### Question 1 / 5

Given images 1 and 2, what is the MOST likely type of refractive surgery undergone by this patient?

- A) Photorefractive keratectomy (PRK)
- B) Implantable collamer lens (ICL)
- C) Laser-assisted in situ keratomileusis (LASIK)
- D) Conductive keratoplasty (CK)
- E) Refractive lens exchange (RLE)

### Question 2 / 5

Which of the following refractive errors is best suited for that specific type of surgery?

- A) +2.25 DS
- B) -2.75 -1.25 x 090
- C) -0.75 -0.75 x 180
- D) -1.25 DS
- E) +1.00 -1.00 x 180
- F) +4.25 -0.75 x 180

### Question 3 / 5

How would you expect a patient's corneal topography to change after undergoing this type of refractive surgery?

- A) Central and mid-peripheral flattening
- B) Central flattening and mid-peripheral steepening
- C) The corneal topography is not expected to change
- D) Central and mid-peripheral steepening
- E) Central steepening and mid-peripheral flattening

### Question 4 / 5

What is the MOST likely cause of the patient's entering visual concern?

- A) Undercorrection; causing an increased focal length and further working distance
- B) Overcorrection; causing an increased focal length and further working distance
- C) Undercorrection; causing a reduced focal length and closer working distance
- D) Overcorrection; causing a reduced focal length and closer working distance

### Question 5 / 5

Which of the following represents the MOST appropriate diagnosis of the patient's retinal findings observed in images 3 and 4?

- A) Reticular pigmentary degeneration
- B) Retinitis pigmentosa
- C) Plaquenil® retinopathy

## D) Lattice degeneration