# Straight-Line Planar Graph Drawing — Part 1


Lecture Graph Drawing Algorithms · 192.053

Martin Nöllenburg 17.04.2018











Planar graphs are an important graph class in graph drawing and graph theory.

What do you know about planar graphs?


- average degree  $\leq 6$ - Enlar characteristic: |V| - |E| + |F| = 2  $|E| \leq 3|V| - 6$ - no  $K_{3,3}$  or  $K_5$  as a minor

- 4 colorable



Planar graphs are an important graph class in graph drawing and graph theory.


**Def:** A planar graph G is a simple graph that can be drawn/ embedded in the plane  $\mathbb{R}^2$  without edge crossings.






Planar graphs are an important graph class in graph drawing and graph theory.

**Def:** A planar graph G is a simple graph that can be drawn/ embedded in the plane  $\mathbb{R}^2$  without edge crossings.



different drawings same embedding?






Planar graphs are an important graph class in graph drawing and graph theory.

Plane graph = planar graph + embedding


**Def:** A planar graph G is a simple graph that can be drawn/ embedded in the plane  $\mathbb{R}^2$  without edge crossings.




**Def:** The **rotation scheme** of a planar drawing is the circular ordering of the edges incident to each vertex.

Two planar drawings have the same embedding if they have the same rotation scheme and the same external face.

different drawings



same embedding?





Planar graphs are an important graph class in graph drawing and graph theory.

**Def:** A planar graph G is a simple graph that can be drawn/ embedded in the plane  $\mathbb{R}^2$  without edge crossings.

**Theorem:** G is planar  $\Leftrightarrow G$  has no  $K_5$  or  $K_{3,3}$  minor.

[Kuratowski 1930]



Planar graphs are an important graph class in graph drawing and graph theory.

**Def:** A planar graph G is a simple graph that can be drawn/ embedded in the plane  $\mathbb{R}^2$  without edge crossings.

**Theorem:** G is planar  $\Leftrightarrow G$  has no  $K_5$  or  $K_{3,3}$  minor.

[Kuratowski 1930]

**Theorem:** Testing whether a simple graph with n vertices is planar can be done in O(n) time. [Hopcroft, Tarjan 1974]



Planar graphs are an important graph class in graph drawing and graph theory.

**Def:** A planar graph G is a simple graph that can be drawn/ embedded in the plane  $\mathbb{R}^2$  without edge crossings.

**Theorem:** G is planar  $\Leftrightarrow G$  has no  $K_5$  or  $K_{3,3}$  minor.

[Kuratowski 1930]

**Theorem:** Testing whether a simple graph with n vertices is planar can be done in O(n) time. [Hopcroft, Tarjan 1974]

**Theorem:** Every planar graph has a planar straight-line drawing. [Wagner 1936, Fáry 1948, Stein 1951]



Planar graphs are an important graph class in graph drawing and graph theory.

**Def:** A planar graph G is a simple graph that can be drawn/ embedded in the plane  $\mathbb{R}^2$  without edge crossings.

**Theorem:** G is planar  $\Leftrightarrow G$  has no  $K_5$  or  $K_{3,3}$  minor.

[Kuratowski 1930]

**Theorem:** Testing whether a simple graph with n vertices is


planar can be done in O(n) time. [Hopcroft, Tarjan 1974]

**Theorem:** Every planar graph has a planar straight-line drawing.

[Wagner 1936, Fáry 1948, Stein 1951]

**Theorem:** G is planar  $\Leftrightarrow$  G has a disk contact representation.

[Koebe 1936]





Planar graphs are an important graph class in graph drawing and graph theory.

**Def:** A planar graph G is a simple graph that can be drawn/ embedded in the plane  $\mathbb{R}^2$  without edge crossings.

**Theorem:** G is planar  $\Leftrightarrow G$  has no  $K_5$  or  $K_{3,3}$  minor.

[Kuratowski 1930]

**Theorem:** Testing whether a simple graph with n vertices is planar can be done in O(n) time. [Hopcroft, Tarjan 1974]

**Theorem:** Every planar graph has a planar straight-line drawing. [Wagner 1936, Fáry 1948, Stein 1951]

**Theorem:** G is planar  $\Leftrightarrow G$  has a disk contact representation. [Koebe 1936]

**Theorem:** G=(V,E) is planar  $\Rightarrow |V|-|E|+|F|=2$  [Euler 1707-1783]

#### Drawing Planar Graphs



How do we find good straight-line planar drawings?

#### Drawing Planar Graphs



How do we find good straight-line planar drawings?

**Theorem:** A straight-line convex drawing of a 3-connected planar graph can be computed in linear time. [Tutte 1963]

But: no polynomial bound on the required area

#### Drawing Planar Graphs




How do we find good straight-line planar drawings?

**Theorem:** A straight-line convex drawing of a 3-connected planar graph can be computed in linear time. [Tutte 1963]


But: no polynomial bound on the required area

**Theorem:** Every n-vertex embedded planar graph has a planar straight-line drawing on an  $(2n-4)\times(n-2)$  grid. [de Fraysseix, Pach, Pollack 1988]





























sufficient to focus on drawing maximally planar (triangulated) graphs






sufficient to focus on drawing maximally planar (triangulated) graphs









sufficient to focus on drawing maximally planar (triangulated) graphs









sufficient to focus on drawing maximally planar (triangulated) graphs








sufficient to focus on drawing maximally planar (triangulated) graphs





#### Overview



#### Canonical ordering

Shift algorithm

**Implementation** 

#### Canonical Ordering



**Def:** Let G = (V, E) be a **triangulated planar embedded graph** with  $n \geq 3$  vertices. An ordering  $\pi = (v_1, v_2, \dots, v_n)$  is called a **canonical ordering**, if the following conditions hold for each k,  $3 \leq k \leq n$ 

 $\blacksquare$   $\{v_1, \ldots v_k\}$  induce 2-connected internally triangulated graph  $G_k$ ,

lacksquare edge  $(v_1,v_2)$  belongs to the outer face of each  $G_k$ ,


for all k < n vertex  $v_{k+1}$  lies in the external face of  $G_k$ , and all neighbors of  $v_{k+1}$  in  $G_k$  appear on the boundary  $C_k$  of  $G_k$  consecutively.

#### Canonical Ordering




**Def:** Let G=(V,E) be a **triangulated planar embedded graph** with  $n\geq 3$  vertices. An ordering  $\pi=(v_1,v_2,\ldots,v_n)$  is called a **canonical ordering**, if the following conditions hold for each k,  $3\leq k\leq n$ 


- $\blacksquare$   $\{v_1, \ldots v_k\}$  induce 2-connected internally triangulated graph  $G_k$ ,
- lacksquare edge  $(v_1,v_2)$  belongs to the outer face of each  $G_k$ ,
- for all k < n vertex  $v_{k+1}$  lies in the external face of  $G_k$ , and all neighbors of  $v_{k+1}$  in  $G_k$  appear on the boundary  $C_k$  of  $G_k$  consecutively.





















































# Example: Canonical Ordering






# Example: Canonical Ordering





# Example: Canonical Ordering







**Lemma.** Every triangulated planar graph G=(V,E) has a canonical ordering.



**Lemma.** Every triangulated planar graph G=(V,E) has a canonical ordering.

**Proof:** inverse induction on k



**Lemma.** Every triangulated planar graph G=(V,E) has a canonical ordering.

**Proof:** inverse induction on k

 $\blacksquare k = n$ :




Induction hypothesis: vertices  $v_n, v_{n-1}, \ldots, v_{k+1}$  were chosen such that they satisfy the three conditions.




**Lemma.** Every triangulated planar graph G=(V,E) has a canonical ordering.

**Proof:** inverse induction on k

 $\blacksquare k = n$ :



- Induction hypothesis: vertices  $v_n, v_{n-1}, \ldots, v_{k+1}$  were chosen such that they satisfy the three conditions.
- lacksquare consider  $G_k$  and find suitable vertex  $v_k$  on  $C_k$




 $\blacksquare$   $G_k$  is 2-connected and internally triangulated




**Lemma.** Every triangulated planar graph G = (V, E) has a canonical ordering.

**Proof:** inverse induction on k

k=n:



- induction hypothesis: vertices  $v_n, v_{n-1}, \ldots, v_{k+1}$  were chosen such that they satisfy the three conditions.
- consider  $G_k$  and find suitable vertex  $v_k$  on  $C_k$




- lacksquare  $G_k$  is 2-connected and internally triangulated
- lacksquare  $v_k$  should not be incident to chord of  $C_k$




**Lemma.** Every triangulated planar graph G = (V, E) has a canonical ordering.

**Proof:** inverse induction on k

k=n:



- induction hypothesis: vertices  $v_n, v_{n-1}, \ldots, v_{k+1}$  were chosen such that they satisfy the three conditions.
- consider  $G_k$  and find suitable vertex  $v_k$  on  $C_k$




- lacksquare  $G_k$  is 2-connected and internally triangulated
- lacksquare  $v_k$  should not be incident to chord of  $C_k$


Is this sufficient?













































We show:  $v_k$  not incident to a chord  $\Rightarrow G_{k-1}$  is 2-connected.




Can we always find a vertex not incident to a chord?






**We show:**  $v_k$  not incident to a chord  $\Rightarrow G_{k-1}$  is 2-connected.



Can we always find a vertex not incident to a chord?



- chords form proper nesting
- innermost chords span at least one such vertex

### Computing a Canonical Ordering



CanonicalOrdering(plane graph G = (V, E))

#### forall $v \in V$ do

 $\lfloor \mathsf{chords}(v) \leftarrow 0$ ;  $\mathsf{out}(v) \leftarrow \mathsf{false}$ ;  $\mathsf{mark}(v) \leftarrow \mathsf{false}$ ;  $\mathsf{out}(v_1)$ ,  $\mathsf{out}(v_2)$ ,  $\mathsf{out}(v_n) \leftarrow \mathsf{true}$ 

for k = n to 3 do

choose  $v \neq v_1, v_2$  with mark(v) = false, out(v) = true, chords(v) = 0


 $v_k \leftarrow v$ ; mark $(v) \leftarrow$  true

$$(w_1 = v_1, w_2, \dots, w_{t-1}, w_t = v_2) \leftarrow C_{k-1}$$

 $(w_p, \dots, w_q) \leftarrow \text{unmarked neighbors of } v_k$ 

 $\operatorname{out}(w_i) \leftarrow \operatorname{true} \text{ for all } p < i < q$ 

update chords( $\cdot$ ) for these  $w_i$  and their neighbors



- $\blacksquare$  mark(v) = true iff vertex v was numbered
- lacksquare out(v)= true iff v is currently in the external face

### Computing a Canonical Ordering



```
lacksquare chord(v) = number of chords incident to v
```

update chords( $\cdot$ ) for these  $w_i$  and their neighbors

 $\blacksquare$  mark(v) = true iff vertex v was numbered

 $(w_1 = v_1, w_2, \dots, w_{t-1}, w_t = v_2) \leftarrow C_{k-1}$ 

 $(w_p, \dots, w_q) \leftarrow \text{unmarked neighbors of } v_k$ 

 $\mathsf{out}(w_i) \leftarrow \mathsf{true} \ \mathsf{for} \ \mathsf{all} \ p < i < q$ 

lacksquare out(v) = true iff v is currently in the external face

**Lemma:** Algorithm CanonicalOrdering computes a canonical ordering of G in O(n) time.

Overview



Canonical ordering


#### Shift algorithm

**Implementation** 

## Shift Algorithm [de Fraysseix, Pach, Pollack 1988]




#### even $L_1$ -distance




[de Fraysseix, Pach, Pollack 1988]



even  $L_1$ -distance




- $lacksquare v_1$  is on (0,0),  $v_2$  is on (2k-6,0)
- boundary  $C_{k-1}$  of  $G_{k-1}$  (minus edge  $(v_1, v_2)$ ) is drawn x-monotone
- each edge of  $C_{k-1}$  (minus edge  $(v_1, v_2)$ ) is drawn with slopes  $\pm 1$




[de Fraysseix, Pach, Pollack 1988]



even  $L_1$ -distance




- $lacksquare v_1$  is on (0,0),  $v_2$  is on (2k-6,0)
- boundary  $C_{k-1}$  of  $G_{k-1}$  (minus edge  $(v_1, v_2)$ ) is drawn x-monotone
- each edge of  $C_{k-1}$  (minus edge  $(v_1, v_2)$ ) is drawn with slopes  $\pm 1$




[de Fraysseix, Pach, Pollack 1988]



even  $L_1$ -distance




- $lacksquare v_1$  is on (0,0),  $v_2$  is on (2k-6,0)
- boundary  $C_{k-1}$  of  $G_{k-1}$  (minus edge  $(v_1, v_2)$ ) is drawn x-monotone
- each edge of  $C_{k-1}$  (minus edge  $(v_1, v_2)$ ) is drawn with slopes  $\pm 1$




[de Fraysseix, Pach, Pollack 1988]



even  $L_1$ -distance




- $lacksquare v_1$  is on (0,0),  $v_2$  is on (2k-6,0)
- boundary  $C_{k-1}$  of  $G_{k-1}$  (minus edge  $(v_1, v_2)$ ) is drawn x-monotone
- each edge of  $C_{k-1}$  (minus edge  $(v_1, v_2)$ ) is drawn with slopes  $\pm 1$




[de Fraysseix, Pach, Pollack 1988]



even  $L_1$ -distance




- $lacksquare v_1$  is on (0,0),  $v_2$  is on (2k-6,0)
- boundary  $C_{k-1}$  of  $G_{k-1}$  (minus edge  $(v_1, v_2)$ ) is drawn x-monotone
- each edge of  $C_{k-1}$  (minus edge  $(v_1, v_2)$ ) is drawn with slopes  $\pm 1$




[de Fraysseix, Pach, Pollack 1988]



even  $L_1$ -distance




- $lacksquare v_1$  is on (0,0),  $v_2$  is on (2k-6,0)
- boundary  $C_{k-1}$  of  $G_{k-1}$  (minus edge  $(v_1, v_2)$ ) is drawn x-monotone
- each edge of  $C_{k-1}$  (minus edge  $(v_1, v_2)$ ) is drawn with slopes  $\pm 1$




[de Fraysseix, Pach, Pollack 1988]



even  $L_1$ -distance




- $lacksquare v_1$  is on (0,0),  $v_2$  is on (2k-6,0)
- boundary  $C_{k-1}$  of  $G_{k-1}$  (minus edge  $(v_1, v_2)$ ) is drawn x-monotone
- each edge of  $C_{k-1}$  (minus edge  $(v_1, v_2)$ ) is drawn with slopes  $\pm 1$




[de Fraysseix, Pach, Pollack 1988]



even  $L_1$ -distance




- $lacksquare v_1$  is on (0,0),  $v_2$  is on (2k-6,0)
- boundary  $C_{k-1}$  of  $G_{k-1}$  (minus edge  $(v_1, v_2)$ ) is drawn x-monotone
- each edge of  $C_{k-1}$  (minus edge  $(v_1, v_2)$ ) is drawn with slopes  $\pm 1$




[de Fraysseix, Pach, Pollack 1988]



even  $L_1$ -distance




- $lacksquare v_1$  is on (0,0),  $v_2$  is on (2k-6,0)
- boundary  $C_{k-1}$  of  $G_{k-1}$  (minus edge  $(v_1, v_2)$ ) is drawn x-monotone
- each edge of  $C_{k-1}$  (minus edge  $(v_1, v_2)$ ) is drawn with slopes  $\pm 1$




[de Fraysseix, Pach, Pollack 1988]



even  $L_1$ -distance




- $lacksquare v_1$  is on (0,0),  $v_2$  is on (2k-6,0)
- boundary  $C_{k-1}$  of  $G_{k-1}$  (minus edge  $(v_1, v_2)$ ) is drawn x-monotone
- each edge of  $C_{k-1}$  (minus edge  $(v_1, v_2)$ ) is drawn with slopes  $\pm 1$




[de Fraysseix, Pach, Pollack 1988]



even  $L_1$ -distance




- $lacksquare v_1$  is on (0,0),  $v_2$  is on (2k-6,0)
- boundary  $C_{k-1}$  of  $G_{k-1}$  (minus edge  $(v_1, v_2)$ ) is drawn x-monotone
- each edge of  $C_{k-1}$  (minus edge  $(v_1, v_2)$ ) is drawn with slopes  $\pm 1$




# Shift Algorithm: Example




























































































































- each internal vertex is covered exactly once
- lacksquare covering relation defines a tree in G
- $\blacksquare$  forest in  $G_i$ ,  $3 \le i \le n-1$






- each internal vertex is covered exactly once
- lacksquare covering relation defines a tree in G
- forest in  $G_i$ ,  $3 \le i \le n-1$






- each internal vertex is covered exactly once
- lacksquare covering relation defines a tree in G
- forest in  $G_i$ ,  $3 \le i \le n-1$






- each internal vertex is covered exactly once
- lacksquare covering relation defines a tree in G
- forest in  $G_i$ ,  $3 \le i \le n-1$





- each internal vertex is covered exactly once
- lacktriangle covering relation defines a tree in G
- forest in  $G_i$ ,  $3 \le i \le n-1$





- each internal vertex is covered exactly once
- lacktriangle covering relation defines a tree in G
- forest in  $G_i$ ,  $3 \le i \le n-1$

**Lemma:** Let  $G_k$  be a planar straight-line grid drawing. Let  $0 < \delta_1 \le \delta_2 \le \cdots \le \delta_t \in \mathbb{N}$ . If we shift each  $L(w_i)$  by  $\delta_i$  to the right, we get another planar straight-line grid drawing.



**Lemma:** Let  $G_k$  be a planar straight-line grid drawing. Let  $0 < \delta_1 \le \delta_2 \le \cdots \le \delta_t \in \mathbb{N}$ . If we shift each  $L(w_i)$  by  $\delta_i$  to the right, we get another planar straight-line grid drawing.



**Lemma:** Let  $G_k$  be a planar straight-line grid drawing. Let  $0 < \delta_1 \le \delta_2 \le \cdots \le \delta_t \in \mathbb{N}$ . If we shift each  $L(w_i)$  by  $\delta_i$  to the right, we get another planar straight-line grid drawing.


#### **Proof**

 $\blacksquare$  induction on i, i.e. we consider  $G_3, \ldots, G_n$ 



**Lemma:** Let  $G_k$  be a planar straight-line grid drawing. Let  $0 < \delta_1 \le \delta_2 \le \cdots \le \delta_t \in \mathbb{N}$ . If we shift each  $L(w_i)$  by  $\delta_i$  to the right, we get another planar straight-line grid drawing.

- $\blacksquare$  induction on i, i.e. we consider  $G_3, \ldots, G_n$
- $\blacksquare$  clearly true for  $G_3$ ; assume the lemma holds for  $G_{k-1}$







**Lemma:** Let  $G_k$  be a planar straight-line grid drawing. Let  $0 < \delta_1 \le \delta_2 \le \cdots \le \delta_t \in \mathbb{N}$ . If we shift each  $L(w_i)$  by  $\delta_i$  to the right, we get another planar straight-line grid drawing.

- $\blacksquare$  induction on i, i.e. we consider  $G_3, \ldots, G_n$
- $\blacksquare$  clearly true for  $G_3$ ; assume the lemma holds for  $G_{k-1}$
- $\blacksquare$  let  $w_1, \ldots, w_p, v_k, w_q, \ldots, w_t$  be the boundary of  $G_k$



**Lemma:** Let  $G_k$  be a planar straight-line grid drawing. Let  $0 < \delta_1 \le \delta_2 \le \cdots \le \delta_t \in \mathbb{N}$ . If we shift each  $L(w_i)$  by  $\delta_i$  to the right, we get another planar straight-line grid drawing.

- $\blacksquare$  induction on i, i.e. we consider  $G_3,\ldots,G_n$
- $\blacksquare$  clearly true for  $G_3$ ; assume the lemma holds for  $G_{k-1}$
- $\blacksquare$  let  $w_1, \ldots, w_p, v_k, w_q, \ldots, w_t$  be the boundary of  $G_k$
- $\blacksquare \text{ let } \delta_1 \leq \cdots \leq \delta_p \leq \delta \leq \delta_q \leq \cdots \leq \delta_t$



**Lemma:** Let  $G_k$  be a planar straight-line grid drawing. Let  $0 < \delta_1 \le \delta_2 \le \cdots \le \delta_t \in \mathbb{N}$ . If we shift each  $L(w_i)$  by  $\delta_i$  to the right, we get another planar straight-line grid drawing.

- $\blacksquare$  induction on i, i.e. we consider  $G_3,\ldots,G_n$
- $\blacksquare$  clearly true for  $G_3$ ; assume the lemma holds for  $G_{k-1}$
- $\blacksquare$  let  $w_1, \ldots, w_p, v_k, w_q, \ldots, w_t$  be the boundary of  $G_k$
- $\blacksquare \text{ let } \delta_1 \leq \cdots \leq \delta_p \leq \delta \leq \delta_q \leq \cdots \leq \delta_t$




**Lemma:** Let  $G_k$  be a planar straight-line grid drawing. Let  $0 < \delta_1 \le \delta_2 \le \cdots \le \delta_t \in \mathbb{N}$ . If we shift each  $L(w_i)$  by  $\delta_i$  to the right, we get another planar straight-line grid drawing.

- $\blacksquare$  induction on i, i.e. we consider  $G_3, \ldots, G_n$
- $\blacksquare$  clearly true for  $G_3$ ; assume the lemma holds for  $G_{k-1}$
- $\blacksquare$  let  $w_1, \ldots, w_p, v_k, w_q, \ldots, w_t$  be the boundary of  $G_k$
- $\blacksquare \text{ let } \delta_1 \leq \cdots \leq \delta_p \leq \delta \leq \delta_q \leq \cdots \leq \delta_t$
- $\blacksquare \text{ define } \delta_i' = \begin{cases} \delta_i & \text{for } 1 \leq i \leq p \\ \delta + 1 & \text{for } p + 1 \leq i \leq q 1 \text{ (covered vertices of } v_k) \\ \delta_i + 2 & \text{for } q \leq i \leq t \end{cases}$
- by induction hypothesis we can move each  $L(w_1),\ldots,L(w_t)$  by  $\delta'_1,\ldots,\delta'_t$ , respectively



**Lemma:** Let  $G_k$  be a planar straight-line grid drawing. Let  $0 < \delta_1 \le \delta_2 \le \cdots \le \delta_t \in \mathbb{N}$ . If we shift each  $L(w_i)$  by  $\delta_i$  to the right, we get another planar straight-line grid drawing.

- $\blacksquare$  induction on i, i.e. we consider  $G_3, \ldots, G_n$
- $\blacksquare$  clearly true for  $G_3$ ; assume the lemma holds for  $G_{k-1}$
- $\blacksquare$  let  $w_1, \ldots, w_p, v_k, w_q, \ldots, w_t$  be the boundary of  $G_k$
- $\blacksquare \text{ let } \delta_1 \leq \cdots \leq \delta_p \leq \delta \leq \delta_q \leq \cdots \leq \delta_t$



- by induction hypothesis we can move each  $L(w_1),\ldots,L(w_t)$  by  $\delta'_1,\ldots,\delta'_t$ , respectively
- complete the drawing of  $G_k$  by placing  $v_k$ ; now  $v_k$  is moved rigidly with  $L(w_{p+1}),\ldots,L(w_{q-1})$  by  $\delta$

Overview



Canonical ordering

Shift algorithm

#### Implementation

# Naïve Implementation



```
Shift(plane graph G = (V, E))
 let v_1, \ldots, v_n canonical ordering of G
 for i = 1 to n do L(v_i) \leftarrow \{v_i\}
 P(v_1) \leftarrow (0,0); P(v_2) \leftarrow (2,0); P(v_3) \leftarrow (1,1)
 for k=4 to n do
   let w_1 = v_1, w_2, \dots, w_{t-1}, w_t = v_2 be vertices of C_{k-1}
   let w_p, \ldots, w_q be neighbors of v_k
   for v \in \bigcup_{i=p+1}^{q-1} L(w_i) do x(v) \leftarrow x(v) + 1
   for v \in \bigcup_{i=a}^t L(w_i) do x(v) \leftarrow x(v) + 2
   P(v_k) \leftarrow \text{intersection point of lines with slope } \pm 1 \text{ from } P(w_p) \text{ and } P(w_q)
   L(v_k) \leftarrow \cup_{j=p+1}^{q-1} L(w_j) \cup \{v_{\mathbf{z}}\}
```

# Naïve Implementation



```
Shift(plane graph G = (V, E))
 let v_1, \ldots, v_n canonical ordering of G
 for i = 1 to n do L(v_i) \leftarrow \{v_i\}
 P(v_1) \leftarrow (0,0); P(v_2) \leftarrow (2,0); P(v_3) \leftarrow (1,1)
 for k=4 to n do
    let w_1 = v_1, w_2, \dots, w_{t-1}, w_t = v_2 be vertices of C_{k-1}
    let w_p, \ldots, w_q be neighbors of v_k
   \begin{array}{l} \text{for } v \in \cup_{j=p+1}^{q-1} L(w_j) \text{ do } x(v) \leftarrow x(v)+1 \\ \text{for } v \in \cup_{j=q}^t L(w_j) \text{ do } x(v) \leftarrow x(v)+2 \end{array} \right\} \quad \text{(a) fine } 
    P(v_k) \leftarrow \text{intersection point of lines with slope} \pm 1 \text{ from } P(w_p) \text{ and } P(w_q)
   L(v_k) \leftarrow \cup_{i=p+1}^{q-1} L(w_i) \cup \{v_i\}
```


#### Running time?

$$\rightarrow O(n^2)$$

# Towards a Faster Implementation



#### Vertex coordinates



$$(x_{k}) = \frac{1}{2}(x(w_{q}) + x(w_{p}) + y(w_{q}) - y(w_{p}))$$

$$y(v_{k}) = \frac{1}{2}(x(w_{q}) - x(w_{p}) + y(w_{q}) + y(w_{p}))$$

$$x(v_{k}) - x(w_{p}) = \underbrace{\frac{1}{2}(x(w_{q}) - x(w_{p}) + y(w_{q}) - y(w_{p}))}_{\times -distance}$$


$$(x_{k}) - x(w_{p}) = \underbrace{\frac{1}{2}(x(w_{q}) - x(w_{p}) + y(w_{q}) - y(w_{p}))}_{\times -distance}$$

$$y = y(\omega_{p}) + 4 \cdot (x - x(\omega_{p}))$$


$$(x_{k}) - x(w_{k}) = \underbrace{\frac{1}{2}(x(w_{q}) - x(w_{p}) + y(w_{q}) - y(w_{p}))}_{\times -distance}$$


$$y = y(\omega_{p}) + 4 \cdot (x - x(\omega_{p}))$$


























lacktriangle in x-distance tree T at each vertex we keep its relative x-distance from its parent and its y-coordinate





relative x-distance tree T




- lacktriangle in x-distance tree T at each vertex we keep its relative x-distance from its parent and its y-coordinate
- If we know the y-coordinates of  $w_p$  and  $w_q$  and the difference  $x(w_p) x(w_q)$ , we can compute the difference  $x(v_k) x(w_p)$  [eq. (3)]



- lacktriangle in x-distance tree T at each vertex we keep its relative x-distance from its parent and its y-coordinate
- If we know the y-coordinates of  $w_p$  and  $w_q$  and the difference  $x(w_p)-x(w_q)$ , we can compute the difference  $x(v_k)-x(w_p)$  [eq. (3)]




- lacktriangle in x-distance tree T at each vertex we keep its relative x-distance from its parent and its y-coordinate
- If we know the y-coordinates of  $w_p$  and  $w_q$  and the difference  $x(w_p) x(w_q)$ , we can compute the difference  $x(v_k) x(w_p)$  [eq. (3)]






- lacktriangle in x-distance tree T at each vertex we keep its relative x-distance from its parent and its y-coordinate
- If we know the y-coordinates of  $w_p$  and  $w_q$  and the difference  $x(w_p) x(w_q)$ , we can compute the difference  $x(v_k) x(w_p)$  [eq. (3)]
- lacksquare  $\Delta_x(w_p,w_q)=\Delta_x(w_{p+1})+\cdots+\Delta_x(w_q)$ , where  $\Delta_x(w_q)$  is x-distance from the parent and  $\Delta_x(w_p,w_q)$  is x-distance of  $w_p$  and  $w_q$
- lacksquare Calculate  $\Delta_x(v_k)$  by eq. (3)
- lacksquare Calculate  $y(v_k)$  by eq. (2)






- lacktriangle in x-distance tree T at each vertex we keep its relative x-distance from its parent and its y-coordinate
- If we know the y-coordinates of  $w_p$  and  $w_q$  and the difference  $x(w_p) x(w_q)$ , we can compute the difference  $x(v_k) x(w_p)$  [eq. (3)]
- lacksquare  $\Delta_x(w_p,w_q)=\Delta_x(w_{p+1})+\cdots+\Delta_x(w_q)$ , where  $\Delta_x(w_q)$  is x-distance from the parent and  $\Delta_x(w_p,w_q)$  is x-distance of  $w_p$  and  $w_q$
- lacksquare Calculate  $\Delta_x(v_k)$  by eq. (3)
- lacksquare Calculate  $y(v_k)$  by eq. (2)






- lacktriangle in x-distance tree T at each vertex we keep its relative x-distance from its parent and its y-coordinate
- If we know the y-coordinates of  $w_p$  and  $w_q$  and the difference  $x(w_p) x(w_q)$ , we can compute the difference  $x(v_k) x(w_p)$  [eq. (3)]
- lacksquare  $\Delta_x(w_p,w_q)=\Delta_x(w_{p+1})+\cdots+\Delta_x(w_q)$ , where  $\Delta_x(w_q)$  is x-distance from the parent and  $\Delta_x(w_p,w_q)$  is x-distance of  $w_p$  and  $w_q$
- lacksquare Calculate  $\Delta_x(v_k)$  by eq. (3)
- lacksquare Calculate  $y(v_k)$  by eq. (2)






- lacktriangle in x-distance tree T at each vertex we keep its relative x-distance from its parent and its y-coordinate
- If we know the y-coordinates of  $w_p$  and  $w_q$  and the difference  $x(w_p) x(w_q)$ , we can compute the difference  $x(v_k) x(w_p)$  [eq. (3)]
- lacksquare  $\Delta_x(w_p,w_q)=\Delta_x(w_{p+1})+\cdots+\Delta_x(w_q)$ , where  $\Delta_x(w_q)$  is x-distance from the parent and  $\Delta_x(w_p,w_q)$  is x-distance of  $w_p$  and  $w_q$
- lacksquare Calculate  $\Delta_x(v_k)$  by eq. (3)
- lacksquare Calculate  $y(v_k)$  by eq. (2)






- lacktriangle in x-distance tree T at each vertex we keep its relative x-distance from its parent and its y-coordinate
- If we know the y-coordinates of  $w_p$  and  $w_q$  and the difference  $x(w_p) x(w_q)$ , we can compute the difference  $x(v_k) x(w_p)$  [eq. (3)]
- lacksquare  $\Delta_x(w_p,w_q)=\Delta_x(w_{p+1})+\cdots+\Delta_x(w_q)$ , where  $\Delta_x(w_q)$  is x-distance from the parent and  $\Delta_x(w_p,w_q)$  is x-distance of  $w_p$  and  $w_q$
- lacksquare Calculate  $\Delta_x(v_k)$  by eq. (3)
- lacksquare Calculate  $y(v_k)$  by eq. (2)






- lacktriangle in x-distance tree T at each vertex we keep its relative x-distance from its parent and its y-coordinate
- If we know the y-coordinates of  $w_p$  and  $w_q$  and the difference  $x(w_p) x(w_q)$ , we can compute the difference  $x(v_k) x(w_p)$  [eq. (3)]
- lacksquare  $\Delta_x(w_p,w_q)=\Delta_x(w_{p+1})+\cdots+\Delta_x(w_q)$ , where  $\Delta_x(w_q)$  is x-distance from the parent and  $\Delta_x(w_p,w_q)$  is x-distance of  $w_p$  and  $w_q$
- lacksquare Calculate  $\Delta_x(v_k)$  by eq. (3)
- lacksquare Calculate  $y(v_k)$  by eq. (2)





- lacktriangle in x-distance tree T at each vertex we keep its relative x-distance from its parent and its y-coordinate
- If we know the y-coordinates of  $w_p$  and  $w_q$  and the difference  $x(w_p) x(w_q)$ , we can compute the difference  $x(v_k) x(w_p)$  [eq. (3)]
- lacksquare  $\Delta_x(w_p,w_q)=\Delta_x(w_{p+1})+\cdots+\Delta_x(w_q)$ , where  $\Delta_x(w_q)$  is x-distance from the parent and  $\Delta_x(w_p,w_q)$  is x-distance of  $w_p$  and  $w_q$
- lacksquare Calculate  $\Delta_x(v_k)$  by eq. (3)
- lacksquare Calculate  $y(v_k)$  by eq. (2)

- $\Delta_x(w_{p+1}) = \Delta_x(w_{p+1}) \Delta_x(v_k)$






- lacktriangle in x-distance tree T at each vertex we keep its relative x-distance from its parent and its y-coordinate
- If we know the y-coordinates of  $w_p$  and  $w_q$  and the difference  $x(w_p) x(w_q)$ , we can compute the difference  $x(v_k) x(w_p)$  [eq. (3)]
- lacksquare  $\Delta_x(w_p,w_q)=\Delta_x(w_{p+1})+\cdots+\Delta_x(w_q)$ , where  $\Delta_x(w_q)$  is x-distance from the parent and  $\Delta_x(w_p,w_q)$  is x-distance of  $w_p$  and  $w_q$
- lacksquare Calculate  $\Delta_x(v_k)$  by eq. (3)
- lacksquare Calculate  $y(v_k)$  by eq. (2)

$$\Delta_x(w_{p+1}) = \Delta_x(w_{p+1}) - \Delta_x(v_k)$$

Once T is completed, compute x-coordinates by a linear-time pre-order traversal



# Summary



**Theorem:** Every n-vertex embedded planar graph G=(V,E) has a straight-line planar drawing on a grid of size  $(2n-4)\times(n-2)$ . [de Fraysseix, Pach, Pollack 1988]

**Theorem:** The corresponding shift algorithm can be implemented to run in O(n) time. [Chrobak, Payne 1995]