

Thermal Correction of VIR Hyperspectral Data for Ceres: Bayesian Approach for simultaneous Parameter Estimation

Aliyya Fathima Rinu¹, Guneshwar Thangjam²

^{1,2}National Institute of Science Education and Research, Bhubaneswar, India

¹aliyyarinu@gmail.com

Abstract

We present a method to simultaneously estimate surface temperature and spectral emissivity from hyperspectral radiance measurements obtained by the VIR instrument onboard NASA's *Dawn* mission during the HAMO phase at Ceres. The dataset spans the visible–infrared region (0.25–5.1 μm) with a spatial resolution of ~ 0.38 km/pixel. Estimating surface temperature from radiance is an underdetermined problem, as it requires determining $N+1$ unknowns from N equations. Previous approaches often imposed fixed emissivity assumptions to retrieve temperature first. Here, we employ a Bayesian framework that allows simultaneous estimation of temperature and emissivity without prior fixed values. Our two-step strategy begins with an uninformative state of knowledge [?] to derive Gaussian priors for the subsequent iterative retrieval using a theory-based numerical method. The final temperature maps deviate by 1–3 K from the first-step estimates, with emissivity uncertainties on the order of 10^{-3} . For Haulani crater, the retrieved temperatures show lower values compared to [?].