# Math 426.2SY Calculus II

University of New Hampshire

May 31, 2017

1 / 16

(UNH) Lecture 4 May 31, 2017

### Outline

1 5.6 - Substitution and Area Between Curves

(UNH) Lecture 4 May 31, 2017 2 / 16

## 5.6 - Substitution and Definite Integrals

There are two methods to evaluate definite integrals using the substitution method.

#### Example of Method 1

$$\int_{-1}^{1} 3x^2 \sqrt[3]{x^3 + 1} \, dx$$

(UNH) Lecture 4 May 31, 2017 3 / 16

# Substitution and Definite Integrals

### Example of Method 2

$$\int_{-1}^{1} 3x^2 \sqrt[3]{x^3 + 1} \, dx$$

ロト 4月ト 4 三 ト 三 ・ の9个

(UNH) Lecture 4 May 31, 2017 4 / 16

## Substitution and Definite Integrals

#### More Examples of Method 2

$$\int_0^{\pi/12} \tan(3x) \sec^2(3x) dx$$

ロト 4月ト 4 三 ト 三 ・ の9个

(UNH) Lecture 4 May 31, 2017 5 / 16

# Substitution and Definite Integrals

### More Examples of Method 2

$$\int_{-\pi/4}^{\pi/4} \tan x \, dx$$

ロト 4月 4 三 ト 4 三 ト 日 めのや

(UNH) Lecture 4 May 31, 2017 6 / 16

# Definite Integrals of Symmetric Functions

#### Definition

- A function f(x) is called **even** if f(-x) = f(x) for all x.
- A function f(x) is called **odd** if f(-x) = -f(x) for all x.
- Examples:

4□ > 4□ > 4□ > 4□ > 4□ > 900

(UNH) Lecture 4 May 31, 2017 7 / 16

# Definite Integrals of Symmetric Functions

#### Theorem

Let f be a function defined on the interval [-a, a].

- If f is odd, then  $\int_{-a}^{a} f(x) dx = 0$
- ② If f is even, then  $\int_{-a}^{a} f(x) dx = 2 \int_{0}^{a} f(x) dx$ 
  - Proof:

ロ ト 4 個 ト 4 重 ト 4 重 ト 9 9 (や)

(UNH) Lecture 4 May 31, 2017 8 / 16

## Definite Integrals of Symmetric Functions

#### Examples

- $\int_{-\pi/3}^{\pi/3} e^{x^2} \sin x \, dx$   $\int_{-2}^{2} x^4 4x^2 + 6 \, dx$

(UNH) May 31, 2017 9 / 16 Lecture 4

• Suppose we want to find the area of a region that is bounded above by the curve y = f(x), below by the curve y = g(x), and on the left and right by the lines x = a and x = b.

4D > 4A > 4B > 4B > B 900

(UNH) Lecture 4 May 31, 2017 10 / 16

If  $f(x) \ge g(x)$  throughout [a, b], then the area of the region between the curves y = f(x) and y = g(x) from a to b is given by:

$$A = \int_{a}^{b} f(x) - g(x) \, dx$$

シャイ ほう イモト イロト

(UNH) Lecture 4 May 31, 2017 11 / 1

### Example

Find the area of the region enclosed by the parabola  $y = 2-x^2$  and the line y = -x.

(UNH) Lecture 4 May 31, 2017 12 / 16

#### Note

If the formula for a bounding curve changes at one or more points, we subdivide the region into subregions that correspond to the formula changes and apply the formula for the area between curves to each subregion.

ロト 4月ト 4 三 ト 1 三 り90

(UNH) Lecture 4 May 31, 2017 13 / 16

#### Example

Find the area of the region in the first quadrant that is bounded above by  $y = \sqrt{x}$  and below by the x-axis and the line y = x-2.

(UNH) Lecture 4 May 31, 2017 14 / 16

#### Example

Find the area of the region enclosed by y = 4x + 3 and  $y = 3 - x^2$ .

4 D > 4 A > 4 E > 4 E > 9 Q A

(UNH) Lecture 4 May 31, 2017 15 / 16

#### Example

Find the area of the region enclosed by  $x + y^2 = 0$  and  $x + 3y^2 = 2$ .

(UNH) Lecture 4 May 31, 2017 16 / 16