Math 426.2SY Calculus II

University of New Hampshire

June 5, 2017

(UNH) Lecture 5 June 5, 2017 1 / 12

Outline

1 6.2 - Method of Cylindrical Shells

2 6.3 - Arc Lenght



(UNH) Lecture 5 June 5, 2017 2 / 12

Introduction

Finding the volume using the Method of Disks/Washers

- Look at a slice of the 2D region **perpendicular** to the axis of revolution.
- Find the volume of this slice after rotation. (Call it ΔV_k)
- Approximate the total volume by $V \approx \sum_{k=1}^{n} \Delta V_k$
- Replace the \sum by \int
- Find the limits of integration by looking at the region.
- Evaluate the definite integral.

Today

- The first step in this procedure (taking slices perpendicular to the axis of rotation) is not always efficient.
- Sometimes its easier to take slice that are **Parallel** to the axis of rotation.
- The rest of the procedure remains unchanged.
- This new method is called the the **Method of Cylindrical Shells**.

Lecture 5 June 5, 2017

Example

The region enclosed by the x-axis and the parabola $y = f(x) = 3x - x^2$ is revolved about the vertical line x = -1 to generate a solid. Find the volume of the solid.

JNH) Lecture 5 June 5, 2017 4 / 12

• Using the Method of Washers requires us to solve $y = f(x) = 3x - x^2$ for x, which leads to complicated formulas.

(UNH) Lecture 5 June 5, 2017 5 / 12

• Instead, let's look at slices of the 2D region **parallel** to the axis of revolution.

(UNH) Lecture 5 June 5, 2017 6 / 12

Example

The region bounded by the curve $y = \sqrt{x}$ the x-axis, and the line x = 4 is revolved about the y-axis to generate a solid. Find the volume of the solid.

(UNH) Lecture 5 June 5, 2017 7 / 12

Example

The region bounded by the curve $x=3-y^2$ and the lines $y=\sqrt{3}, x=3$ is revolved about the x-axis to generate a solid. Find the volume of the solid.

(UNH) Lecture 5 June 5, 2017 8 / 12

The Washer Method


Example

The region bounded by the curve $x=3-y^2$ and the lines $y=\sqrt{3}, x=3$ is revolved about the x-axis to generate a solid. Find the volume of the solid.

(UNH) Lecture 5 June 5, 2017 9 / 12

6.3 - Arc Lenght

- We know what is meant by the length of a straight line segment.
- Using Calculus, we can find a precise denfition for the length of a general curve.



Arc Lenght

Example

Find the length of $y = \ln x - x^2/8$ from x = 1 to x = 2.



(UNH) Lecture 5 June 5, 2017 11 / 12

Arc Lenght

Example

Find the length of $y = \int_{-2}^{x} \sqrt{3t^4 - 1} dt$ from x = -2 to x = -1.



(UNH) Lecture 5 June 5, 2017 12 / 12